

Climate Change Vulnerability Assessment
Washington Natural Heritage Program
June 27, 2014

Species	Artemisia campestris var. wormskioldii	Index Result:	Moderately Vulnerable
English Name	northern wormwood	Confidence:	Very High
Taxonomic Group	Vascular Plants	Assessor:	John Gamon
Geographic Area	Washington		
Range Rel.	Entire range		
Cave/Ground Water Obligate	No		
GRank	G5T1		
SRank	S1		

Climate Change Vulnerability Index Values:

Temperature Scope	A >5.5F	0
	A 5.1F	0
	A 4.5F	0
	A 3.9F	100
	A <3.9F	0
Hamon AET:PET Moisture Metric Scope	< -0.119	0
	-0.119	0
	-0.096	0
	-0.073	0
	-0.05	100
	>-0.028	0
Sea level rise	B1	N
Natural barriers	B2a	N
Anthropogenic barriers	B2b	Inc
Climate Change mitigation	B3	SI
Dispersal/Movement	C1	SI
Historical thermal niche	C2ai	N
Physiological thermal niche	C2aii	N
Historical hydrological niche	C2bi	SI
Physiol. hydrological niche	C2bii	N
Disturbance dependence	C2c	SI
Ice/snow dependence	C2d	N
Physical habitat restrictions	C3	N
Other spp create habitat	C4a	N
Dietary Versatility	C4b	N/A
Pollinator Versatility	C4c	U
Other spp for dispersal	C4d	N
Other spp interaction	C4e	N
Genetic variation	C5a	U
Genetic bottleneck	C5b	U
Phenological response	C6	U
Documented response	D1	U
Modeled change	D2	U
Modeled overlap	D3	U
Modeled protected Areas	D4	U

Affect on Vulnerability:
GI = Greatly Increase
Inc = Increase
SI = Somewhat Increase
N = Neutral
SD = Somewhat Decrease
Dec = Decrease
U = Unknown

Index Scores:

Extremely Vulnerable:

Abundance and/or range extent within geographical area assessed extremely likely to substantially decrease or disappear by 2050.

Highly Vulnerable: Abundance and/or range extent within geographical area assessed likely to decrease significantly by 2050.

Moderately Vulnerable:

Abundance and/or range extent within geographical area assessed likely to decrease by 2050.

Not Vulnerable/Presumed

Stable: Available evidence does not suggest that abundance and/or range extent within the geographical area assessed will change (increase/decrease) substantially by 2050. Actual range boundaries may change.

Not Vulnerable/Increase Likely:

Available evidence suggests that abundance and/or range extent within geographical area assessed is likely to increase by 2050.




Figure 1. Exposure of the known occurrences of *Artemisia campestris* var. *wormskioldii* to projected local climate change: temperature. Both occurrences are projected to experience a temperature increase of 3.9 – 4.4° F by the mid 2050s.




Figure 2. Exposure of known occurrences of *Artemisia campestris* var. *wormskioldii* to projected local climate change: predicted annual change in Hamon AET:PET moisture metric, 2040-2069. Values represent percent change in the metric; negative values indicate net drying. Both occurrences are projected to experience a net drying in the range of -0.028 to -0.050.




Figure 3. Historical thermal niche (exposure to past variations in temperature) of the Washington occurrences of *Artemisia campestris* var. *wormskioldii*. This factor measures large-scale temperature variation that a species has experienced in recent historical times (i.e., the past 50 years), as approximated by mean seasonal temperature variation (difference between highest mean monthly maximum temperature and lowest mean monthly minimum temperature) for occupied cells within the assessment area. It is a proxy for species' temperature tolerance at a broad scale. Considering the mean seasonal temperature variation for occupied cells, the species has experienced **average (57.1 - 77° F/31.8 - 43.0° C)** temperature variation in the past 50 years. This range of variation is characterized as being **Neutral** to climate change.




Figure 4. Historical hydrological niche (exposure to past variations in precipitation) of the Washington occurrences of *Artemisia campestris var. wormskioldii*. This factor measures large-scale precipitation variation that a species has experienced in recent historical times (i.e., the past 50 years), as approximated by mean annual precipitation variation across occupied cells within the assessment area. Considering the range of mean annual precipitation across occupied cells, the species has experienced **slightly lower than average (11 - 20 inches/255 - 508 mm)** precipitation variation in the past 50 years. This range of precipitation variation is characterized as indicating a **Somewhat Increased Vulnerability** to climate change.