



---

# Chapter 3

## Affected Environment

---

- 3.1 Introduction
- 3.2 Land Ownership and Use
- 3.3 Air Quality
- 3.4 Geology, Soils, and Erosional Processes
- 3.5 Water Resources
- 3.6 Vegetation
- 3.7 Riparian and Wetland Processes
- 3.8 Fish and Fish Habitat
- 3.9 Amphibians and Amphibian Habitat
- 3.10 Birds, Mammals, Other Wildlife, and Their Habitats
- 3.11 Recreation
- 3.12 Visual Resources
- 3.13 Archeological, Historical, Cultural and Indian Trust Resources
- 3.14 Social and Economic Environment





3. AFFECTED ENVIRONMENT

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43

3.	AFFECTED ENVIRONMENT .....	3-1
3.1	INTRODUCTION .....	3-2
3.2	LAND OWNERSHIP AND USE .....	3-5
3.2.1	Introduction .....	3-5
3.2.2	Existing Habitat Conservation Plans.....	3-9
3.2.3	Land Ownership and Use by Region .....	3-9
3.2.4	Timber Harvest Rates for Western and Eastern Washington.....	3-14
3.2.5	Forestland Conversion .....	3-16
3.3	AIR QUALITY .....	3-23
3.4	GEOLOGY, SOILS, AND EROSIONAL PROCESSES.....	3-25
3.4.1	Geology and Soils.....	3-25
3.4.2	Erosion .....	3-28
3.5	WATER RESOURCES .....	3-37
3.5.1	Surface Water Quality.....	3-37
3.5.2	Surface Water Quantity.....	3-43
3.5.3	Groundwater Quantity and Quality .....	3-50
3.6	VEGETATION.....	3-53
3.6.1	Forest Vegetation.....	3-53
3.6.2	Riparian Vegetation .....	3-58
3.6.3	Disturbance Agents.....	3-59
3.6.4	Threatened and Endangered Plants.....	3-63
3.6.5	Invasive Plants.....	3-63
3.7	RIPARIAN AND WETLAND PROCESSES .....	3-65
3.7.1	Riparian Areas .....	3-65
3.7.2	Wetlands .....	3-71
3.8	FISH AND FISH HABITAT.....	3-77
3.8.1	Introduction .....	3-77
3.8.2	Fish Status in Washington .....	3-77
3.8.3	Life History of Covered and Affected Fish Species .....	3-78
3.8.4	The Freshwater Aquatic Ecosystem .....	3-94
3.8.5	Fish and Fish Habitat by Analysis Region .....	3-102
3.9	AMPHIBIANS AND AMPHIBIAN HABITAT .....	3-111
3.9.1	Introduction .....	3-111
3.9.2	Amphibian Distribution, Status, and Habitat .....	3-112
3.9.3	Review of Timber Harvest Effects on Amphibians.....	3-118
3.10	BIRDS, MAMMALS, OTHER WILDLIFE AND THEIR HABITATS .....	3-123
3.10.1	Introduction .....	3-123
3.10.2	Federally Threatened or Endangered Wildlife Species .....	3-125
3.10.3	Importance of Riparian Habitats to Wildlife.....	3-132
3.10.4	Wildlife in Upland Forested Habitats.....	3-134



# Chapter 3

1	3.11 RECREATION.....	3-137
2	3.11.1 Introduction.....	3-137
3	3.11.2 Estimates of Future Recreational Needs.....	3-138
4	3.11.3 Private Lands and Recreation.....	3-139
5	3.11.4 Forest Road-related Recreation.....	3-139
6	3.12 VISUAL RESOURCES.....	3-141
7	3.12.1 Introduction.....	3-141
8	3.12.2 Visual Resources and the Current Washington Forest Practices Rules ..	3-143
9	3.13 ARCHEOLOGICAL, HISTORICAL, CULTURAL AND INDIAN TRUST	
10	RESOURCES.....	3-145
11	3.13.1 Introduction.....	3-145
12	3.13.2 Archaeological Overview.....	3-145
13	3.13.3 Cultural and Trust Resources of Native American Tribes.....	3-148
14	3.13.4 Overview of Regional History.....	3-150
15	3.13.5 Washington State Protective Structure.....	3-151
16	3.14 SOCIAL AND ECONOMIC ENVIRONMENT.....	3-153
17	3.14.1 Introduction.....	3-153
18	3.14.2 Population.....	3-153
19	3.14.3 Employment and the Economy.....	3-154
20	3.14.4 Environmental Justice.....	3-161

21

## 3.1 INTRODUCTION

22  
23 This chapter describes the affected environment to provide background for the assessment  
24 of the environmental effects of the alternatives in Chapter 4 (Environmental Effects) and  
25 Chapter 5 (Cumulative Effects). The affected environment sections describe the resources  
26 and their current conditions against which the anticipated environmental effects of the  
27 alternatives described in Chapter 2 (Alternatives) are evaluated. The first section describes  
28 land ownership and use within the State, to provide context for the description of the other  
29 sections. The remaining sections present the physical environment first, followed by the  
30 biological environment, and then the social environment. The specific order of the sections  
31 is as follows:

- 32 • Land Ownership and Use (subsection 3.2)
- 33 • Air Quality (subsection 3.3)
- 34 • Geology, Soils, and Erosional Processes (subsection 3.4)
- 35 • Water Resources (subsection 3.5)
- 36 • Vegetation (subsection 3.6)
- 37 • Riparian and Wetland Processes (subsection 3.7)
- 38 • Fish and Fish Habitat (subsection 3.8)
- 39 • Amphibian and Amphibian Habitat (subsection 3.9)
- 40 • Birds, Mammals, Other Wildlife and Their Habitats (subsection 3.10)
- 41 • Recreation (subsection 3.11)
- 42 • Visual Resources (subsection 3.12)
- 43 • Cultural Resources and Indian Trust Resources (subsection 3.13)
- 44 • Socioeconomic Conditions (subsection 3.14)



1 The study area that defines the affected environment includes the majority of the State of  
2 Washington. The proposed action and the alternatives would directly affect the forested  
3 lands that are covered by the Washington Forest Practices Rules. These lands include the  
4 non-Federal and non-tribal forestlands of the State (Figure 3-1). These lands are referred  
5 to as the “covered lands” or the lands subject to Washington Forest Practices Rules in this  
6 EIS (See also the SEPA *Final EIS on Alternatives for Forest Practices Rules for: Aquatic*  
7 *and Riparian Resources* dated April 2001, Washington Forest Practices Board).

8 In addition to displaying the covered lands, Figure 3-1 displays 12 analysis regions, which  
9 are similar to the 10 regions identified in the Forest Practices Alternatives SEPA EIS Rules  
10 for Aquatic and Riparian Resources (Washington Forest Practices Board 2001c, 2002).  
11 However, to more fully capture the diverse landscape of the Puget Sound Region, this  
12 Region was divided into three smaller regions in this document. Detailed maps of each  
13 analysis region that illustrate rivers, lakes, highways, and more local place names are  
14 provided in the Regional Summaries (DEIS Appendix A).

15 The 12 analysis regions are referenced in this EIS to describe some of the regional aspects  
16 of the affected environment. This information is used in Chapters 4 and 5 to assess the  
17 indirect effects of the alternatives described in Chapter 2 (Alternatives). The regions were  
18 defined based on three factors: the distribution of threatened and endangered salmonids,  
19 Water Resource Inventory Area (WRIA) boundaries, and the physiographic regions of the  
20 State. The 12 analysis regions consist of 7 western Washington regions and 5 in eastern  
21 Washington as follows:

### 22 Western Washington Analysis Regions

- 23 • North Puget Sound
- 24 • South Puget Sound
- 25 • West Puget Sound
- 26 • Islands
- 27 • Olympic Coast
- 28 • Southwest
- 29 • Lower Columbia

### 30 Eastern Washington Analysis Regions

- 31 • Middle Columbia
- 32 • Upper Columbia – Downstream of Grand Coulee Dam
- 33 • Upper Columbia – Upstream of Grand Coulee Dam
- 34 • Snake River
- 35 • Columbia Basin


36 To provide further background and detail for the affected environment descriptions and the  
37 evaluation of effects, detailed summaries of land ownership and use and physical and  
38 biological factors were developed for each of the analysis regions. These descriptions are  
39 provided in DEIS Appendix A.

40



# Chapter 3

Figure 3-1. Analysis Regions and Covered Lands in Washington



3-4



## 1   **3.2 LAND OWNERSHIP AND USE**

### 2   **3.2.1 Introduction**

3   The State of Washington is approximately 43,272,000 acres in size. Federal lands  
4   comprise approximately 30 percent of the State, with slightly more than one-third of these  
5   lands (11 percent of the State) classified as wilderness, national parks, or wildlife refuges.  
6   State and tribal lands comprise approximately 9 percent and 7 percent of the State,  
7   respectively, with county and city lands accounting for approximately 1 percent. The  
8   remaining 53 percent of the lands in Washington are in private ownership (Table 3-1).

9   Slightly more than half of Washington State (53 percent) is forested (Table 3-2).  
10   Forestland accounts for 83 percent of western Washington and just 36 percent of eastern  
11   Washington. Eastern Washington is, however, considerably larger than western  
12   Washington, accounting for 64 percent of the State. Approximately 9 million acres in  
13   eastern Washington are forested, compared to 13 million acres in western Washington.  
14   Shrubland and grassland comprise approximately 23 percent of the State, with the majority  
15   of these lands (97 percent) located in eastern Washington. Agricultural lands account for  
16   approximately 18 percent of the State. Freshwater and wetlands account for 2 percent of  
17   the State; ice, snow, and bare rock account for another 2 percent; with residential and  
18   commercial lands covering the remaining 2 percent (Table 3-2).

19   Approximately 28 percent of forestlands in the State are Federal and State lands not  
20   managed for timber production. Federal and tribal lands available for timber management  
21   comprise approximately 22 percent of the forestland in the State. The remaining 50  
22   percent are State, county, city, and private lands that are potentially available for timber  
23   management under the Washington Forest Practices Rules (Table 3-3). These State,  
24   county, city, and private lands account for approximately 26 percent of total State lands.  
25   State, county, city, and private lands potentially available for timber management under the  
26   Washington Forest Practices Rules account for approximately 62 percent of forestlands in  
27   western Washington and 34 percent in eastern Washington (Table 3-3).

28   Land ownership and use across Washington State is heavily affected by the distribution  
29   and size of the human population. Approximately 5.9 million people resided in  
30   Washington State in 2000, an increase of approximately 21 percent or one million people  
31   since 1990 (U.S. Census Bureau 2000). Population projections anticipate continued  
32   population growth in the State, with the total population projected to reach 7.5 million by  
33   2020 (Washington Office of Financial Management 2002a). As the population of the State  
34   continues to increase, land ownership and land use are affected, and development in the  
35   form of urban growth and low-density residential areas is likely to continue to encroach on  
36   the State's forestlands, farmlands, and fish and wildlife habitat. Population trends are  
37   discussed further in subsection 3.14 (Social and Economic Environment).

38   The remainder of this section is divided into four subsections that address existing Habitat  
39   Conservation Plans (HCPs), land ownership and use by region, timber harvest rates, and  
40   forestland conversion, respectively.



## Chapter 3

**Table 3-1.** Land Ownership Acreage in Washington State by Analysis Region.

Analysis Region	Federal Wildernesses, National Parks, and Wildlife Refuges	Other Federal Lands	State Parks and Wildlife Areas	Washington DNR and Other State Lands	County and City Lands	Tribal Lands	Private Lands	Total
<b>Western Washington</b>								
North Puget Sound	1,323,585	987,699	20,707	493,568	31,761	40,785	1,499,857	4,397,962
South Puget Sound	200,046	279,906	8,003	161,563	122,175	23,351	1,383,184	2,178,228
West Puget Sound	454,466	235,362	13,154	176,166	9,194	15,349	831,597	1,735,288
Islands	1,990	8,425	11,417	12,895	1,321	0	210,775	246,822
Olympic Coast	529,794	196,674	1,673	309,147	7,752	234,990	476,837	1,756,868
Southwest	12,132	124,872	12,582	304,062	40,062	4,623	1,814,921	2,313,254
Lower Columbia	327,355	750,238	14,033	325,013	2,512	95	1,653,166	3,072,412
<b>Western Washington Total</b>	<b>2,849,368</b>	<b>2,583,176</b>	<b>81,569</b>	<b>1,782,414</b>	<b>214,777</b>	<b>319,193</b>	<b>7,870,337</b>	<b>15,700,834</b>
<b>Percent of Western Washington Total</b>	<b>18%</b>	<b>16%</b>	<b>1%</b>	<b>11%</b>	<b>1%</b>	<b>2%</b>	<b>50%</b>	<b>100%</b>
<b>Eastern Washington</b>								
Middle Columbia	355,338	1,302,933	178,826	464,006	1,388	1,255,467	2,939,158	6,497,115
Upper Columbia - Downstream of Grand Coulee	1,203,796	2,043,164	183,062	573,642	1,237	431,539	1,964,137	6,400,577
Upper Columbia - Upstream of Grand Coulee	82,706	1,477,635	33,649	345,066	10,293	1,084,900	2,713,551	5,747,801
Snake River	125,263	338,433	44,592	231,230	795	0	3,835,556	4,575,868
Columbia Basin	34,358	353,942	65,990	254,332	214	0	3,641,362	4,350,198
<b>Eastern Washington Total</b>	<b>1,801,461</b>	<b>5,516,107</b>	<b>506,119</b>	<b>1,868,276</b>	<b>13,927</b>	<b>2,771,906</b>	<b>15,093,764</b>	<b>27,571,559</b>
<b>Percent of Eastern Washington Total</b>	<b>7%</b>	<b>20%</b>	<b>2%</b>	<b>7%</b>	<b>0%</b>	<b>10%</b>	<b>55%</b>	<b>100%</b>
<b>STATE TOTAL</b>	<b>4,650,830</b>	<b>8,099,284</b>	<b>587,687</b>	<b>3,650,689</b>	<b>228,705</b>	<b>3,091,098</b>	<b>22,964,102</b>	<b>43,272,394</b>
<b>State Total Percent</b>	<b>11%</b>	<b>19%</b>	<b>1%</b>	<b>8%</b>	<b>1%</b>	<b>7%</b>	<b>53%</b>	<b>100%</b>

Source: Washington DNR Major Public Lands and WRIA GIS layers 2004.



## Chapter 3



**Table 3-2.** Acreage of Washington State in General Land Cover/Use Categories by Analysis Region.

Region	Forestland	Shrubland	Grassland	Water and Wetlands	Ice, Snow, and Bare Rock	Residential and Commercial	Agricultural	Total
<b>Western Washington</b>								
North Puget Sound	3,427,441	105,972	132,537	75,138	269,681	83,594	303,607	4,397,968
South Puget Sound	1,532,465	32,194	29,988	59,762	47,315	362,055	114,443	2,178,221
West Puget Sound	1,522,350	19,320	22,272	23,033	31,877	70,114	46,333	1,735,299
Islands	180,345	3,959	1,822	4,136	1,285	14,020	41,242	246,808
Olympic Coast	1,671,082	11,358	6,887	35,546	25,431	2,015	4,542	1,756,860
Southwest	2,057,953	15,866	8,157	52,161	4,319	30,522	144,258	2,313,236
Lower Columbia	2,615,876	45,966	24,032	90,067	58,793	65,890	171,772	3,072,396
<b>Western Washington Total</b>	<b>13,007,512</b>	<b>234,635</b>	<b>225,695</b>	<b>339,843</b>	<b>438,701</b>	<b>628,210</b>	<b>826,197</b>	<b>15,700,788</b>
<b>Percent of Western Washington Total</b>	<b>83%</b>	<b>1%</b>	<b>1%</b>	<b>2%</b>	<b>3%</b>	<b>4%</b>	<b>5%</b>	<b>100%</b>
<b>Eastern Washington</b>								
Middle Columbia	2,691,433	1,828,024	620,496	92,678	52,851	76,321	1,135,336	6,497,138
Upper Columbia-Downstream of Grand Coulee	2,773,960	1,633,415	1,113,611	125,289	153,690	30,129	570,473	6,400,567
Upper Columbia-Upstream of Grand Coulee	4,084,093	360,939	386,504	131,226	3,355	81,352	700,332	5,747,801
Snake River	376,315	1,343,585	369,787	63,445	913	30,922	2,390,898	4,575,865
Columbia Basin	12,842	1,696,451	175,870	111,574	873	46,573	2,306,014	4,350,198
<b>Eastern Washington Total</b>	<b>9,938,643</b>	<b>6,862,414</b>	<b>2,666,268</b>	<b>524,212</b>	<b>211,682</b>	<b>265,297</b>	<b>7,103,053</b>	<b>27,571,569</b>
<b>Percent of Eastern Washington Total</b>	<b>36%</b>	<b>25%</b>	<b>10%</b>	<b>2%</b>	<b>1%</b>	<b>1%</b>	<b>26%</b>	<b>100%</b>
<b>STATE TOTAL</b>	<b>22,946,156</b>	<b>7,097,049</b>	<b>2,891,963</b>	<b>864,054</b>	<b>650,382</b>	<b>893,507</b>	<b>7,929,249</b>	<b>43,272,359</b>
<b>Percent of State Total</b>	<b>53%</b>	<b>16%</b>	<b>7%</b>	<b>2%</b>	<b>2%</b>	<b>2%</b>	<b>18%</b>	<b>100%</b>

Source: U.S. Geological Survey Land Use/Land Cover and WRIA GIS layers 2004.



## Chapter 3

**Table 3-3.** Ownership and Management of Forestlands in Washington State by Analysis Region.

Region	Federal and State Protected Lands Not Managed for Timber Production <sup>1/</sup>	Federal and Tribal Lands Available for Timber Management <sup>2/</sup>	Forestlands Available for Timber Management and Subject to Washington Forest Practices Rules			Total Forested Lands
			Washington DNR and Other State Lands <sup>3/</sup>	Private, County, and City Lands	Total	
<b>Western Washington</b>						
North Puget Sound	1,644,519	235,028	472,932	1,074,910	1,547,842	3,427,389
South Puget Sound	291,193	122,903	148,349	970,000	1,118,349	1,532,444
West Puget Sound	631,196	22,105	168,691	700,204	868,896	1,522,198
Islands	11,706	3,607	10,562	154,405	164,967	180,280
Olympic Coast	684,287	228,828	307,170	450,786	757,957	1,671,071
Southwest	140,690	3,933	294,684	1,618,539	1,913,223	2,057,847
Lower Columbia	719,253	262,482	313,523	1,320,459	1,633,982	2,615,716
<b>Western Washington Total</b>	<b>4,122,844</b>	<b>878,886</b>	<b>1,715,911</b>	<b>6,289,303</b>	<b>8,005,216</b>	<b>13,006,945</b>
<b>Percent of Western Washington Total</b>	<b>32%</b>	<b>7%</b>	<b>13%</b>	<b>48%</b>	<b>62%</b>	<b>100%</b>
<b>Eastern Washington</b>						
Middle Columbia	879,862	867,469	231,650	712,447	944,097	2,691,428
Upper Columbia-Downstream of Grand Coulee	1,267,217	1,034,605	214,305	257,835	472,140	2,773,963
Upper Columbia-Upstream of Grand Coulee	102,588	2,177,129	284,808	1,519,518	1,804,326	4,084,043
Snake River	95,725	149,067	12,791	118,730	131,522	376,314
Columbia Basin	36	120	1,481	11,205	12,687	12,843
<b>Eastern Washington Total</b>	<b>2,345,428</b>	<b>4,228,390</b>	<b>745,035</b>	<b>2,619,735</b>	<b>3,364,772</b>	<b>9,938,591</b>
<b>Percent of Eastern Washington Total</b>	<b>24%</b>	<b>43%</b>	<b>7%</b>	<b>26%</b>	<b>34%</b>	<b>100%</b>
<b>STATE TOTAL</b>	<b>6,468,273</b>	<b>5,107,277</b>	<b>2,460,947</b>	<b>8,909,039</b>	<b>11,369,986</b>	<b>22,945,536</b>
<b>Percent of State Total</b>	<b>28%</b>	<b>22%</b>	<b>11%</b>	<b>39%</b>	<b>50%</b>	<b>100%</b>

<sup>1/</sup> Federal and State Protected Lands not Managed for Timber Production includes forestlands set aside for wilderness, late successional reserves, managed late successional reserves, adaptive management areas, national wildlife refuges, national parks, Washington State parks, and Washington Department of Fish and Wildlife lands.

<sup>2/</sup> Federal and Tribal Lands Available for Timber Management include U.S. Forest Service Matrix lands, other U.S. Forest Service lands, BLM lands, Department of Defense lands, and all tribal lands.

<sup>3/</sup> Washington DNR and Other State Lands include all Washington DNR, Department of Corrections, and University lands.

Source: U.S. Geological Survey Land Use/Land Cover and WRIA GIS layers 2004



1     **3.2.2     Existing Habitat Conservation Plans**

2     As of June 2004, there were 11 HCPs in the State of Washington that had been approved  
3     by the Services (Table 3-4). Although the specific activities covered and the mitigation  
4     requirements vary under each plan depending on the interests of the landowners, most were  
5     developed for forest management activities. The only exception to this is the Daybreak  
6     Mine HCP, which covers floodplain-adjacent mining. The largest HCP, completed in  
7     1997, covers approximately 1.6 million acres of State trust lands managed by Washington  
8     DNR.

9     **3.2.3     Land Ownership and Use by Region**

10    The amount of forestland by region ranges from approximately 13,000 acres in the  
11    Columbia River Basin analysis region to just over 4 million acres in the Upper Columbia –  
12    Upstream of Grand Coulee Region (Table 3-2). Land ownership and use is summarized by  
13    analysis region in Tables 3-1, 3-2, and 3-3 and discussed in the following subsections.  
14    More detailed descriptions are found in the Region Descriptions in DEIS Appendix A. The  
15    analysis regions are shown in Figure 3-1, which also identifies those forestlands that are  
16    subject to the Washington Forest Practices Rules.

17    **3.2.3.1   Western Washington**

18    **North Puget Sound**

19    The North Puget Sound Region is approximately 4,398,000 acres in size. Approximately  
20    3,427,000 acres, or 78 percent, of this area is forestland. Agricultural lands make up 7  
21    percent of the Region, and residential and commercial land uses make up 2 percent.  
22    Ice/snow and bare rock makes up 6 percent. Other land use/land cover types each make up  
23    3 percent or less (Table 3-2).

24    Approximately 48 percent of the forestlands in this Region are managed under a Federal or  
25    State protected status that generally does not allow timber production. Approximately 7  
26    percent of the forestlands are under other Federal or tribal management and are potentially  
27    available for timber production; the remaining 45 percent of the forestlands are State,  
28    county, city, and private lands that are potentially available for timber management under  
29    the Washington Forest Practices Rules (Table 3-3).

30    **South Puget Sound**

31    The South Puget Sound Region is approximately 2,178,000 acres in size. Approximately  
32    1,532,000 acres, or 70 percent, of this area is forestland. Developed residential and  
33    commercial lands make up 17 percent of the Region (primarily the Seattle-Tacoma area),  
34    and agricultural lands make up 5 percent. Other land use/land cover types each make up 3  
35    percent or less (Table 3-2).

36    Approximately 19 percent of the forestlands in the Region are managed under a Federal or  
37    State protected status that generally does not allow timber production. Approximately 8  
38    percent of the forestlands are under other Federal or tribal management and are potentially  
39    available for timber production; the remaining 73 percent of the forestlands are State,

40



## Chapter 3

1 **Table 3-4.** Completed Habitat Conservation Plans in Washington State (as of  
2 October 2004).

Name	Species	Approximate HCP Start Date <sup>1/</sup>	Status	Acres <sup>2/</sup>
West Fork Timber <sup>3/</sup>	Spotted Owl	1992	Completed 1993	53,500
West Fork Timber	All Species	1994	Completed 1995	See Above
Scofield	Spotted Owl	1996	Completed 1996 <sup>4/</sup>	40
Plum Creek (Cascades)	All Vertebrates	1993	Completed 1996	170,000
Port Blakely (Robert B. Eddy)	All Species	1994	Completed 1996	7,500
Washington DNR State Trust Lands	All Species <sup>6/</sup>	1993	Completed 1997	1,600,000 <sup>6/</sup>
Seattle Public Utilities	Multiple Species	1994	Completed 2000	91,000
Green Diamond Resource Co. <sup>5/</sup>	Multiple Species	1997	Completed 2000	262,000
City of Tacoma/Tacoma Water	Multiple Species	1997	Completed 2001	15,000
Boise Cascade	Spotted Owl	2001	Completed 2001	620
Day Break Mine (Storehdahl)	Aquatic Species	1999	Completed 2004	300

<sup>1/</sup> Start dates are approximate. Applicants often prepare in advance of initiating active involvement with the Services.

<sup>2/</sup> Acres presented here are rounded from acres reported in the original HCP documents. In some cases, lands have been added to or subtracted from that reported in the original documents, and actual acres managed presently under the HCPs may be slightly different.

<sup>3/</sup> Previously known as the Murray-Pacific Corporation; name was changed to the original company name.

<sup>4/</sup> The original documents were completed in 1996. However, unlike the other completed HCPs, this resulted in a short-term (1 year) permit, which has since expired. The mitigation continues in the form of a perpetual deed restriction.

<sup>5/</sup> Previously known as the Simpson Resource Company.

<sup>6/</sup> Aquatic species are not covered on approximately 228,000 acres of State lands on the eastside of the Cascade Crest.

Source: USFWS 2004a.

3  
4 county, city, and private lands that are potentially available for timber management under  
5 the Washington Forest Practices Rules (Table 3-3).

### 6 **West Puget Sound**

7 The West Puget Sound Region is approximately 1,735,000 acres in size. Approximately  
8 1,522,000 acres, or 88 percent, of this area is forestland. Developed residential and  
9 commercial lands make up 4 percent of the Region, and agricultural lands make up 3  
10 percent. Other land use/land cover types each make up 2 percent or less (Table 3-2).

11 Approximately 41 percent of the forestlands in the Region are managed under a Federal or  
12 State protected status that generally does not allow timber production. Approximately 1



1 percent of the forestlands are under other Federal or tribal management and are potentially  
2 available for timber production; the remaining 57 percent of the forestlands are State,  
3 county, city, and private lands that are available for timber management under the  
4 Washington Forest Practices Rules (Table 3-3).

### 5 **Islands**

6 The Islands Region is approximately 247,000 acres in size. Approximately 180,000 acres,  
7 or 73 percent, of this area is forestland. Agricultural lands make up 17 percent of the  
8 Region, and developed residential and commercial lands make up 6 percent. Other land  
9 use/land cover types each make up 2 percent or less (Table 3-2).

10 Approximately 6 percent of the forestlands in this Region are managed under a Federal or  
11 State protected status that generally does not allow timber production. Approximately 2  
12 percent of the forestlands are under other Federal or tribal management and are potentially  
13 available for timber production; the remaining 92 percent of the forestlands are State,  
14 county, city, and private lands that are available for timber management under the  
15 Washington Forest Practices Rules (Table 3-3).

### 16 **Olympic Coast**

17 The Olympic Coast Region is approximately 1,757,000 acres in size. Approximately  
18 1,671,000 acres, or 95 percent, of this area is forestland. All other land use/land cover  
19 types each make up 2 percent of the area or less (Table 3-2).

20 Approximately 41 percent of the forestlands in this Region are managed under a Federal or  
21 State protected status that generally does not allow timber production. Approximately 14  
22 percent of the forestlands are under other Federal or tribal management and are potentially  
23 available for timber production; the remaining 45 percent of the forestlands are State,  
24 county, city, and private lands that are available for timber management under the  
25 Washington Forest Practices Rules (Table 3-3).

### 26 **Southwest**

27 The Southwest Region is approximately 2,313,000 acres in size. Approximately 2,058,000  
28 acres, or 89 percent, of this area is forestland. Agricultural lands make up 6 percent of the  
29 Region, and all other land use/land cover types each make up 2 percent of the area or less  
30 (Table 3-2).

31 Approximately 7 percent of the forestlands in this Region are managed under a Federal or  
32 State protected status that generally does not allow timber production. Less than 1 percent  
33 of the forestlands are under other Federal or tribal management and are potentially  
34 available for timber production; the remaining 93 percent of the forestlands are State,  
35 county, city, and private lands that are potentially available for timber management under  
36 the Washington Forest Practices Rules (Table 3-3).

### 37 **Lower Columbia**

38 The Lower Columbia Region is approximately 3,072,000 acres in size. Approximately  
39 2,616,000 acres, or 85 percent, of this area is forestland. Agricultural lands make up 6  
40 percent of the Region, residential and commercial lands make up 2 percent and water and



## Chapter 3

---

1 wetlands make up 3 percent. Other land use/land cover types each make up 2 percent or  
2 less (Table 3-2).

3 Approximately 27 percent of the forestlands in this Region are managed under a Federal or  
4 State protected status that generally does not allow timber production. Approximately 10  
5 percent of the forestlands are under other Federal or tribal management and are potentially  
6 available for timber production; the remaining 62 percent of the forestlands are State,  
7 county, city, and private lands that are potentially available for timber management under  
8 the Washington Forest Practices Rules (Table 3-3).

### 9 **3.2.3.2 Eastern Washington**

#### 10 **Middle Columbia**

11 The Middle Columbia Region is approximately 6,497,000 acres in size. Approximately  
12 2,691,000 acres, or 41 percent, of this area is forestland. Shrubland and grassland  
13 combined make up 38 percent of the Region, and agricultural lands make up 17 percent.  
14 Other land use/land cover types each make up 1 percent or less (Table 3-2).

15 Approximately 33 percent of the forestlands in this Region are managed under a Federal or  
16 State protected status that generally does not allow timber production. Approximately 32  
17 percent of the forestlands are under other Federal or tribal management and are potentially  
18 available for timber production; the remaining 35 percent of the forestlands are State,  
19 county, city, and private lands that are potentially available for timber management under  
20 the Washington Forest Practices Rules (Table 3-3).

#### 21 **Upper Columbia River Downstream of Grand Coulee**

22 The Upper Columbia – Downstream of Grand Coulee Dam Region is approximately  
23 6,401,000 acres in size. Approximately 2,774,000 acres, or 43 percent, of this area is  
24 forestland. Shrubland and grassland combined make up 43 percent of the Region, and  
25 agricultural lands make up 9 percent. Other land use/land cover types each make up 2  
26 percent or less (Table 3-2).

27 Approximately 46 percent of the forestlands in this Region are managed under a Federal or  
28 State protected status that generally does not allow timber production. Approximately 37  
29 percent of the forestlands are under other Federal or tribal management and are potentially  
30 available for timber production; the remaining 17 percent of the forestlands are State,  
31 county, city, and private lands that are potentially available for timber management under  
32 the Washington Forest Practices Rules (Table 3-3).

#### 33 **Upper Columbia River Upstream of Grand Coulee**

34 The Upper Columbia – Upstream of Grand Coulee Dam Region is approximately  
35 5,748,000 acres in size. Approximately 4,084,000 acres, or 71 percent, of this area is  
36 forestland. Shrubland and grassland combined make up 13 percent of the Region, and  
37 agricultural lands make up 12 percent. Other land use/land cover types each make up 2  
38 percent or less (Table 3-2).

39 Approximately 3 percent of the forestlands in this Region are managed under a Federal or  
40 State protected status that generally does not allow timber production. Approximately 53



1 percent of the forestlands are under other Federal or tribal management and are potentially  
2 available for timber production; the remaining 44 percent of the forestlands are State,  
3 county, city, and private lands that are potentially available for timber management under  
4 the Washington Forest Practices Rules (Table 3-3).

### 5 **Snake River**

6 The Snake River Region is approximately 4,576,000 acres in size. Approximately 376,000  
7 acres, or 8 percent, of this area is forestland. Shrubland and grassland combined make up  
8 37 percent of the Region, and agricultural lands make up 52 percent. Other land use/land  
9 cover types each make up 1 percent or less (Table 3-2).

10 Approximately 25 percent of the forestlands in this Region are managed under a Federal or  
11 State protected status that generally does not allow timber production. Approximately 40  
12 percent of the forestlands are under other Federal or tribal management and are potentially  
13 available for timber production; the remaining 35 percent of the forestlands are State,  
14 county, city, and private lands that are potentially available for timber management under  
15 the Washington Forest Practices Rules (Table 3-3).

### 16 **Columbia Basin**

17 The Columbia Basin Region is approximately 4,350,000 acres in size. Approximately  
18 13,000 acres, or less than 1 percent, of this area is forestland. Shrubland and grassland  
19 combined make up 43 percent of the Region, and agricultural lands make up 53 percent.  
20 Open water and wetlands make up 3 percent and other land use/land cover types each make  
21 up 1 percent or less (Table 3-2).

22 Less than 1 percent of the forestlands in this Region are managed under a Federal or State  
23 protected status that generally does not allow timber production. Approximately 1 percent  
24 of the forestlands are under other Federal or tribal management and are potentially  
25 available for timber production; the remaining 99 percent of the forestlands are State,  
26 county, city, and private lands that are potentially available for timber management under  
27 the Washington Forest Practices Rules (Table 3-3).

### 28 **3.2.3.3 Summary**

29 The distribution of lands potentially available for timber management under the  
30 Washington Forest Practices Rules is shown graphically in Figure 3-1. Approximately 70  
31 percent of these lands are located in western Washington. Four of the 12 analysis regions  
32 each account for more than 10 percent of these lands. The Southwest, Upper Columbia-  
33 Upstream of Grand Coulee Dam, Lower Columbia, and North Puget Sound Regions  
34 accounted for 17 percent, 16 percent, 14 percent, and 14 percent of the statewide total,  
35 respectively (Figure 3-1).

36 Lands potentially available for timber management under the Washington Forest Practices  
37 Rules comprise a relatively large share of total forestlands in the Southwest (79 percent),  
38 South Puget Sound (63 percent), Lower Columbia (50 percent), and Upper Columbia-  
39 Upstream of Grand Coulee Dam (44 percent) Regions.



## Chapter 3

---

### 3.2.4 Timber Harvest Rates for Western and Eastern Washington

As discussed in subsection 3.6.1 (Forest Vegetation), forest stand conditions in western and eastern Washington vary in terms of levels of precipitation, stand composition, densities, and disturbance regimes. As a result, different harvest strategies and harvest levels, as well as different land uses are typically seen in each half of the State. The following subsections address western and eastern Washington in turn.

#### 3.2.4.1 Western Washington

Approximately 13 million acres or 83 percent of the land base in western Washington is forestland. Approximately 4.1 million acres, or 32 percent, of the total forested acres are generally unavailable for harvest due to some form of protected status leaving 8.9 million acres (68 percent) potentially available for timber harvest (Table 3-3). Forestlands in western Washington account for about 57 percent of the forestland in the State (Table 3-2), but have historically provided over 80 percent of the total timber harvest (Adams et al. 1992). Annual harvest levels for 1990 through 2002 are displayed by ownership in subsection 3.14 (Social and Economic Environment). These data indicate that 75 percent of the statewide harvest occurred in western Washington in 2002, with the remaining 25 percent taking place in eastern Washington (Washington DNR 2004b).

Private landowners (including both large and small forest landowners) account for roughly 6.1 million acres or about 47 percent of the total westside forestland base. Historically, private landowners have accounted for a large share of the overall westside timber harvest, with rates between 1949 and 2002 averaging around 72 percent of the total westside harvests (Table 3-5), and 59 percent of the total statewide harvest (Table 3-6). Private landowners accounted for 84 percent of the total westside harvest in 2002, as well as 84 percent of total statewide harvest (Washington DNR 2004b).

#### 3.2.4.2 Eastern Washington

Forestlands comprise a much smaller portion of the total land base in eastern Washington than they do in western Washington. This is primarily due to a combination of drier growing conditions and relatively high percentages of agricultural lands and naturally occurring shrub-steppe lands in eastern Washington (Table 3-2). Approximately 9.9 million acres or 36 percent of the land base in eastern Washington is forestland (Table 3-2). Approximately 2.3 million acres or 24 percent of this forestland is unavailable for harvest due to some form of protection status, leaving 7.6 million acres of forestland on the eastside available for harvest (Table 3-2). Forestlands in eastern Washington account for about 43 percent of the forestland in the State (Table 3-2), but have historically provided less than 20 percent of the total timber harvest (Adams et al. 1992). Eastern Washington accounted for 25 percent of statewide timber harvest in 2002 (Washington DNR 2004b).

Private landowners (including both large and small forest landowners) account for roughly 2.6 million acres or about 26 percent of the total eastside forestland base. Historically, private landowners in eastern Washington have contributed a large percentage of the overall eastside timber harvest, with rates between 1949 and 2002 averaging around 64 percent of the total eastside harvests (Table 3-7). Private landowners accounted for 84 percent of the total eastside harvest in 2002 (Washington DNR 2004b).



# Chapter 3



1 **Table 3-5.** Western Washington Timber Harvests by Ownerships, 1949-2002.

Owner Class	Total Harvest in MBF for 1985-2002	Percent of Total Harvest for 1985-2002 (%)	Total Harvest in MBF for 1949-1984	Percent of Total Harvest for 1949-1984 (%)	Total Harvest in MBF for 1949-2002	Percent of Total Harvest for 1949-2002 (%)
Native American	691,098	1.0	4,625,909	2.7	5,317,007	2.2
Forest Industry	33,744,611	47.2	102,038,423	60.5	135,783,034	56.5
Private, Large	10,878,685	15.2	5,033,071	3.0	15,911,756	6.6
Private, Small	9,430,217	13.2	7,151,184	4.2	16,581,401	6.9
<b>Total Private</b>	<b>54,744,611</b>	<b>76.5</b>	<b>118,848,587</b>	<b>70.4</b>	<b>173,593,198</b>	<b>72.2</b>
State	10,073,411	14.1	16,460,816	9.8	26,534,227	11.0
Other Non-Federal	406,856	0.6	690,527	0.4	1,097,383	0.5
National Forest	6,130,203	8.6	32,147,529	19.1	38,277,732	15.9
Other Federal	193,927	0.3	631,276	0.4	825,203	0.3
<b>Total Public</b>	<b>16,804,397</b>	<b>23.5</b>	<b>49,930,148</b>	<b>29.6</b>	<b>66,734,545</b>	<b>27.8</b>
<b>Total All Ownerships</b>	<b>71,549,008</b>	<b>--</b>	<b>168,778,735</b>	<b>--</b>	<b>240,327,743</b>	<b>--</b>

Source: Washington DNR's Washington Timber Harvest 2002 report published in 2004.

2

3 **Table 3-6.** Washington Timber Harvests in Thousand Board Feet (MBF) by  
4 Ownerships, 1949-2002.

Owner Class	Total MBF Harvested Eastern Washington	Total MBF Harvested Western Washington	Statewide Total MBF Harvested	Percent of Total Statewide Harvest (MBF) (%)
Native American	11,406,533	5,317,007	16,723,540	5.7
Forest Industry	14,726,109	135,783,034	150,509,143	51.3
Private, Large	2,299,286	15,911,756	18,211,042	6.2
Private, Small	5,633,936	16,581,401	22,215,337	7.6
<b>Total Private</b>	<b>34,065,864</b>	<b>173,593,198</b>	<b>207,659,062</b>	<b>70.8</b>
State	3,328,364	26,534,227	29,862,591	10.2
Other Non-Federal	210,296	1,097,383	1,307,679	0.5
National Forest	15,085,700	38,277,732	53,363,432	18.2
Other Federal	207,346	825,203	1,032,549	0.4
<b>Total Public</b>	<b>18,831,706</b>	<b>66,734,545</b>	<b>85,566,251</b>	<b>29.2</b>
<b>Total All Ownership</b>	<b>52,897,570</b>	<b>240,327,743</b>	<b>293,225,313</b>	<b>--</b>

Source: Washington DNR's Washington Timber Harvest 2002 report published in 2004.

5

6



# Chapter 3

1 **Table 3-7.** Eastern Washington Timber Harvests by Ownerships, 1949-2002.

Owner Class	Total Harvest in MBF for 1985-2002	Percent of Total Harvest for 1985-2002 (%)	Total Harvest in MBF for 1949-1984	Percent of Total Harvest for 1949-1984 (%)	Total Harvest in MBF for 1949-2002	Percent of Total Harvest for 1949-2002 (%)
Native American	3,793,798	19.7	7,612,735	22.6	11,406,533	21.6
Forest Industry	5,256,791	27.3	9,469,318	28.1	14,726,109	27.8
Private, Large	1,277,501	6.6	1,021,785	3.0	2,299,286	4.4
Private, Small	3,538,796	18.4	2,095,140	6.2	5,633,936	10.7
<b>Total Private</b>	<b>13,866,886</b>	<b>72.1</b>	<b>20,198,978</b>	<b>60.0</b>	<b>34,065,864</b>	<b>64.4</b>
State	1,556,742	8.1	1,771,622	5.3	3,328,364	6.3
Other Non-Federal	61,035	0.3	149,261	0.4	210,296	0.4
National Forest	3,730,678	19.4	11,355,022	33.7	15,085,700	28.5
Other Federal	27,972	0.2	179,374	0.5	207,346	0.4
<b>Total Public</b>	<b>5,376,427</b>	<b>27.9</b>	<b>13,455,279</b>	<b>40.0</b>	<b>18,831,706</b>	<b>35.6</b>
<b>Total All Ownership</b>	<b>19,243,313</b>	<b>--</b>	<b>33,654,257</b>	<b>--</b>	<b>52,897,570</b>	<b>--</b>

Source: Washington DNR's Washington Timber Harvest 2002 report published in 2004.

2

### 3 3.2.5 Forestland Conversion

4 As the population of Washington State continues to grow, lands are being converted from  
 5 forestland to other uses. Comprehensive tracking of forestland conversion rates began in  
 6 the late 1970s, with the USDA Forest Service Forest Inventory and Analysis data  
 7 (Bolsinger et al. 1997). The Forest Service conducted inventories in 1978 and 1979 and  
 8 again in 1988 and 1989. Historic data for years prior to 1978 tend to be less consistent.  
 9 Some information is, however, available from the 1930s through the 1970s, and this  
 10 information may be used to approximate general trends leading to present day conditions.

11 The following discussion is divided into three parts that address land conversion trends  
 12 from the 1930s through 1991, statewide data compiled from 1997 through 2003 from the  
 13 Washington DNR Forest Practices Application Review System, and data compiled for  
 14 King County since 1982.

#### 15 3.2.5.1 Forestland Conversion from the 1930s to 1991

16 This subsection is divided into three parts: Washington forestlands in the 1930s, forestland  
 17 conversion from 1945 to 1970, and forestland conversion from 1978 to 1991.



1 **Washington Forestlands in the 1930s**  
 2 Data compiled from early forest inventory surveys conducted in the 1930s indicate there  
 3 were approximately 26.5 million acres of forestlands in Washington State at that time  
 4 (Table 3-8). These data were collected using a combination of methods including field  
 5 surveys, analyses of aerial photos, assessment of tax information followed up by field  
 6 verification, and review of county records and stocking classification of previously logged  
 7 areas. Approximately 12.5 million acres or 47 percent of forestlands in the State were in  
 8 private ownership, with 9.1 million acres or 73 percent of private forestlands located in  
 9 western Washington (Table 3-8). Private forestlands made up 54 percent and 32 percent of  
 10 forested lands in western and eastern Washington, respectively.

11 The total acres of forestlands available for harvest statewide in all ownerships was  
 12 approximately 25.2 million acres, of which 15.2 million acres (60 percent) occurred in the  
 13 western half of the State, and 10.0 million (40 percent) occurred in the eastern half.  
 14 Federal and State forested reserve lands accounted for 804,848 acres statewide or 3 percent  
 15 of the total acres of forestland. Of this, 341,178 acres (42 percent) were in western  
 16 Washington, and 463,670 acres (58 percent) were in eastern Washington (Table 3-8).

17 Comparison between these data and estimated forestlands in Washington in 2004 (Tables  
 18 3-2 and 3-3) suggests that there has been a net loss of approximately 3.5 million acres of  
 19 forestland since the 1930s, with the majority of this loss (80 percent or 2.8 million acres)  
 20 occurring in western Washington. These data also suggest that reductions in the amount of  
 21 privately owned forestland accounted for the majority of this loss.

22 **Table 3-8.** Acres of Forestland in Washington State by Ownership in the early  
 23 1930s.

<b>Land Ownership</b>	<b>Western Washington (Acres)</b>	<b>Eastern Washington (Acres)</b>	<b>Total (Acres)</b>
Private	9,055,874	3,432,730	12,488,604
State- Available for harvest	865,346	617,910	1,483,256
State Reserved	11,882	1,680	13,562
County/City	344,882	347,995	692,877
Tribal- Available for harvest	250,648	1,516,490	1,767,138
Federal-Available for harvest	4,652,531	4,075,220	8,727,751
Federal, Other Forestland <sup>1/</sup>	356,330	190,685	547,015
Federal Reserved	329,296	461,990	791,286
<b>Total</b>	<b>15,849,289</b>	<b>10,644,700</b>	<b>26,493,989</b>

<sup>1/</sup>Includes railroad lands, tribal land grants, and miscellaneous lands.

Source: Table modified from Harrington 2003.



## Chapter 3

### 1 **Forestland Conversion from 1945 to 1970**

2 As the population of Washington State continues to grow, lands are being converted from  
 3 forestland to other uses. Comprehensive tracking of forestland conversion rates began in  
 4 the late 1970s with the U.S. Forest Service Inventory and Analysis data (also known as  
 5 FIA), which are available in the U.S. Forest Service Resource Bulletin PNW-RB-46  
 6 (Bolsinger et al. 1997). Data presented in the bulletin estimates that 630,000 acres of  
 7 commercial forestlands were converted to non-forest uses in Washington over this period,  
 8 with conversion to urban-industrial, agricultural, and road use (Table 3-9). By 1970 there  
 9 were an estimated 23.1 million acres of forestland in Washington State, with  
 10 approximately 4.7 million acres in some kind of reserve status (1.6 million acres) or  
 11 considered not capable of growing commercial timber (3.1 million acres) (Table 3-10).

12 **Table 3-9.** Conversion of Commercial Forestland in Washington State by  
 13 Ownership and Land Use, 1945-1970 (In Thousand Acres).

Ownership	Roads	Reservoirs	Powerlines	Farms <sup>1/</sup>	Urban-Industrial	Miscellaneous	Total
National Forest	33	10	3	0	0	0	46
Other Public	22	1	33	-18 <sup>2</sup>	35	0	73
Forest Industry	50	0	38	13	8	0	109
Farm and Miscellaneous Private	27	9	16	135	214	1	402
<b>Total</b>	<b>132</b>	<b>20</b>	<b>90</b>	<b>130</b>	<b>257</b>	<b>1</b>	<b>630</b>

<sup>1/</sup> Farms include lands converted to both agricultural farming and Christmas tree farms.

<sup>2/</sup> Minus indicates a gain of forestland.

Source: Table modified from Bolsinger 1973.

15 **Table 3-10.** Acres of Forestland by Land Class in Washington State, 1970  
 16 (In Thousand Acres).

Land Class	Acres of Forestland
Commercial Forest	18,401
Commercial Reserved Forest	1,446
Noncommercial Forest <sup>1</sup>	3,108
Deferred Forest <sup>2</sup>	143
<b>Total</b>	<b>23,098</b>

<sup>1/</sup> This report defined Noncommercial forest to mean forestland that is not capable of growing 20 cubic feet of industrial wood per year, or is too steep and rocky for harvesting and growing timber crops.

<sup>2/</sup> This report defined Deferred forest to mean commercial forestland within National Forests that was being considered for wilderness designation in 1973.

Source: Table modified from Bolsinger 1973.



## 1 Forestland Conversion from 1978 to 1991

2 Between 1978 and 1991, Bolsinger et al. (1997) estimated that lands available for timber  
 3 production in Washington outside of the National Forests decreased by 488,000 acres.  
 4 This figure includes approximately 117,000 net acres of private timberlands that were  
 5 transferred to the National Forest System, and an additional 92,000 acres (mostly tribal)  
 6 that were reclassified to reserve status, meaning that they were not available for timber  
 7 harvest but were still forested lands. Conversions to another use accounted for  
 8 approximately 279,000 acres. Clearing for rights-of-way for roads, pipelines, and other  
 9 uses accounted for 155,000 acres or 56 percent of this total (Table 3-11). Most lands  
 10 converted to rights-of-way were for construction of roads or landings used for logging and  
 11 forest access, or in the case of other private lands, for connection of private properties, or  
 12 construction of new or expanding highways or freeways (Bolsinger et al. 1997).  
 13 Conversions to urban development accounted for 89,000 acres, with the majority of this  
 14 type of conversion (92 percent) occurring on private, non-industrial forestlands  
 15 (Table 3-11).

16 **Table 3-11. Changes in Forestland Area by Ownership and Land Use in**  
 17 **Washington State from 1978-1991.**

	Other Public <sup>1/</sup>	Forest Industry <sup>2/</sup>	Other Private <sup>3/</sup>	Total
Rights-of-way	-12,000	-95,000	-48,000	-155,000
Urban	0	-7,000	-82,000	-89,000
Agriculture <sup>4/</sup>	0	0	-53,000	-53,000
Reclaimed Forest	0	6,000	12,000	18,000
<b>Total</b>	<b>-12,000</b>	<b>-96,000</b>	<b>-171,000</b>	<b>-279,000</b>

<sup>1/</sup> Lands administered by public agencies other than the USDA Forest Service.

<sup>2/</sup> Lands owned by companies that grow timber for industrial uses.

<sup>3/</sup> Private lands not owned by the forest industry, but including tribal lands, farmer-owned lands, and miscellaneous other private lands.

<sup>4/</sup> Agricultural lands, including Christmas tree farms, converted to forestland.

Source: Bolsinger et al. 1997

18  
 19 Bolsinger et al. (1997) also considered changes over this period in the “primary forest  
 20 zone,” which they defined as large tracts of forestlands with no more than one development  
 21 per 640 acres and containing roads that are used primarily for resource extraction and are at  
 22 least one-quarter mile apart. Uses of this land are primarily restricted to timber production,  
 23 grazing, watershed, and wildlife protection. Forestlands within this zone were estimated in  
 24 the 1978 to 1980 inventory to be 7,143,000 acres in western Washington and 6,397,000  
 25 acres in eastern Washington. Approximately 10 years later, the 1988 to 1991 inventory  
 26 estimated approximately 6,729,000 acres in this zone in western Washington (a loss of  
 27 414,000 acres or 6 percent) and 6,384,000 acres in eastern Washington (a loss of 13,000  
 28 acres or 0.2 percent) (Bolsinger et al. 1997).

29 Adams et al. (1992) found that between 1980 and 1990 all ownership groups had a net loss  
 30 of forest acres except for non-industrial private landowners, which reported a net gain of  
 31 169,000 acres. However, it was not clear if this gain was from lands converted back into



## Chapter 3

1 forest production or if it was simply land purchases. Adams et al. (1992) attributed 90  
 2 percent of the total acreage of forestland conversions to non-forest uses between 1980 and  
 3 1990 to development such as urban uses (130,000 acres) or rights-of-way development  
 4 (50,000 acres).

### 5 **3.2.5.2 Washington Department of Natural Resources Data**

6 Four classes of forest practices have been established based on the potential of planned  
 7 activities to adversely impact public resources. Forest practices are classed as I, II, III, or  
 8 IV, with Class I having no potential of damaging public resources and Class IV having the  
 9 greatest potential. Class IV forest practices are further distinguished as Class IV-Special or  
 10 Class IV-General by rule of the Washington Forest Practices Board. Class IV-Special  
 11 forest practices are those that have been determined to have potential for a substantial  
 12 impact on the environment. Class IV-General are forest practices on lands platted after  
 13 January 1, 1960, or on lands being converted to another use, or that will not be reforested  
 14 because of a likelihood of future conversion to urban development.

15 Conversion information available from Washington DNR's Forest Practices Application  
 16 Review System database indicate that 53,821 acres of forestland were converted to other  
 17 uses between 1997 and 2003, with an average of 7,687 acres per year statewide (Table  
 18 3-12). These data are based on Class IV-General forest practices applications approved by  
 19 the Washington DNR. Some margin of error is expected due to the fact that not all  
 20 landowners report conversions of lands that were harvested under a different application  
 21 class, and not all landowners who apply to convert their lands actually follow through with  
 22 the land conversion.

23 **Table 3-12. Acres of Forestlands Converted to Other Uses by Year.**

Year	Total Acres Western Washington	Total Acres Eastern Washington	Total Acres Statewide
1997	4,059.2	3,095.6	7,154.8
1998	3,748.1	3,675.8	7,423.9
1999	3,320.7	3,351.1	6,702.2
2000	3,143.8	1,863.8	5,007.6
2001	7,374.2	1,258.6	8,632.8
2002	2,223.6	3,794.9	6,018.5
2003	10,083.5	2,797.3	12,880.8
<b>Total</b>	<b>33,953</b>	<b>19,837</b>	<b>53,821</b>

Source: Washington DNR FPARS Data Base, January 2004.

24

### 25 **Transfer of Authority for Class IV – General Applications**

26 While the Forest Practices Board is the entity responsible for establishing forest practices  
 27 standards that serve as the basis for the Forest Practices Regulatory Program, and the  
 28 Washington DNR is the agency charged with managing Program implementation, some



1 local governments have authority over Class IV-General forest practices within their  
2 jurisdiction. In accordance with the Revised Code of Washington (RCW) (RCW Chapter  
3 76.09.240), the Washington DNR is in the process of working with local governments to  
4 transfer jurisdiction of Class IV-General forest practices.

5 Each city and county in the State is required to adopt ordinances by December 31, 2005 to  
6 regulate Class IV-General forest practices. The city or county's ordinances or regulations  
7 must meet or exceed the standards set forth in the current Washington Forest Practices  
8 Rules. Washington DNR, in consultation with Washington Department of Ecology  
9 (Ecology), may approve or disapprove the regulations in whole or in part (RCW Chapter  
10 76.09.240(3)). To date, Washington DNR has transferred authority to regulate Class IV-  
11 General forest practices to four counties: Thurston, King, Spokane, and Clark Counties.  
12 Generally, forest practices inside designated urban growth areas are likely to be associated  
13 with future conversion to another land use, while forest practices outside urban growth  
14 areas are not. In high population growth areas, development pressures may result in more  
15 Class IV-General conversions outside the urban growth area than in areas with lower  
16 growth rates.

### 17 **3.2.5.3 King County Data**

18 King County, the most populated county in the State, has kept relatively complete data on  
19 Class IV-General conversions of forestland since about 1982. The average conversion rate  
20 from 1982 to 1987 was about 675 acres per year, which grew in the late 1980s to an  
21 average of 1,500 acres per year (Personal Communication, Chandler Felt, King County  
22 Department of Development and Environmental Studies, February 19, 2004). Conversion  
23 rates in King County slowed to an average of about 600 acres per year after the adoption of  
24 the 1990 Growth Management Act and because of economic slowdowns also experienced  
25 at that time. In 1998, another short-lived conversion boom occurred (1,500 acres), after  
26 which the average dropped back down to 490 acres per year (Personal Communication,  
27 Chandler Felt, King County Department of Development and Environmental Studies,  
28 February 19, 2004) (King County 2000) (Table 3-13). Additionally, in 1994, King County  
29 designated Forest Production Districts under the Growth Management Act, which  
30 established zoning restrictions and other regulations designed to encourage timber  
31 production in those districts.

32



## Chapter 3

1 **Table 3-13.** Forest Practices Applications and Acres of Converted Forestland in  
 2 King County, 1990 through 1998.

Year	Total Acres Converted	Inside Forest Production Areas		Outside Forest Production Areas	
		Forested Acres Converted	Number of Conversion Applications	Forested Acres Converted	Number of Conversion Applications
1990	728	1	5	727	56
1991	426	71	12	355	17
1992	445	7	1	438	27
1993	1,131	13	4	1,118	96
1994	306	0	0	306	32
1995	674	0	0	674	41
1996	754	4	1	750	55
1997	541	58	3	483	34
1998	1,483	145	5	1,338	27
<b>Total</b>	<b>6,488</b>	<b>299</b>	<b>31</b>	<b>6,189</b>	<b>385</b>

Source: King County 2000 Annual Growth Report; Personal Communication, Chandler Felt, King County Department of Development and Environmental Services, February 19, 2004.

3



**3.3 AIR QUALITY**

Air quality is regulated by the Federal Clean Air Act, which requires the Environmental Protection Agency (EPA) to set national ambient air quality standards for pollutants considered harmful to public health and the environment. “Ambient air” refers to that portion of the atmosphere, external to buildings, to which the general public has access. An air quality standard establishes values for maximum acceptable concentration, exposure time, and frequency of occurrence of one or more air contaminants in the ambient air. Ambient air quality standards have been set for six principal pollutants: carbon monoxide, nitrogen dioxide, ozone, lead, particulate matter, and sulfur dioxide. The State of Washington has an approved State Implementation Plan, which regulates, among other pollutants and emissions from prescribed burning (Washington Department of Ecology 1999a). The State Implementation Plan also addresses particulate matter (including “PM<sub>10</sub>”), visibility, and smoke management.

Prescribed burning on forestland is regulated by Washington DNR’s Resource Protection Division, which requires a permit for burning. Washington DNR’s federally mandated Smoke Management Plan (Washington Administrative Code [WAC] 173-425-120; State adopted October 18, 1990, EPA effective January 15, 1993) ensures that forest management-related burning activities comply with the Clean Air Act and provides regulatory direction, operating procedures, and information regarding the management of smoke and fuels on the forestlands of Washington (Washington DNR 1993). The plan coordinates and facilitates the statewide regulation of prescribed burning on Federal and non-Federal forestlands and participating tribal lands. Washington DNR follows the guidelines set forth in this plan when issuing and regulating burning permits for open fire when such fires are for the following:

1. Abating a forest fire hazard
2. Preventing a fire hazard in a forested area
3. Instructing public officials in methods of forest fire fighting
4. Any silvicultural operation to improve the forestlands of the State

In particular, three potential sources of particulate air pollution associated with forest management activities are slash burning, wildfire, and road use. Managers of non-Federal forestlands in Washington typically use slash burning as part of their site preparation activities, usually by concentrating slash in piles and burning the piles (slash concentrated burning) rather than by broadcast burning, as was more common in the past. Broadcast burning is the practice of burning logging slash scattered throughout a recently harvested unit to prepare the site for planting and/or to reduce dangerous fuel loads. Burning is usually done in the spring or fall under wetter conditions when fewer particulates are emitted than would be the case if the same fuels burned in a wildfire. Particulate emissions from wildfires are, on average, three to four times higher than from prescribed burning (Washington DNR 1996).

The use of logging roads during dry weather conditions generates air-borne dust. Air-borne dust is regulated through road maintenance standards of the Washington Forest Practices Board (WAC 222-24) and safety standards of the Washington Department of



## Chapter 3

---

1 Labor and Industries (WAC 296-54). The amount of air-borne dust is a function of road  
2 use and surfacing material. Gravel can reduce dust on dirt roads (Washington Department  
3 of Ecology 2001) as can water and chemical dust suppressants. In general, the adverse  
4 effects of air-borne dust are localized and short term (Washington DNR 1996).

5 The main sources of air pollution in western Washington include motor vehicles (55  
6 percent), industrial (13 percent), and wood stoves (9 percent). Approximately 4 percent is  
7 generated from outdoor burning, a portion of which comes from forest management  
8 activities (Washington Department of Ecology 2003). Air quality in Washington is  
9 generally good or moderate, although some areas do not meet Federal standards on some  
10 days. Air quality has improved greatly since 1987, when Washington violated air quality  
11 standards on 150 days. This figure dropped to 7 percent in 1999 (Washington Department  
12 of Ecology 2003).

13 One of the ecological benefits of forested lands is the enhancement of air quality. Plants  
14 enhance air quality by emitting oxygen and consuming carbon dioxide, the gas most  
15 associated with global warming. In addition, trees retard the spread of air-borne  
16 particulates by trapping the material on their leaf surfaces and by slowing the wind speed  
17 to the point that particulates cannot remain suspended. Timber harvesting temporarily  
18 removes the air quality benefits provided by trees (Washington DNR 1996).

19



1 **3.4 GEOLOGY, SOILS, AND EROSIONAL PROCESSES**


2 **3.4.1 Geology and Soils**

3 The geology and soils of Washington are highly variable and complex. Because lands  
4 covered by the Forest Practices Habitat Conservation Plan (FPHCP) are widely distributed  
5 throughout the State, the geologic and soil conditions on those lands are similarly complex.  
6 The physiographic provinces of the State are shown in Figure 3-2. Overviews of the  
7 geologic and soil conditions on covered lands are provided below and have been grouped  
8 according to geologically similar analysis regions (using the analysis regions described in  
9 Section 3.1, Introduction). Region-specific descriptions of the geology and soils can be  
10 found in DEIS Appendix A.

11 **3.4.1.1 North Puget, South Puget, West Puget, Olympic, and Islands**  
12 **Regions**

13 The North Puget, South Puget, West Puget, Olympic, and Islands Regions encompass the  
14 entire Olympic Mountains physiographic province and parts of the Northern Cascades,  
15 Southern Cascades, Puget Lowland, and Willapa Hills physiographic provinces as defined  
16 by Lasmanis (1991). The physiography of these five regions reflects widespread glacial  
17 activity that occurred during the Fraser Glaciation approximately 25,000 to 10,000 years  
18 ago. Alpine glaciers extending out of the Cascade and Olympic mountain ranges carved  
19 U-shaped river valleys while continental glaciers pushed through what are now Puget  
20 Sound and the Strait of Juan de Fuca.

21 **Figure 3-2.** Physiographic Provinces of Washington State.



22  
23  
24

Source: Lasmanis 1991.



## Chapter 3

---

1 In addition to shaping the topography of the regions, glaciers blanketed large areas with  
2 till, outwash, and lake sediments. These glacial deposits served as parent materials on  
3 which many of the regions' soils have developed. Glaciers left behind landforms that  
4 range from nearly flat glacial plains in the Puget Lowland to very long, steep mountain  
5 slopes in the Northern and Southern Cascades and Olympic Mountains physiographic  
6 provinces.

7 While Quaternary glacial deposits dominate areas that lie within the Puget Lowland  
8 physiographic province, other areas contain highly diverse rock types. The most common  
9 geologic units in the Northern Cascades and Olympic Mountains include sedimentary and  
10 volcanic rocks from the Lower Tertiary Period and Mesozoic Era and intrusive igneous  
11 rocks from the Tertiary Period.

12 The relatively short time since deglaciation has limited the degree of soil development in  
13 many parts of these regions. The glacial deposits and other parent materials remaining  
14 after deglaciation have not undergone the higher levels of physical and chemical  
15 weathering and related soil formation found in non-glaciated areas of the State. Soils  
16 developed on glaciated terrain tend to have much lower levels of organic matter in their  
17 surface horizons and less horizon development than older, more heavily weathered soils in  
18 other parts of Washington. Soils developed from glacial parent materials are common at  
19 low elevations, as are alluvial soils along major rivers and streams. The primary types of  
20 glacial parent materials, in order of their relative coverage, are glacial till, glacial outwash,  
21 and glacial lake sediments. At moderate to high elevations, soils are more commonly  
22 formed from a mixture of colluvial bedrock materials, glacial drift deposits, and volcanic  
23 ash (Washington DNR 1996).

### 24 **3.4.1.2 Southwest and Lower Columbia Regions**

25 The Southwest and Lower Columbia Regions encompass the Willapa Hills and Portland  
26 Basin physiographic provinces and parts of the Olympic Mountains, Puget Lowland, and  
27 Southern Cascades provinces as defined by Lasmanis (1991). Unlike analysis regions to  
28 the north, many parts of the Southwest and Lower Columbia Regions were never glaciated.  
29 As a result, highly weathered rocks that are relatively easily erodible remain widespread,  
30 particularly in the Southwest Region and the western portions of the Lower Columbia  
31 Region. Erosion of these rocks has produced hills and ridges that are rounded with short,  
32 steep slopes and low gradient stream channels.

33 The eastern part of the Lower Columbia Region includes portions of the Southern  
34 Cascades physiographic province that was subject to alpine glaciation during the  
35 Pleistocene Epoch. This area is similar to the glaciated northern regions where broad, U-  
36 shaped valleys and long, steep slopes are common. Typical geologic units in the  
37 Southwest and Lower Columbia Regions are Quaternary Period sediments (including  
38 alluvial and coastal deposits), Tertiary Period sedimentary rocks, and Lower Tertiary  
39 volcanic rocks.

40 Due to the limited influence of glacial activity in these regions, soils tend to be older,  
41 deeper, finer in texture, and have a higher nutrient status relative to soils in the northern



1 analysis regions described earlier. Due to these soil characteristics and the generally  
2 favorable climatic conditions, the average potential productivity of covered lands in the  
3 Southwest and Lower Columbia Regions tends to be higher than in other regions.

4 Most non-alluvial soils have formed on parent materials derived from the underlying  
5 bedrock. Topography strongly influences the character and behavior of these soils. On  
6 steeper slopes, soils tend to be shallower, have higher gravel contents and lower potential  
7 productivities relative to soils on more gentle terrain. This is primarily due to the increased  
8 potential for surface erosion and mass wasting on steeper slopes (Washington DNR 1996).

9 **3.4.1.3 Middle and Upper Columbia (Downstream of Grand Coulee Dam)**  
10 **Regions**

11 The Middle and Upper Columbia (Downstream of Grand Coulee Dam) Regions include  
12 portions of the Southern Cascades, Northern Cascades, and Columbia Basin physiographic  
13 provinces as defined by Lasmanis (1991). These Regions largely consist of eastward  
14 trending river valleys that are deeply dissected and separated by sharp ridge crests.  
15 Volcanic rocks of the Columbia River Basalt Group and younger Quaternary volcanics,  
16 including andesite and basalt flows, dominate the geology in the Middle Columbia Region.  
17 Deposits of alpine glacial sediments are also scattered throughout the Region at higher  
18 elevations.

19 Further north, the Upper Columbia Region (downstream of Grand Coulee Dam) lies within  
20 the rugged Northern Cascades where the influence of glaciation is more apparent. All but  
21 the highest peaks in the Region have been heavily glaciated and the valleys of Columbia  
22 River tributaries have relatively flat bottoms and steep walls. Dominant geologic units  
23 include Lower Tertiary Period and Mesozoic Era sedimentary and intrusive igneous rocks.  
24 Both Regions transition into the Columbia Basin physiographic province near their eastern  
25 boundaries where topographic relief decreases. Non-glacial Quaternary Period sediments  
26 and rocks of the Columbia River Basalt Group dominate the geology of the Columbia  
27 Basin.

28 Climatic differences between the analysis regions of western Washington described earlier  
29 and those of eastern Washington have produced substantial differences in soils. The  
30 Middle and Upper Columbia (downstream of Grand Coulee Dam) Regions lie in the rain  
31 shadow of the Cascades and the low precipitation levels limit soil profile development and  
32 forest productivity (Washington DNR 1996).

33 Areas closer to the eastern edges of the analysis regions have the lowest levels of soil  
34 development and potential productivity because they have the lowest average annual  
35 precipitation. Areas in the western portions of the analysis regions tend to have greater soil  
36 development and higher productivity than in the eastern portions because of their higher  
37 average annual precipitation. However, productivity is limited at higher elevations due to  
38 the shorter growing season (Washington DNR 1996).



## Chapter 3

---

### 3.4.1.4 Upper Columbia (Upstream of Grand Coulee Dam) Region

The Upper Columbia (upstream of Grand Coulee Dam) Region roughly coincides with the Okanogan Highlands physiographic province as defined by Lasmanis (1991). The Region is characterized by rounded mountains and deep, narrow valleys. The eastern half of the Region contains the oldest rocks in the State. Dominant geologic units include Paleozoic Era and Precambrian sedimentary rocks and Mesozoic Era intrusive igneous rocks. The western half of the Region was formed by offshore deposition of sediments and volcanic rocks west of the continental margin during the Lower Tertiary Period, Glacial ice covered the Region during the Quaternary Period, reshaping the landscape and forming lakes in the Columbia and Pend Oreille River valleys.

Soil productivity in the Upper Columbia (upstream of Grand Coulee Dam) Region is tied closely to annual precipitation. The areas with higher annual precipitation generally have deeper, more productive soils than drier areas.

### 3.4.1.5 Snake River Region

The Snake River Region includes the entire Blue Mountains physiographic province and part of the Columbia Basin province as defined by Lasmanis (1991). The Blue Mountains are characterized by a broad geologic uplift reaching elevations of more than 6,000 feet above sea level. Rocks of the Columbia River Basalt Group dominate the Region's geology. Quaternary sediments derived largely from glacial outburst floods and eolian deposits are common at lower elevations in the Columbia Basin province.

Many soils in the Blue Mountains have developed from wind-deposited sediments known as loess. These soils tend to be fine-textured and moderately productive. Soil productivity is somewhat influenced by topography, with soils on steeper slopes tending to be more shallow and less productive than soils on gentler terrain.

### 3.4.2 Erosion

Erosion is the detachment and movement of soil particles either individually, in small aggregates, or in large masses (Brooks et al. 1991). The two dominant processes of erosion on forestlands are surface erosion and mass wasting. Surface erosion is the detachment and subsequent removal of soil particles and small aggregates from land surfaces by wind or water. Mass wasting includes erosion in which cohesive masses of soil are displaced. Debris avalanches, debris flows, and debris torrents are mass wasting processes that transport material rapidly. Debris avalanches are shallow, rapid landslides on steep hillslopes (Chatwin et al. 1991). If enough water is present, debris avalanches become debris flows that rapidly transport soil, rocks, and organic material directly to the valley floor and occasionally to stream channels (Chatwin et al. 1991). Where debris avalanches and debris flows reach steep stream channels during peak flow periods, debris torrents occur, involving the rapid movement of large volumes of water-charged soil, rock, and debris (Chatwin et al. 1991). Rates of movement are very high (feet/second) and damage can be extensive. Another class of mass wasting processes includes deep-seated landslides that, unlike debris avalanches, debris flows, and debris torrents, tend to have failure planes greater than 10 feet below the ground surface. Also, deep-seated landslides



1 generally have slower rates of movement (i.e., feet/year to feet/day) although rapid  
2 movement (i.e., feet/minute to feet/second) is possible. Under natural conditions, mass  
3 wasting is the more common form of erosion on forestlands in the Pacific Northwest,  
4 particularly in steep terrain (Sidle et al. 1985).

5 Dam break floods occur when water impounded behind material such as woody debris  
6 and/or coarse sediment breaks through the “dam” releasing a large discharge of water  
7 through the channel. These events can release sediment, scour and temporarily change  
8 channel morphology.

9 The relative importance of different erosion processes on covered lands within the various  
10 analysis regions is summarized below. This information largely reflects findings of the  
11 approximately 60 watershed analyses completed in Washington since 1993.

### 12 **3.4.2.1 Erosion Processes on Covered Lands**

#### 13 **North Puget, South Puget, West Puget, Olympic, and Islands Regions**

14 Mass wasting is the dominant form of erosion in these glaciated regions (Washington DNR  
15 1993, 1994a, 1995a, and 1996b). Steep, mountainous terrain combined with high  
16 precipitation levels produces high rates of mass wasting. Debris avalanches, debris flows,  
17 debris torrents, and dam-break floods are common in most regions (the exception being the  
18 Islands Region) and are most often associated with large magnitude rain or rain-on-snow  
19 events that occur during the fall and winter months (Washington DNR 1998a). Debris  
20 torrents and dam-break floods are usually associated with high-gradient, confined  
21 headwater streams and may translate downstream for several miles (Coho and Burgess  
22 1991). Deep-seated landslides are fairly common and occur most frequently in areas of  
23 structurally weak bedrock or glacial lake deposits (Washington DNR 1998b). Even though  
24 many soils are shallow, high organic matter contents generally produce high infiltration  
25 capacities, so surface erosion is limited to disturbed and/or compacted soils.

#### 26 **Southwest and Lower Columbia Regions**

27 The Southwest and Lower Columbia Regions are similar to those described above in that  
28 mass wasting is the dominant form of erosion (Washington DNR 1994b, 1995b, and  
29 1998c). While topographic relief is low relative to the more heavily glaciated regions to  
30 the north, steep slopes and high precipitation levels produce high rates of mass wasting.  
31 Debris avalanches and debris flows are very common; however, short slope lengths and  
32 low channel gradients make debris torrents and dam-break floods less common. High  
33 frequencies of deep-seated landslides occur in certain geologic units (Washington DNR  
34 1998c). Soils generally have high infiltration capacities, so surface erosion is limited to  
35 disturbed and compacted soils.

#### 36 **Middle and Upper Columbia and Snake River Regions**

37 The lower annual precipitation and more moderate topography east of the Cascade crest  
38 generally results in a lower incidence of mass wasting than on the westside; however,  
39 debris avalanches, debris flows, debris torrents, and dam-break floods all occur to varying  
40 degrees. On the east slopes of the Cascade Range and in the Columbia River Gorge, these  
41 mass wasting processes are usually driven by rain or rain-on-snow events during the fall



## Chapter 3

---

1 and winter months (Washington DNR 1998d, 1998e). In the Okanogan Highlands of  
2 northeastern Washington, rapid spring snowmelt and spring thunderstorms typically  
3 initiate mass wasting (Personal Communication, Jack Powell, Washington DNR, March  
4 2004). Deep-seated landslides are most often associated with interbeds of sediments  
5 between basalt flows of the Columbia River Basalt Group throughout eastern Washington  
6 and glacial lake sediments in the northeastern part of the State (Personal Communication,  
7 Jack Powell, Washington DNR, March 2004). Many soils have low organic matter  
8 contents, making infiltration capacities low relative to soils in western Washington. Also,  
9 relatively low annual precipitation makes vegetation recovery slower than in western  
10 Washington. Relatively low infiltration capacities and longer revegetation periods make  
11 surface erosion fairly common, particularly in areas with disturbed and compacted soils  
12 (Washington DNR 1995c, 1997d).

### 13 **3.4.2.2 Forest Practices Effects on Erosion and Sedimentation**

14 Forest practices such as timber harvesting and road construction have the potential to  
15 accelerate the rate of erosion by disturbing soils, reducing infiltration and increasing  
16 surface runoff (Swanson et al. 1987). Accelerated rates of erosion can lead to increased  
17 sediment delivery to channel networks where it can negatively affect aquatic resources.  
18 Increased sediment supply to streams can reduce the quantity and quality of habitat for  
19 aquatic organisms such as fish, amphibians, and macroinvertebrates (Bisson et al. 1987).  
20 The following discussion provides an overview of various forest practices activities and  
21 their effects on erosion, sediment delivery, and aquatic resources.

#### 22 **Road-Related Surface Erosion**

23 Road-related surface erosion is a function of the intensity of road use; road surfacing  
24 materials, construction, and maintenance types; the intensity and amount of precipitation;  
25 and other factors (Reid and Dunne 1984; Megahan and Kidd 1972). Road construction,  
26 use, maintenance, abandonment, and drainage all play important roles in the production  
27 and delivery of sediment to streams, ponds, lakes, wetlands, and marine areas. Surface  
28 erosion from roads tends to be a chronic source of fine sediment to the drainage network.  
29 Chronic sources of fine sediment can adversely impact the physical habitat of the aquatic  
30 system and certain life stages of fish, aquatic insects and amphibians, and also degrade  
31 water quality.

32 Road density can be used to help understand the potential for impacts from road surface  
33 erosion, drainage, and sediment delivery to streams. As mentioned earlier, many factors  
34 affect the degree of impact to aquatic resources from roads, and there can be a greater  
35 possibility of adverse impacts as road density in a watershed increases. However, the body  
36 of research on this subject indicates the relationship between the degree of impacts to  
37 aquatic resources and road density is not simple and linear. Nonetheless, road density may  
38 be used as an indicator variable. Road densities were estimated within the FPHCP covered  
39 lands for the 12 analysis regions. DEIS Appendix D, Table D.1 shows the average road  
40 density by region and by WRIA, as well as the overall average road density for covered  
41 lands. Estimated average road density in the 12 regions ranged from 2.5 to 4.9 miles of  
42 road per square mile of covered lands with an overall average road density of 3.4 miles per





1 square mile. A description of the process to estimate road density is included with DEIS  
2 Appendix D, Table D.1.

3 Studies of the hydrologic impacts of forest roads indicate that the dominant source of road  
4 runoff is intercepted subsurface storm water flow, where permeable soil intercepts  
5 relatively impermeable bedrock (LaMarche and Lettenmaier 2001; Bowling and  
6 Lettenmaier 2001). Sediment delivery from road surfaces to streams is dependent on  
7 connectivity between streams and road surfaces. Adequate road construction,  
8 maintenance, and abandonment would accommodate most storm events, minimize road  
9 erosion by traffic and precipitation events, and minimize delivery of road surface sediment  
10 to the stream network.

11 Currently, Road Maintenance and Abandonment Plans (RMAPs) are required for large  
12 landowners subject to the Washington Forest Practices Rules. The total road mileage  
13 accounted for under RMAPs as of March 2004 is approximately 40,000 miles.  
14 Approximately 11,000 miles of those are in eastern Washington, and 29,000 miles are in  
15 western Washington. Under current rules, large landowners must have all of their roads  
16 under approved RMAPs by 2006. Small landowners, other than 20-acre exempt  
17 landowners, have the option to submit an RMAP checklist or RMAP when they submit a  
18 forest practices application. The small landowner RMAP checklist submitted with the  
19 forest practices application must cover the haul roads being used under the forest practices  
20 application and does not have to cover other roads the landowners may own. The 20-acre  
21 exempt landowner does not have to submit an RMAP or RMAP checklist. Correction of  
22 all road problems for large landowners must occur by 2016. Correction of road problems  
23 for small landowners generally occurs at the time a landowner files a forest practices  
24 application that includes a road-related problem, and the Washington Forest Practices  
25 Rules require them to maintain roads used for timber hauling in a condition that prevents  
26 damage to public resources. Quantification of the risks of road-related sediment  
27 production and delivery to streams is not completed. However, an adaptive management  
28 research and effectiveness monitoring study is under development to evaluate road  
29 sedimentation under the current Washington Forest Practices Rules.

### 30 **Road-Related Mass Wasting**

31 Debris avalanches, debris flows, and debris torrents are episodic sources of fine and coarse  
32 sediment to the aquatic system. These erosion processes can be greatly accelerated by road  
33 management practices. Many studies have shown that on a unit area basis, roads have the  
34 greatest effect on mass wasting of all activities on forestlands (Sidle et al. 1985).  
35 However, some research suggests that harvest-related mass wasting occurs with  
36 approximately the same frequency as road-related landslides (Montgomery et al. 1998).  
37 Road location, drainage, design, construction, and maintenance are all-important factors in  
38 effective road design, but can be contributing factors to road-related failure. New road  
39 construction and engineering design has reduced road-related mass wasting relative to  
40 roads constructed more than 15 to 20 years ago (Toth 1991; Robison et al. 1999).



## Chapter 3

---

### 1 **Orphan Roads**

2 Orphan roads are roads constructed prior to 1974 that have not been used for forest  
3 practices since that time (WAC 222-24-052\*(4)). Such roads are typically not maintained,  
4 and many were constructed without a requirement to consider public resource and channel  
5 impacts. The mileage of orphan roads in Washington is unknown; however, the associated  
6 hazards have been identified. The concern with orphan roads is the lack of knowledge  
7 about their location and potential for failure and initiation of debris avalanches, debris  
8 flows, and debris torrents. Large landowners are inventorying and assessing their orphan  
9 roads and showing the orphan roads' locations in their RMAPs. The goal of the orphan  
10 roads inventory is to locate the roads and assess the condition of the roads to determine  
11 what work needs to be done to protect resources from orphan road failures and how the  
12 cost of that work might be funded.

### 13 **Harvest-Related Surface Erosion**

14 Surface erosion is dependent on many variables. The primary variables are slope, soil  
15 texture, and vegetation cover (Benda et al. 1998).

16 Harvest activities such as ground skidding or cable yarding can cause some degree of soil  
17 disturbance. Typically, ground-based systems compact and disturb more soils than non-  
18 ground-based harvest systems (Graham et al. 1991). The harvest systems most likely to  
19 cause greater levels of disturbance (from greatest to lowest) are ground-based systems,  
20 cable yarding, and aerial systems (Beschta and Boyle 1995). Clearcuts tend to create the  
21 greatest area of soil disturbance (Hermann 1978); however, disturbance from felling,  
22 yarding, and skid trails in partial cuts can also cause ground disturbance and compaction.  
23 Cromack et al. (1978) found levels of soil disturbance in clearcut and partial cut areas to be  
24 comparable because of the need for equivalent access through a harvest unit.

25 Streamside buffers can substantially reduce fine sediment that is transported overland  
26 (Rashin et al. 1999). The capacity of riparian buffers to control sediment inputs from  
27 surface erosion depends on several site characteristics including the presence of vegetation  
28 or organic litter, slope, soil type, and drainage characteristics. Additionally, the filtering  
29 capacity is affected by timber harvest activities within the buffer. Although soil  
30 disturbance generally increases the sediment delivery potential, the addition of obstructions  
31 on the forest floor from tree limbs and boles (limb-free trunks) associated with partial  
32 logging can offset diminished filtration (Burroughs and King 1989; Benoit 1979). These  
33 factors influence the ability of buffers to trap sediment by controlling the infiltration rate of  
34 water and the velocity of overland flow.

35 Riparian protection measures should also include practices for minimizing sediment  
36 contributions from outside the riparian area. Timber harvest activities can disturb the  
37 upper soil layers, exposing the subsoil to erosion. A study of sediment delivery from  
38 timber harvest was conducted by Rashin et al. (1999) between 1992 and 1995, prior to the  
39 current Washington Forest Practices Rules. This study evaluated specific harvest practices  
40 on State and private lands across Washington. Harvest sites were evaluated for soil  
41 exposed and sediment delivered, and was categorized by a number of parameters, including  
42 harvest method and distance from streams. They found that in areas where there were no



1 buffers, best management practices (BMPs) for timber harvest were not effective in  
2 preventing soil disturbance or preventing sediment from reaching streams. They also  
3 found that no-harvest buffers at least 30 feet wide were effective in filtering sediment,  
4 although they caution use of these results because of low precipitation and storm events  
5 during the study period.

### 6 **Harvest-Related Mass Wasting**

#### 7 **Debris Avalanches and Debris Flows**

8 Debris avalanches and debris flows are the most common types of landslides in steep  
9 forestlands. Still, only 2 percent of the landscape is affected at any given time (Ketcheson  
10 and Froelich 1978; Ice 1985). Debris avalanches and debris flows are typically initiated by  
11 high magnitude rain or rain-on-snow events during the fall and winter months (Swanson et  
12 al. 1987). Sidle et al. (1985) summarized several studies (Swanston 1970, 1974;  
13 O’Loughlin 1974; Ziemer and Swanston 1977; Burroughs and Thomas 1977; Gray and  
14 Megahan 1981; Ziemer 1981) indicating that slope stability depends partly on  
15 reinforcement from tree roots, especially when soils are partly or completely saturated.  
16 Clearcut timber harvesting on unstable slopes or landforms decreases rooting strength,  
17 increasing the potential frequency and magnitude of debris avalanches and debris flows  
18 (Ziemer and Swanston 1977; Wu and McKinnell III 1978).

#### 19 **Debris Torrents**

20 Debris avalanches and debris flows can turn into debris torrents. Debris torrents usually  
21 transport more material than the initiating event, due to scouring action on the slope or in  
22 the channel. Debris torrents stop moving when the channel gradient decreases  
23 substantially or when the torrent encounters a sharp bend in the channel. Debris torrents  
24 contain substantial amounts of wood and can travel varying distances, which can result in  
25 variable degrees of impact depending upon channel gradient, confinement, layout of the  
26 channel network, and other characteristics (Fannin and Rollerson 1993). Debris torrents  
27 can have long-lasting effects on stream channels. The channel location and cross-section  
28 can be radically altered in such a way that normal flows and normal peak flows cannot  
29 reconfigure the channel easily (Lamberti et al. 1991). This is important because even  
30 though mass wasting in general may affect only 1 percent of a watershed, debris torrents  
31 can affect up to 10 percent of the stream system because of their mobility (Swanson et al.  
32 1987). These channel alterations from debris torrents, however, are not always negative.  
33 For example, Benda et al. (2003) found that channel morphology and habitat complexity  
34 (e.g., pool density, substrate texture, and channel widths) increased in proximity to low-  
35 order (See Glossary) tributary confluences prone to debris flows. In addition to having  
36 impacts on the stream channel, debris torrents can also affect riparian buffer functions and  
37 streamside forests when bank scour is so great that streamside vegetation is removed.

#### 38 **Streambank Stability**

39 The roots of riparian vegetation help bind soil together, making soils more resistant to  
40 erosion and slope failure (Wu and Sidle 1995). The stability of streambanks is largely  
41 determined by the size, type and cohesion of bank material, vegetation cover, and the  
42 amount of bedload carried by the channel (Sullivan et al. 1987). Riparian vegetation can



## Chapter 3

---

1 also provide hydraulic roughness elements that dissipate stream energy during high or  
2 overbank flows, which further reduces bank erosion. In most cases, vegetation  
3 immediately adjacent to a stream channel is most important in maintaining bank integrity  
4 (FEMAT 1993); however, in wide valleys with migrating channels, floodplain vegetation  
5 may be important in minimizing bank erosion over longer time periods.

### 6 **3.4.2.3 History of Forest Practices Affecting Erosion and Sedimentation**

7 Prior to the adoption of the Washington Forest Practices Act in 1974, there were few rules  
8 or regulations that governed timber management activities on State and private forestlands.  
9 During the early years of logging in the Pacific Northwest, stream and river channels were  
10 used to move logs downstream to accumulation sites. Most streams of sufficient size in  
11 western Oregon and Washington were cleared of obstructions for log drives during high  
12 water (Sedell et al. 1991). On streams too small for log drives, splash dams of stacked log  
13 cribs were used to raise a head of water for sluicing (transporting with water) logs (Sedell  
14 and Luchessa 1982). By about 1900, more than 300 major splash dams and many  
15 undocumented small dams operated in Oregon and Washington.

16 During the first part of the 20<sup>th</sup> century, railroads were constructed along large channels,  
17 and logs were yarded down the small tributaries to the rail bed. In this way, impacts  
18 extended to headwater channels. Whole watersheds were logged as convenience,  
19 beginning in the lower watershed and progressing upstream until all valuable timber was  
20 extracted. Logs were yarded downhill, moving debris and sediment into stream channels.  
21 Streams were protected from being used for yarding beginning in the 1950s, but clearcut  
22 harvesting to the streambank was commonplace until the adoption of the Forest Practices  
23 Act in 1974.

24 The construction of an extensive network of logging roads began with the advent of the log  
25 truck. Beginning in the 1930s and continuing through the 1970s, many roads were  
26 constructed across unstable slopes with little regard for mass wasting potential. Operators  
27 typically employed cut-and-fill construction techniques that involved sidecasting of excess  
28 fill material. Such approaches resulted in the construction of hundreds of miles of unstable  
29 roads throughout Washington. Once constructed, few roads were properly maintained to  
30 minimize surface erosion and mass wasting.

31 Like logging road construction, early timber harvest operations were conducted without  
32 consideration of impacts to unstable slopes. In addition, the development of road networks  
33 beginning in the 1950s allowed harvesting to occur in many areas that were formerly  
34 inaccessible by rail. Clearcut harvesting of unstable landforms increased the rate of  
35 management-related mass wasting in many areas prior to adoption of Washington Forest  
36 Practices Rules protecting unstable slopes in 1987.

37 The Forest Practices Act of 1974 was the first comprehensive law aimed at minimizing the  
38 effects of forest practices on aquatic resources in Washington. The Act established the  
39 State's Forest Practices Board and directed the agency to adopt operating standards and  
40 rules for activities on State and private forestlands. These first rules focused on  
41 minimizing sediment delivery and controlling stream temperature increases by requiring



1 the retention of riparian buffers along large streams. In 1987, buffer requirements were  
2 expanded in light of new information related to the importance of instream large woody  
3 debris (LWD). The 1987 rules also granted the Washington DNR regulatory authority  
4 over operations on unstable slopes and formally recognized the collaborative Timber, Fish,  
5 and Wildlife (TFW) forum as the preferred approach to solving forest resource  
6 management problems. Washington Forest Practices Rules related to aquatic resources  
7 were again revised in 1992 when the Forest Practices Board endorsed Watershed Analysis  
8 as a way of addressing cumulative watershed effects involving sediment, water, woody  
9 debris, temperature, and hydrology.

10 The most recent revision of Washington Forest Practices Rules occurred in 2001 when the  
11 Forest Practices Board adopted sweeping changes based on the 1999 Forests and Fish  
12 Report (FFR) as the current Washington Forest Practices Rules. Compared to the 1974  
13 rules, the current Washington Forests Practices Rules include several mechanisms to  
14 further reduce sediment-related impacts to aquatic resources. These mechanisms include:  
15 1) better screening and identification of unstable slopes and landforms during the forest  
16 practices application review process, 2) more rigorous review of operations proposed on  
17 unstable slopes and landforms (including review under the State Environmental Policy  
18 Act), 3) expanded buffer requirements for fish-bearing waters, 4) Equipment Limitation  
19 Zones adjacent to non-fish-bearing waters to minimize soil disturbance and sediment  
20 delivery, and 5) increased road maintenance and abandonment planning and  
21 implementation requirements.

22 Many of the sediment minimization prescriptions in the current Washington Forest  
23 Practices Rules arose from information learned through watershed analyses conducted  
24 across the State beginning in 1992. More than 60 watershed analyses have been conducted  
25 in Washington; however, the majority of watersheds in the State have not undergone  
26 analysis, partially because the current Washington Forest Practices Rules were derived  
27 from effective Watershed Analysis prescriptions. Once the Washington Forest Practices  
28 Rules were adopted, the need for new watershed analyses decreased. Nevertheless, two  
29 modules within Watershed Analysis, mass wasting and surface erosion, provided  
30 substantial landscape-scale data related to forest practices effects on landslide processes  
31 and road and hillslope erosion. These data were gathered from more than 60 different  
32 Watershed Administrative Units that include over 3,000 square miles of forestland.  
33 Additional information related to the effectiveness of forestry BMPs in controlling  
34 sediment delivery came from Rashin et al. (1999). In the study, the authors evaluated  
35 timber harvesting, road construction, and road maintenance practices in light of water  
36 quality standards. The report included recommendations for improvement where practices  
37 were determined to be ineffective.

38 Current Washington Forest Practices Rules intended to reduce erosion and sedimentation  
39 will be evaluated through adaptive management. Under the rules, monitoring projects are  
40 currently being developed that will evaluate the effectiveness of unstable slopes protection  
41 measures and implementation of RMAPs. In addition, two adaptive management projects  
42 are already underway to identify and improve the detection of unstable slopes and



## Chapter 3

---


1 landforms (i.e., the Regional Landform Identification Project and the Landslide Hazard  
2 Zonation Project) (CMER 2004b).

3 Under current Washington Forest Practices Rules, when harvest is proposed on potentially  
4 unstable slopes, one of three outcomes usually occurs (Personal Communication, Jeff  
5 Grizzel, Washington DNR, April 7, 2004). If the landowner recognizes he/she is  
6 proposing harvest or construction on an unstable slope or landform before submitting the  
7 forest practices application, then he/she hires a qualified (licensed) individual, as defined in  
8 the Washington Forest Practices Rules, who conducts the assessment and writes a report.  
9 The report is then submitted with the forest practices application. The report must include  
10 an analysis of the risks and any mitigation involving the potentially unstable slope or  
11 landform. Once the Washington DNR confirms the proposal includes harvest or  
12 construction on an unstable slope/landform through field review, the forest practices  
13 application is classified as Class IV-Special. If a landowner does not recognize he/she is  
14 proposing harvest or construction on an unstable slope or landform and submits the forest  
15 practices application to the Washington DNR, then the Washington DNR initially classifies  
16 the forest practices application as a Class II or III.

17 The Washington DNR screens the forest practices application for unstable slope presence  
18 using GIS-based screening tools. The Washington DNR forest practices forester relies on  
19 his/her personal knowledge of the project area along with the results of the GIS-screen to  
20 assess the need for field review. If a field review is conducted and the forester discovers an  
21 unstable slope/landform, the forest practices application is disapproved, reclassified as a  
22 Class IV-Special, and is sent back to the landowner letting him/her know that the forest  
23 practices application would be a Class IV-Special if it is resubmitted without modification.  
24 The Washington DNR also explains that a qualified (licensed) individual is required to  
25 conduct a geotechnical assessment before resubmitting the forest practices application. In  
26 some instances in this second scenario, the landowner will modify his/her proposal so the  
27 operation avoids the unstable slope/landform, thereby eliminating the need for the  
28 geotechnical assessment. For landowners, especially small landowners, the need to have a  
29 geotechnical assessment and potentially design mitigation is generally a disincentive to  
30 harvest on unstable landforms and potentially unstable slopes.

31 Therefore, the current Washington Forest Practices Rules provide a means for identifying  
32 unstable landforms and potentially unstable slopes, and mitigating for those effects. The  
33 process of identification may lead to elimination of potentially unstable areas from the  
34 proposed harvest. In addition, the process of identification of unstable landforms and  
35 potentially unstable slopes can be refined over time under adaptive management to be more  
36 efficient and certain.

37



## 1 **3.5 WATER RESOURCES**

2 Wet Pacific weather systems combined with the rain shadow effect produced by the  
3 Cascade Mountains, produce heavy rains on the western slopes of the Cascades and drier  
4 conditions east of the Cascades. As a result, a myriad of surface water conditions occur in  
5 Washington State. Literally all forested lands in Washington have distinct surface water  
6 features, ranging from small, intermittent streams to the very large Columbia and Snake  
7 Rivers. Most of these rivers and streams support complex aquatic ecosystems, including  
8 stocks of endangered Pacific salmon and numerous other aquatic communities. Many of  
9 the major rivers and streams on the westside of the Cascades and the eastside of the  
10 Olympics drain into Puget Sound, a complex and valuable marine resource to Washington  
11 State.

12 This section is divided into three subsections: Surface Water Quality (subsection 3.5.1),  
13 Surface Water Quantity (subsection 3.5.2), and Groundwater Quantity and Quality  
14 (subsection 3.5.3). Specific information related to water resources by analysis region can  
15 be found in DEIS Appendix A.

### 16 **3.5.1 Surface Water Quality**

#### 17 **3.5.1.1 Introduction**

18 Water quality is measured by many parameters. The physical properties and chemical  
19 constituents of water serve as the primary means for monitoring and evaluating water  
20 quality. Forest practices have the greatest potential effect on the following water  
21 parameters: 1) stream water temperature, 2) sediment-related water quality parameters  
22 such as turbidity, 3) dissolved oxygen, 4) pesticides/herbicides, and 5) nutrients. The  
23 Washington Forest Practices Rules must comply with the Clean Water Act to meet State  
24 water quality standards for surface waters and groundwater. Moreover, they must provide  
25 for adequate water quality protection for fish and wildlife habitat. This subsection briefly  
26 describes the issue of water quality and the current water quality status of lands subject to  
27 Washington Forest Practices Rules.

#### 28 **3.5.1.2 Water Quality Parameters for Surface Waters**

##### 29 **Temperature**

30 Stream temperature is influenced by many factors including latitude, altitude, season, time  
31 of day, flow, channel width and depth, groundwater flow, stream shading from topography  
32 or vegetation, and coastal fog (MacDonald et al. 1991). Temperature plays an integral role  
33 in the biological productivity of streams. Aquatic life can be very sensitive to water  
34 temperature fluctuations. Because salmonids and some amphibians are sensitive to water  
35 temperatures, they are often used as indicator species regarding water temperature and  
36 water quality. Salmonid temperature requirements can vary by species and life stage  
37 (Bjornn and Reiser 1991; Hicks 2002; Environmental Protection Agency 2003). However,  
38 in general, juvenile salmon and trout may be susceptible to sublethal adverse effects when  
39 the average stream temperature is above about 59°F (15°C) (Hicks 2002). Bull trout,  
40 especially juveniles, may be susceptible when average temperatures are greater than about  
41 50°F (10°C). The upper lethal temperature for salmonids common in the Pacific



## Chapter 3

---

1 Northwest ranges from 73 to 79°F (23 to 26°C) (Bjornn and Reiser 1991). The preferred  
2 range for most salmon and trout is 54 to 57°F (12 to 14°C) (Bjornn and Reiser 1991).  
3 However, available information suggests that optimal temperatures for bull trout juvenile  
4 growth and rearing are below 59°F (15°C) and probably closer to 50 to 54°F (10 to 12°C)  
5 (Buchannan and Gregory 1997; Oregon Department of Environmental Quality 1995).

6 Stream water temperature is regulated by heat exchange between the stream water and the  
7 aerial and subsurface conditions. Heat energy is transferred to and from streams by direct  
8 solar radiation (short wave), long-wave radiation, convective mixing with air, evaporation,  
9 conduction with the streambed, and advective mixing with inflow from groundwater or  
10 tributary streams (Beschta et al. 1987; Sullivan et al. 1990). Streams exhibit a natural  
11 warming trend as water flows from headwaters to the sea (Sullivan et al. 1990). However,  
12 changes in environmental conditions along a reach can modify temperatures beyond the  
13 normal trend (Johnson and Jones 2000). In small- to intermediate-size streams of forested  
14 regions, incoming solar radiation represents the dominant form of energy input to streams  
15 during the summer, with convection, conduction, evaporation, and advection playing  
16 relatively minor roles (Brown 1980; Beschta et al. 1987; Sullivan et al. 1990). In larger  
17 streams, the effects of riparian shading and advective mixing diminish and, as a result,  
18 evaporative heat-loss processes increase. In small streams, groundwater discharge may  
19 also be important where it provides a large percentage of the overall discharge, particularly  
20 in the summer months during low flows.

21 Brosofske et al. (1997) suggested that groundwater and stream temperatures could increase  
22 due to heating of upslope soils in clearcuts. In their study, stream temperatures were  
23 correlated with shallow (4 inches) upslope soil temperatures. However, the Brosofske et  
24 al. (1997) study was focused on microclimate gradients in riparian zones rather than water  
25 heating and watershed hydrology; no measurements of interflow (horizontal movements of  
26 water above the water table) and groundwater temperatures were taken. St. Hilaire et al.  
27 (2000) incorporated interflow in their mechanistic stream-heating model. Their unverified  
28 modeling predictions suggested that less than a 1°F (0.4°C) increase would occur during a  
29 tropical (warm front) storm if 50 percent of the watershed was harvested. Overall, the  
30 magnitude of effects of upslope clearcuts on stream temperatures, if any, is uncertain.  
31 More detailed information pertaining to water temperature concerns by analysis region is  
32 available in DEIS Appendix A.

### 33 **Sediment**

34 Two of the most common water quality parameters measured and monitored for sediment  
35 are suspended sediment and turbidity. Both are related to sediment delivery and transport  
36 in hydrologic systems. Streams that exceed water quality objectives for sediment would  
37 have high suspended-sediment delivery rates and/or turbidity. Suspended sediment is the  
38 portion of the sediment load suspended in the water column. The grain size of suspended  
39 sediment is usually less than 1 mm in diameter (Sullivan et al. 1987). Turbidity refers to  
40 the amount of light scattered or absorbed by a fluid and is measured in nephelometric  
41 turbidity units (NTUs). In streams, turbidity is usually a result of suspended particles of





1 silts and clay, but also organic matter, colored organic compounds, plankton, and  
2 microorganisms.

3 Biological effects of increased turbidity may include a decrease in primary productivity of  
4 algae and periphyton due to the decrease in light penetration. Declines in primary  
5 productivity can adversely affect the productivity of higher trophic levels such as  
6 macroinvertebrates and fish (Gregory et al. 1987). Siltation and turbidity have also been  
7 shown to affect fish adversely at every stage in their life cycle (Iwamoto et al. 1978).  
8 Deposited sediments tend to have a greater impact on fish than suspended sediment;  
9 spawning and incubation habitats are most directly affected (Spence et al. 1996).  
10 However, suspended sediment can settle out of suspension in the water column and  
11 become part of the bedload (i.e., sediment carried along the bed of the stream). DEIS  
12 Appendix A provides a summary of currently available information on sedimentation  
13 impacts by analysis region. See also subsections 3.8.3.1 (Coarse Sediment) and 3.8.3.2  
14 (Fine Sediment) for a discussion of the attributes of sediment relative to the aquatic  
15 ecosystem.

### 16 **Dissolved Oxygen**

17 High water temperatures can contribute to low dissolved oxygen because warm water  
18 cannot hold as much oxygen in solution as cold water can. Dissolved oxygen is often  
19 lower in surface waters that have a large percentage of their volume coming from poorly  
20 oxygenated groundwater, especially during summer low flows. Excess nutrients,  
21 especially phosphorus, from sediments, fertilization or waste products, can stimulate algae  
22 and aquatic plant growth in surface water. When dead plant material decays, and when  
23 plants are taking up oxygen at night, dissolved oxygen levels decline.

24 Salmon (all life stages) and other aquatic life need sufficient levels of dissolved oxygen to  
25 survive. As temperature increases, salmon metabolism increases and the demand for  
26 oxygen also increases. For maintenance of the health of salmonids and many other aquatic  
27 organisms, levels of dissolved oxygen concentrations in the water should approach  
28 saturation (Deas and Orlob 1999). Water with dissolved oxygen in the range of 8 to 9  
29 milligrams per liter is needed to ensure that the normal physiological functions of  
30 salmonids are not impaired (Spence et al. 1996).

### 31 **Pesticides**

32 Pesticides used in forest management include a wide variety of chemicals introduced to the  
33 forest environment with the intent of controlling or halting the proliferation of nuisance  
34 organisms. Pesticides are commonly grouped according to one of three target organisms:  
35 plants (herbicides), insects (insecticides), and fungi (fungicides). In general, pesticide  
36 application rates on forested lands are fairly infrequent, with roughly one to two  
37 applications every 40 to 60 years (Washington Department of Ecology 1993a). The active  
38 ingredients usually determine the effects of individual pesticides. In addition, prior to  
39 application, most pesticides are combined with a surfactant (i.e., a surface-active agent) or  
40 other adjuvant (i.e., a pharmacological agent added to increase or aid the pesticide's effect)  
41 to control and improve the desired effect. Although these additives typically present lesser  
42 threats to the environment than the active ingredients in the pesticides, their impacts can be



## Chapter 3

---

1 important, and in some cases the impacts are greater than those associated with the active  
2 ingredients.

3 Pesticides used in the forest environment can become water contaminants if they are  
4 transported to surface waters or groundwater. Transportation to surface waters would most  
5 likely occur through wind drift; however, heavy rains can result in pesticide transport in  
6 stormwater runoff or through contaminated soil erosion. Pesticides can also be directly  
7 applied to surface waters by overspray and spills. Groundwater contamination can occur  
8 through contaminated surface water recharge and through the direct transport of pesticides  
9 from the soil surface by rainwater. Most pesticides that have been detected in streams and  
10 groundwater are present at very low concentrations, usually well below regulatory drinking  
11 water criteria (U.S. Geological Survey 1996a, 1996b, 1996c, 1997b). However, some  
12 pesticides have been detected at concentrations that exceed the more restrictive guidelines  
13 for the protection of aquatic life (freshwater chronic criteria) or health advisories for  
14 drinking water (U.S. Geological Survey 1996c; Washington Department of Ecology  
15 1993a). Although studies focused specifically on forestry applications have found  
16 violations of applicable water quality standards resulting from chemical applications, these  
17 violations usually resulted from the lack of spray buffers or from applications over dry or  
18 ephemeral streams (Neary and Michael 1996; Washington Department of Ecology 1993a).  
19 Finally, although low levels of pesticide contamination in surface water and groundwater  
20 have been found throughout Washington State, the source of the contamination (e.g., forest  
21 applications, agriculture, urban activity) is difficult to identify and cannot be linked  
22 directly to forest applications, unless no other possible sources exist.

23 Bortleson and Ebbert (2000) reported the occurrence of pesticides in streams and  
24 groundwater in the Puget Sound basin and found more detections and higher  
25 concentrations of distinct compounds in surface water samples during storms than in  
26 groundwater. Groundwater concentrations of pesticides decreased with increasing depth of  
27 well, and compounds were linked to urban and agricultural land use. None of their  
28 detections in groundwater exceeded water quality standards.

### 29 **Nutrients**

30 Inorganic nutrients commonly associated with forest practice activities, such as nitrogen or  
31 phosphorus, may show moderate increases following timber harvest although levels are  
32 typically limited to no more than 10 years following harvest (Hicks et al. 1991; Chamberlin  
33 et al. 1991). Urea is a commonly used forest fertilizer in the Pacific Northwest and rapidly  
34 hydrolyzes to ammonium, which gradually oxidizes to nitrate. The nitrate may leach out  
35 of the soil into streams (Binkley et al. 1999). Water quality criteria for nitrogen are rarely  
36 exceeded, but the criteria are based on human health (nitrates) and fish toxicity (ammonia)  
37 rather than ecological disturbances (Anderson 2002). Forest fertilizers, especially when  
38 combined with high water temperatures, have the potential to result in algal blooms  
39 (Chamberlin et al. 1991). Algal response to additional nutrients could be greatest in  
40 downstream reaches (Anderson 2002). Excessive algal growth can lead to high pH from  
41 photosynthesis (Erickson 1999) and to low oxygen from decay of dead algae (Roberts et al.  
42 2004). High pH and low dissolved oxygen can be harmful to salmon (Bell 1991).



### 3.5.1.3 Regulatory Background

The Washington Forest Practices Rules must comply with the Clean Water Act to meet State water quality standards for surface waters and groundwater (Table 3-14). Water quality standards are set to provide for the protection of beneficial uses such as public water supplies, aquatic habitat, and recreation. The Forest Practices Act of 1974 authorizes the adoption of regulations establishing water quality standards for forest practices, and ESHB 2091 (1999) changed Ecology’s role; the Forest Practices Board must now reach agreement with Ecology’s director (or designee) prior to adopting Washington Forest Practices Rules related to water quality protection.

**Table 3-14.** Washington State Water Quality Standards for the Major Non-Chemical Parameters of Concern.<sup>1/</sup>

Water Quality Parameter	Washington State Standard (Class AA, Excellent)	Washington State Standard (Class A, Good)
Temperature	Shall not exceed 16.0°C (61°F) due to human activities. When natural conditions exceed 16°C (61°F), no temperature increase greater than 0.3°C (0.5°F) is allowed. Incremental temperature changes from nonpoint source activities shall not exceed 2.8°C (5°F).	Shall not exceed 18.0°C (64°F) due to human activities. When natural conditions exceed 18°C (64°F), no temperature increase greater than 0.3°C (0.5°F) is allowed. Incremental temperature changes from nonpoint source activities shall not exceed 2.8°C (5°F).
Sediment	Per WAC 173-201A-200(e)(i), implementation of Best Management Practices (BMPs) will meet water quality criteria and support characteristic water uses. BMPs are described in Forest Practices Board Manual Section 3, Guidelines for Forest Roads.	Same as Class AA.
Turbidity <sup>2/</sup>	Shall not exceed 5 NTU (nephelometric turbidity units) over background when the background level is 50 NTU or less, nor increase more than 10% of background when the background level is 50 NTU or more.	Same as Class AA.
Dissolved Oxygen	Freshwater - dissolved oxygen shall exceed 9.5 mg/L.	Freshwater – dissolved oxygen shall exceed 8.0 mg/L.
All Parameters	Whenever waters are of a higher quality than the assigned criteria, actions reducing water quality shall not be allowed except as described in WAC 173-201A-070 (1997).	Same as Class AA.

<sup>1/</sup> New water quality standards have been adopted and are awaiting approval by EPA. The new standards for temperature would generally be lower and more specific to fish populations (Hicks 2003).

<sup>2/</sup> Nephelometric turbidity units are the measurement units of turbidity using a nephelometer (light reflected by particles in suspension at a right angle to the original source).

Source: WAC 173-210A.



## Chapter 3

---

### 3.5.1.4 Existing Water Quality

The primary water quality problem on forestlands throughout the State is temperature, which also happens to be the most prominent water quality problem for the State's water bodies (DEIS Appendix A). There is no readily available specific information on the number of impaired water bodies on forestlands throughout the State. Although Ecology has compiled 303(d) listings for the State, land use associated with impairments are not included with the information about water quality parameters in violation of State standards. However, elevated water temperature generally occurs in areas where riparian vegetation has been removed or reduced, thereby reducing shade levels, which can affect water temperature. Other problems include erosion from road building, construction, and agriculture, which increases sediment in streams. Increased heat and sediment and reduced flow can also lower dissolved oxygen.

Ecology has prepared a draft Water Quality Assessment of Washington's Surface Waters (Washington Department of Ecology 2004). One category of the list is "polluted waters that require a TMDL (Total Maximum Daily Load)," also known as the "303(d) list" after Section 303(d) of the Federal Clean Water Act. As mentioned in Chapter 1 (Purpose and Need), a TMDL is the maximum amount of pollution or "pollutant load" that a water body can assimilate without violating water quality standards. The draft 303(d) list includes over 700 freshwater segments that have been identified as impaired due to high temperature on over 400 streams statewide. This is approximately a 30 percent increase over the number of segments listed for temperature in 1998, probably due to additional monitoring and reporting rather than an actual increase in the number of impaired waters. Some of these streams may be naturally elevated for temperature; however, that determination will not be made until a TMDL is developed for the stream. Even with this additional monitoring, Ecology does not have data for the majority of streams in Washington. Ecology's 303(d) list also does not differentiate between land uses; therefore, it is difficult to determine how many of the listings are related to forest management and how many are a result of other land uses, such as urban development or agriculture, or due to natural conditions.

For other parameters, there are over 200 freshwater segments listed for dissolved oxygen, and several listings each for turbidity and fine sediment. These numbers are similar to the numbers of listings in 1998. An analysis of 303(d) 1998 data for forested watersheds with greater than 5 percent State ownership or management indicated that dissolved oxygen and fine sediment impacted far fewer such watersheds than did temperature. Many of the elevated values for dissolved oxygen can be linked to temperature effects (dissolved oxygen decreases with increasing temperature) (Washington Department of Ecology 2004).

Segments impaired due to low instream flow are included in a separate category, "impaired by a non-pollutant," and do not require TMDLs. The effect of nutrient input on surface waters as a function of land use is illustrated by the findings of a U.S. Geological Survey study of nutrient transport (U.S. Geological Survey 2000). This study reported that forested watersheds in the Puget Sound Region have much smaller yields of organic nitrogen and nutrients than do watersheds that are urbanized or have agricultural land uses,



1 due to higher inputs from these developed areas. Phosphorous yields were not related to  
2 phosphorous inputs, however, because phosphorous adsorbs to soil particles.

### 3 **3.5.1.5 Water Quality Conditions by Analysis Region**

4 DEIS Appendix A provides a summary of physical conditions of streams within each  
5 analysis region. Although causes of pollutants and impairments is unknown in many  
6 instances, common factors such as temperature, turbidity, and dissolved oxygen, as well as  
7 physical impairments to instream flow and fish habitat are commonly noted, to varying  
8 degrees, in all of the regions analyzed.

9 A common factor for each analysis region is that past timber harvests generally have  
10 resulted in too little riparian vegetation being retained along streams (Kuttel 2001, 2002;  
11 Haring 2000; Correa 2002), thereby reducing down woody debris recruitment at least in  
12 the near-term, as well as shade, which can affect water temperature. Temperature concerns  
13 caused by loss of streamside vegetation are the greatest for medium width streams as  
14 opposed to small or large streams. This is because recovery from past harvests tends to be  
15 rapid in small stream channels, because small trees can provide adequate shade. Similarly,  
16 temperature impacts from riparian harvest along large streams or rivers are less substantial,  
17 because even under natural conditions, large streams or rivers are typically only partially  
18 shaded by riparian trees. However, trees do make a difference on large streams and rivers;  
19 thus, temperature recovery from riparian timber harvest takes longer on large streams  
20 (Washington State Conservation Commission 1999a). Sullivan et al. (1990) found that a  
21 loss of riparian trees will increase water temperature where the open channel is less than  
22 108 feet (33 meters) wide.

23 Another factor attributable to historic timber harvest on western Washington forestlands is  
24 that many riparian buffers have regenerated as hardwood-dominated stands (i.e., greater  
25 than 70 percent hardwoods) (Marshall and Assoc. 2000), with most of this being red alder.  
26 Because alder has a short life span (80 years), limited height (50 to 90 feet depending on  
27 soil and climate) and size potential, and lacks the foliage density of conifers, it is less  
28 effective in providing LWD or shading to wider channels. Also, alder generally has less  
29 longevity as LWD in streams than coniferous LWD (Marshall and Assoc. 2000).

### 30 **3.5.2 Surface Water Quantity**

31 Three primary factors affect surface water quantity in forested watersheds:

- 32 • Climate: Precipitation amount and form (snow or rain) determine the rates of delivery  
33 of water to a watershed. These processes are largely controlled by climate.
- 34 • Vegetation: Interception, condensation, evapotranspiration, and canopy snowmelt  
35 influence delivery of water to the forest floor. These processes are controlled mainly  
36 by vegetation.
- 37 • Transport Pathways: Surface and subsurface pathways transport runoff from the forest  
38 floor to the streams. These pathways are controlled by the interaction of condensation,  
39 precipitation, evapotranspiration, interception, snowmelt, and other physical and  
40 biological factors.



## Chapter 3

---

1 The hydrologic functions of a watershed are dependent upon these processes. When these  
2 processes are individually or cumulatively altered by road construction, harvesting, or  
3 other forest practices, the hydrologic continuity of the watershed is altered (Montgomery  
4 1994; Rashin et al. 1999; USDA Forest Service 2001). Three major areas of hydrologic  
5 concern-annual water yields, low flows, and peak flows-are discussed in this subsection.

### 6 **3.5.2.1 Areas of Concern–Surface Water Quantity**

#### 7 **Water Yield (Annual)**

8 Water yield is the amount of water that is transported from a watershed. Various studies  
9 (Helvey 1980; Bosch and Hewlett 1982; Harr 1983; Kattlemann et al. 1983; Troendle  
10 1983; King and Tennyson 1984; Trimble and Weirich 1987; Keppeler and Ziemer 1990)  
11 have shown short-term increases in water yields following timber harvest. However, the  
12 increase in water yield is usually beneficial to the aquatic system (unless it results in  
13 increased peak flows - see below) and will not be addressed further in this subsection.  
14 Although in general, forests act to lower average stream flows, forests may also reduce  
15 peak flows and increase flows during dry seasons. This is because forested lands tend to  
16 have better infiltration capacity and a high capacity to retain water than nonforested lands  
17 (Jones and Grant 1996; Intergovernmental Panel on Climate Change 2003).

#### 18 **Low Flows**

19 Low flows are often referred to as base flows, dry-weather flows, and groundwater flows.  
20 Low flows are the flows provided by groundwater to the streams during the lowest  
21 precipitation months of the year in the summer. In western Oregon, increases in low flow  
22 are generally short-term (5 years) following clearcut timber harvest (Rothacher 1970; Harr  
23 et al. 1982). In a northern California study, summer low flows were increased following  
24 selection harvest and then declined irregularly for 5 years until they were indistinguishable  
25 from low flows prior to harvest (Keppeler and Ziemer 1990). Because an increase in low  
26 flows (i.e., more water in the stream) for summer months generally does not adversely  
27 affect aquatic life, it will not be discussed further. Small volumetric increases may provide  
28 improved habitat conditions (lower stream temperature, increased instream wetted area and  
29 volume) and survivability of aquatic species.

#### 30 **Peak Flows**

31 Peak flow is the maximum instantaneous discharge measured in stream channels during  
32 high flow periods. Management activities can affect peak flows based upon their site-  
33 specific effect, elevational location within a watershed, and proportion of basin forest that  
34 has been altered by timber-related activities, such as roads and timber harvest (Bauer and  
35 Mastin 1997). Peak flows are addressed in greater detail throughout the remainder of this  
36 section.

### 37 **3.5.2.2 Existing Hydrologic Conditions**

38 For general perspective, Table 3-15 provides a breakdown of the number of stream miles in  
39 Washington by analysis region and ownership category. The mileage of streams on all  
40 forestlands in the State, and on the forestlands potentially affected by the FPHCP are  
41 presented by analysis region in Table 3-16.



**Table 3-15.** Stream Miles by Analysis Region and Ownership in Washington State.

	Federal <sup>1/</sup>	Tribal <sup>1/</sup>	State	County	City	Private	Total
North Puget Sound	13,072	230	4,263	164	163	10,761	28,653
South Puget Sound	3,004	121	1,486	81	983	8,157	13,832
West Puget Sound	2,921	100	1,115	3	67	4,908	9,114
Islands	47	-	98	5	4	855	1,009
Olympic Coast	5,154	1,797	3,158	60	-	4,792	14,959
Southwest	1,011	45	3,790	358	127	23,274	28,607
Lower Columbia	7,846	0	3,819	32	-	17,948	29,645
Middle Columbia	8,586	5,567	3,602	10	6	15,109	32,878
Upper Columbia-Downstream of Grand Coulee	19,693	2,618	4,175	11	-	12,372	38,869
Upper Columbia-Upstream of Grand Coulee	9,346	8,302	1,984	57	12	14,215	33,917
Snake River	3,533	-	1,176	3	4	14,773	19,488
Columbia Basin	1,787	-	1,065	-	1	11,304	14,157
<b>Total</b>	<b>76,000</b>	<b>18,779</b>	<b>29,732</b>	<b>783</b>	<b>1,367</b>	<b>138,467</b>	<b>265,129</b>

<sup>1/</sup>Note that stream miles on Federal and tribal ownership are likely underrepresented.

Source: Washington DNR GIS Hydrography layer, 2004, and Major Public Lands layer, 2004.



## Chapter 3

**Table 3-16.** Total and Forested Stream Miles and Stream Miles on Lands Subject to Washington Forest Practices Rules by Analysis Region in Washington State.

Analysis Region	Total Stream Miles	Forested Stream Miles	Streams on Lands Subject to Washington Forest Practices Rules (FPRs)		
			FPR-Regulated Stream Miles	Percent of Total (%)	Percent of Forested (%)
North Puget Sound	28,653	21,534	11,283	39	52
South Puget Sound	13,832	11,039	8,535	62	77
West Puget Sound	9,114	7,669	4,879	54	64
Islands	1,009	549	497	49	91
Olympic Coast	14,959	13,629	7,480	50	55
Southwest	28,607	25,726	24,654	86	96
Lower Columbia	29,645	25,407	18,647	63	73
Middle Columbia	32,878	15,567	6,490	20	42
Upper Columbia-Downstream of Grand Coulee	38,869	19,321	4,389	11	23
Upper Columbia-Upstream of Grand Coulee	33,917	25,302	10,474	31	41
Snake River	19,488	3,156	1,035	5	33
Columbia Basin	14,157	73	70	<1	97
<b>Total</b>	<b>265,129</b>	<b>168,972</b>	<b>98,433</b>	<b>37</b>	<b>58</b>

Note: < = less than

Sources: Washington DNR GIS Hydrography layer, 2004, and Major Public Lands layer, 2004; U.S. Geological Survey National Land Cover Data Layer, 2004.





1 Based on the Washington DNR's statewide stream mapping, the State has approximately  
2 265,000 miles of streams (Table 3-15). This estimate, and all estimates in this paragraph,  
3 should be considered approximate since the State's stream mapping has not been  
4 completed at a consistent level of detail across the State and in all land ownerships. Based  
5 on this mapping, about 47 percent of the stream miles are in western Washington and 53  
6 percent are in eastern Washington. Approximately 169,000 miles (or 64 percent) of the  
7 total stream miles are on forestlands; about 84 percent of the western Washington streams  
8 are on forestland compared to about 46 percent of the eastern Washington streams (Table  
9 3-16). Approximately 98,000 (or 40 percent) of the total stream miles are on forestlands  
10 subject to Washington Forest Practices Rules; about 60 percent of the western Washington  
11 streams and 16 percent of the eastern Washington streams are on forestlands subject to  
12 Washington Forest Practices Rules (Table 3-16).

### 13 **3.5.2.3 Western Washington Peak Flows**

14 Western Washington (and much of eastern Washington) receives moderate to high  
15 precipitation and is influenced by rain-on-snow events. The significance of rain-on-snow  
16 events is the increase in water delivered to the stream system during these events compared  
17 to rainfall alone. When warm air and rain occur on areas with a snowpack, rapid melting  
18 of the snow can occur, resulting in a pulse of water into the drainage network. Rain-on-  
19 snow events can occur on mountain slopes in the transient snow zone, which extends from  
20 altitudes of approximately 1,000 feet to 3,000 feet above sea level (Harr 1986), but can  
21 shift upward or downward during any given storm due to varying meteorological  
22 conditions.

23 Peak flow events associated with rain-on-snow can be of greater magnitude than rain-only  
24 events because the rainfall is augmented by snowmelt. The direct effects of peak flows  
25 include stream channel alteration, bank erosion, redistribution of sediment and large  
26 organic debris, and flooding. In addition to the direct effect of peak flows, rain-on-snow  
27 events generate large inputs of water to the soils and can generate unstable conditions on  
28 hillslopes by increasing the pore-water pressure, which decreases the strength of the soil  
29 (Sidle et al. 1985); a reduction in soil strength increases the potential for slope failure.

### 30 **3.5.2.4 Eastern Washington Peak Flows**

31 In eastern Washington, the buildup of snowpack over winter contributes to large amounts  
32 of spring runoff. Rain-on-snow events are less common. In forested areas east of the  
33 Cascades, snowmelt is the dominant mechanism for producing peak flows, most commonly  
34 in February to July depending upon location and elevation. Snowpack depths are often  
35 greater in forest openings in eastern Washington forests, as conducted in similar studies  
36 (Kattlemann et al. 1983; Troendle 1983). Peak flows are predominantly generated by  
37 snowmelt and may account for most of the 2- to 10-year peak flows. The timing of  
38 snowmelt runoff is important for many eastern Washington watersheds because this runoff  
39 is vital for irrigation supplies and fish habitat.



## Chapter 3

---

### 3.5.2.5 Management Influences on Peak Flows

#### Roads

The design, construction, and maintenance of roads interact with watershed characteristics of soil, topography, and geology and natural disturbances (such as large storms) to determine the effects of roads on the hydrology of a particular watershed. The interception of surface runoff during storms and interception of shallow groundwater flow by a road prism can affect the routing of surface water, extend the channel network (Wemple et al. 1996), increase the potential for higher peak flows, and increase the potential for mass wasting (Montgomery 1994). In a general sense, roads can act as extensions of the drainage network if the roads drain to streams. Road-influenced peak flows have been demonstrated in small watersheds (Ziemer and Lisle 1998); however, the effects of roads on a river basin scale are less understood (Jones and Grant 1996; Beschta and Boyle 1995).

The FFR (1999) set an objective to “maintain surface and groundwater hydrologic regimes (magnitude, frequency, timing, and routing of stream flows) by disconnecting road drainage from the stream network, preventing increases in peak flows from causing scour, and maintaining hydrologic continuity of wetlands.” Two summaries of recent research studies on roads in forested areas demonstrate that roads can have significant effects on peak flows if roads are improperly constructed and if their drainage networks are allowed to become connected to the stream network through improper construction or inadequate maintenance or abandonment procedures (USDA Forest Service 2001; CMER 2004).

CMER (2004) has issued a draft review of published literature that suggests that roads alter runoff processes by the production of overland flow, by interception of subsurface stormwater flow, and by piracy of streams by road ditches. Road runoff may create additional channels in the stream network by incision (Montgomery 1994), potentially leading to adverse consequences for both hydrology and slope stability, depending on the road location.

A road maintenance survey conducted by the Washington DNR (1999) indicated that in 36 subbasins surveyed, the majority were out of compliance with road maintenance objectives for road surface drainage, ditch drainage, and water crossing structures. The authors stated that the rules in effect on January 1, 1999 were “subjective and inadequate,” and recommended that outsloping or water dips would be the first priority in improving performance of road drainage on forest roads. Based on recent research and modeling studies, the U.S. Forest Service (2001) made the following recommendations on a national scale: that forest road locations should be carefully chosen based on a geologic study; that road design should minimize interception, diversion and concentration of water using outsloping and drainage structures; that crossings should be designed to pass all stream materials (including water, fish, debris and sediment); that design and removal of roads should take into account the sensitivity and orientation of the road; and that roads should be designed with failure in mind.

Rashin et al. (1999) recommended a variety of improvements to road BMPs. Those that apply to reduction of peak flows and direct input of road runoff to streams include: minimization of road location within 485 feet (150 meters) of streams, revised spacing of



1 relief culverts, adequate sediment traps, and spreading or dissipation measures to prevent  
2 incision from runoff. They also recommended against the use of ambiguous language in  
3 BMPs.

### 4 **Timber Harvest**

5 The best understood effect of timber harvest is its influence on stream flow relating to  
6 altering snow accumulation and melt rate. Increased peak flows can occur in the winter,  
7 when a warm wet storm brings rain after a cold storm deposits substantial amounts of  
8 snow. Many floods in Washington, mostly on the westside of the Cascades, have occurred  
9 as a result of rain-on-snow events. While rain-on-snow events are a natural occurrence,  
10 their effects can be exacerbated when a watershed has been logged in a short amount of  
11 time (25 to 30 years) (Coffin and Harr 1992; Troendle and Leaf 1981).

12 The two most important watershed variables that affect rain-on-snow events are elevation  
13 and extent of timber harvest. Timber harvest has the potential to alter snow accumulation  
14 and melt rates in any portion of a watershed, but predominantly in the “rain-on-snow”  
15 zone. The rain-on-snow zone in western Washington typically occurs between 1,200 and  
16 4,000 feet in elevation (Washington DNR 1997a). Forest openings are conducive to  
17 increased snow pack accumulations because more snow reaches the ground as a result of  
18 less snow interception by the tree canopy. Once rainfall associated with a storm occurs,  
19 the forest openings are more conducive to higher rates of convection and condensation to  
20 the snow pack than the surrounding forest. The combination of greater snow accumulation  
21 and increased melt rates can lead to a greater rate of moisture available at the soil surface  
22 in forest openings during a rain-on-snow event than occurs in the adjacent forest (Coffin  
23 and Harr 1992). The net result is that an increase in runoff is expected from forest  
24 clearcuts in areas where rain-on-snow is prevalent.

25 Although not as well understood, timber harvest may increase snowmelt peak flows  
26 (Benda et al. 1998). Because timber harvest can cause increased snow accumulation in  
27 openings, areas where runoff is dominated by snowmelt can theoretically experience  
28 increased peak flows. However, research in the Pacific Northwest has not consistently  
29 demonstrated this effect. While Cheng (1989) found as much as a 35 percent increase in  
30 peak flows with 30 percent clearcuts in British Columbia, Fowler et al. (1987) found no  
31 effect in small watersheds in Oregon. In perhaps the most comprehensive study, Anderson  
32 and Hobba (1959) found an 11 percent increase in spring peak flows across 21 watersheds  
33 in eastern Oregon. This area is analogous to eastern Washington.

34 Rain-dominated watersheds, such as along the coast, may also be subject to increased peak  
35 flows, but for different reasons. Studies that have shown peak flow increases in rain-  
36 dominated watersheds (Harr et al. 1975; Harr 1986) have correlated the increases with soil  
37 compaction, rather than timber harvest. Yet other studies indicate no change in peak flow  
38 after harvest (Benda et al. 1998). If a peak flow following harvest occurs, small basins  
39 seem to be more likely to experience effects than large basins. From a mass balance  
40 perspective the contribution to the total discharge should be more significant in smaller  
41 basins.



## Chapter 3

---

1 Most recent research suggests that peak flow changes due to forest practices are difficult to  
2 detect on large river systems; effects of peak flow changes due to forest practices in small  
3 basins are highly variable, but small peaks are apparently affected more than large peaks  
4 (e.g., Thomas and Megahan 1998; Beschta et al. 2000).

### 5 **Roads in Riparian Areas**

6 Historic road routing and construction practices have, at times, led to substantial impacts  
7 on riparian systems. Roads were often built along flat floodplains, which resulted in the  
8 removal of riparian vegetation. In narrow canyons with limited floodplains, roads were  
9 commonly located on the sideslope within the riparian zone or encroached on the stream  
10 channel; some roads even used the actual stream channels. Even in the absence of these  
11 longitudinal impacts, the continuity of the riparian corridor has been interrupted at each  
12 bridge and culvert crossing (Kondolph et al. 1996). Consequently, roads built in riparian  
13 areas have changed riparian forest structure and composition and caused permanent land  
14 disturbance.

15 Stream-adjacent roads can cause the loss of some or all riparian functions within riparian  
16 lands depending on where road construction has occurred. One example is the loss of  
17 LWD recruitment potential from trees on the upland side of roads within riparian areas.  
18 Most of the trees on the upland side are not be recruited as LWD because they are typically  
19 removed when the tree falls onto the road. Major changes to the aquatic system have also  
20 occurred from riparian land modifications due to road development, including the  
21 straightening or simplification of the stream channel system (Knutson and Naef 1997;  
22 Oregon Department of Forestry 1999a).

23 Currently, limited information on statewide road density or distribution of roads in riparian  
24 areas is available. However, for a general perspective, the U.S. Forest Service has  
25 quantified the number of roads built on National Forest System lands in Washington. By  
26 1907, only 147 miles of road had been built in all of Washington's National Forest System  
27 lands. By 1962, the length of roads on National Forest System lands in Oregon and  
28 Washington had risen to 22,000-24,000 miles, and to over 90,000 miles in 1990. In the  
29 late 1990s, it was estimated that about 3,000 miles of new roads were being constructed  
30 annually in the western-forested area of the United States (Knutson and Naef 1997).

31 In eastern Washington, increased roading has allowed greater access for forest  
32 management and some types of recreation, and has contributed to the protection of the  
33 forest from the spread of fires and catastrophic outbreaks of insects. Railroads were also  
34 built into some areas, and eventually many railroad grades were converted to roads. The  
35 decision of where and when to build roads has always hinged on the logistics of timber  
36 harvesting (Oliver et al. 1994). As the density of roads increases, road impacts on riparian  
37 areas will inevitably increase (Knutson and Naef 1997).

### 38 **3.5.3 Groundwater Quantity and Quality**

39 As a fundamental component of the hydrologic cycle, groundwater plays a critical role in  
40 maintaining the health of riparian and wetland ecosystems. Groundwater sustains stream  
41 and river baseflows and influences the thermal regime of many surface waters in



1 Washington. From a human perspective, groundwater supplies more than one-quarter of  
2 the State's water demand and is estimated to provide at least 65 percent of the drinking  
3 water for the State's residents (Washington Department of Ecology 2002b). In large areas  
4 east of the Cascade Range, 80 to 100 percent of available drinking water is obtained from  
5 groundwater resources. In addition, some areas of the State, including most of Island and  
6 San Juan Counties, rely solely on groundwater sources for potable water. Of the total  
7 number of public water supply systems in Washington, over 95 percent use groundwater as  
8 their primary water source (U.S. Environmental Protection Agency 1999).


9 Washington has three principal groundwater aquifers (Washington Department of Ecology  
10 2002b). They include: 1) the basalts and overlying unconsolidated deposits of the Central  
11 Columbia Plateau in southeastern Washington, 2) the unconsolidated glacial deposits of the  
12 Puget Sound Lowland, and 3) the glacial outwash deposits of the Spokane-Rathdrum  
13 Prairie Aquifer in northeast Washington. These larger aquifer systems are typically  
14 composed of multiple water-bearing units that underlie the surface, often extending many  
15 hundreds of feet below ground. A number of smaller aquifer systems also exist throughout  
16 the State, commonly located within river valleys.

17 When used for consumptive purposes, groundwater is usually accessed by shallow wells  
18 drilled into unconsolidated-deposit aquifers that consist primarily of sand and gravel but  
19 contain variable quantities of clay and silt (U.S. Geological Survey 1994). Many high-  
20 yield public supply and irrigation wells and thousands of domestic wells are completed in  
21 these types of aquifers and most are located on privately owned lands. In many places,  
22 deeper wells produce water from underlying volcanic rocks such as basalt.

23 Groundwater is often connected directly or indirectly to rivers, streams, lakes, and other  
24 surface water bodies, with exchange and mixing occurring between the sources.  
25 Contaminants entering groundwater can therefore affect surface waters (and vice versa)  
26 and associated aquatic organisms. Depending on the geologic and hydrologic conditions of  
27 the aquifer, contaminated groundwater may reach surface waters within days or may take  
28 hundreds or even thousands of years (U.S. Environmental Protection Agency 1986).

29 Currently, there is no comprehensive, statewide groundwater monitoring program in  
30 Washington. Groundwater monitoring efforts cover only a small portion of the primary  
31 aquifers in the State and vary in parameters measured, methods of measurement, data  
32 quality, and degree of long-term commitment (Washington Department of Ecology 2002b).  
33 As a result, it is difficult to assess the general status of groundwater in Washington.  
34 Nevertheless, issues concerning the potential effects of forest practices on groundwater,  
35 including changes in quantity and quality, are addressed in Chapter 4 (Environmental  
36 Effects).

37



## **Chapter 3** \_\_\_\_\_

1 This page is intentionally left blank.

2



## 1 3.6 VEGETATION

### 2 3.6.1 Forest Vegetation

3 About half (22.9 million acres) of the 43.3 million acres that make up Washington State are  
4 mapped as forestland (Table 3-2). In terms of broad categories of land cover, the State  
5 consists of 53 percent forestland, 16 percent shrubland, 7 percent grassland, 18 percent  
6 agricultural lands, 2 percent freshwater and wetlands, 2 percent perennial ice/snow and  
7 rock, and 2 percent developed residential and commercial lands (Table 3-2). Of the  
8 forestlands, approximately 50 percent are covered by the Washington Forest Practices  
9 Rules; these primarily include privately owned and state-managed forestlands, although  
10 forestlands managed by local governments are also covered by the rules. The spatial  
11 distribution of covered lands is shown along with the analysis regions in Figure 3-1. Table  
12 3-3 lists the estimated acreage of covered lands and other forestlands in the State by  
13 analysis region and ownership category.

14 Due to their wide distribution throughout the State, covered lands vary markedly in terms  
15 of their physical characteristics. Franklin and Dyrness (1973) mapped six forested  
16 vegetation zones in Washington (Figure 3-3). Covered lands are present in each of the six  
17 zones. The six zones include: 1) the Sitka spruce (*Picea sitchensis*) zone, 2) the western  
18 hemlock (*Tsuga heterophylla*) zone, 3) the ponderosa pine (*Pinus ponderosa*) zone, 4) the  
19 grand fir (*Abies grandis*) and Douglas-fir (*Pseudotsuga menziesii*) zones, 5) the subalpine  
20 forest zones [including the Pacific silver fir (*Abies amabilis*), mountain hemlock (*Tsuga*  
21 *mertensiana*), and subalpine fir (*A. lasiocarpa*) zones], and 6) the Willamette Valley zone.  
22 An overview of the vegetative and climatological characteristics of each zone is provided  
23 below, based on Franklin and Dyrness (1973).

24 **Sitka Spruce Zone** – The Sitka spruce zone stretches the length of the Washington coast  
25 and is generally only a few miles wide except where it extends up river valleys  
26 (Figure 3-3). The zone is much broader along the westside of the Olympic Peninsula  
27 where an extensive coastal plain exists. The zone typically lies below 500 feet in  
28 elevation, although it may be found up to 2,000 feet in elevation where mountains are  
29 adjacent to the ocean. Approximately 1.3 million acres of covered lands (12 percent of all  
30 covered lands) lie within the Sitka spruce zone.

31 The Sitka spruce zone has the mildest climate of any forest region in Washington.  
32 Extremes in moisture and temperature are minimal due to the proximity to the ocean.  
33 Annual precipitation is high, averaging between 80 and 120 inches; most of this falls as  
34 rain during the fall and winter months. Frequent fog and low clouds during the relatively  
35 dry summer months are probably important in reducing moisture stresses on trees and other  
36 vegetation.

37 Dominant tree species include Sitka spruce (*Picea sitchensis*), western hemlock (*Tsuga*  
38 *heterophylla*), western redcedar (*Thuja plicata*), Douglas-fir (*Pseudotsuga menziesii*),  
39 grand fir (*Abies grandis*), and Pacific silver fir (*Abies amabilis*). Red alder (*Alnus rubra*) is  
40 common on disturbed sites and shore pine (*Pinus contorta*) is common along the ocean.



## Chapter 3

---

1 **Western Hemlock Zone** – The western hemlock zone is the most extensive forest region  
2 in western Washington. The region is famous for its sub-climax forests of Douglas-fir and  
3 climax forests of western hemlock and western redcedar. The zone extends from British  
4 Columbia south through the Olympic Peninsula, Coast Ranges, Puget Trough, and Cascade  
5 physiographic provinces (Figure 3-3). In the Cascades, the western hemlock zone is found  
6 from sea level to 2,200 feet in the north and from 400 to 3,000 feet in the south. The zone  
7 lies between 500 and 1,800 feet on the western slopes of the Olympic Mountains but  
8 ranges from sea level to 3,700 feet on the drier eastern slopes. The western hemlock zone  
9 encompasses the largest proportion of covered lands at 6.3 million acres (55 percent of all  
10 covered lands).

11 The western hemlock zone has a wet, mild marine climate. Because the zone lies further  
12 from the ocean, temperature and moisture extremes are greater than in the Sitka spruce  
13 zone. Within the zone, climatic variation is high due to differences in latitude, elevation,  
14 and location with respect to mountain ranges. Annual precipitation ranges from 60 to  
15 120 inches, most of which falls as rain during the fall and winter months. Between 6 and 9  
16 percent of annual precipitation occurs during the summer.

17 Douglas-fir, western hemlock, and western redcedar are the dominant tree species. Pacific  
18 silver fir is common near the upper elevation limits and even well within the western  
19 hemlock zone in the North Cascades and Olympic Mountains. Grand fir, Sitka spruce, and  
20 western white pine (*Pinus monticola*) occur sporadically. Red alder, bigleaf maple (*Acer*  
21 *macrophyllum*), and black cottonwood (*Populus balsamifera* spp. *trichocarpa*) are  
22 common in riparian areas. Red alder is also common on disturbed sites.


23 **Willamette Valley Zone** – The Willamette Valley zone is part of the Interior Valley zone  
24 or Pine-Oak-Douglas-fir zone of Franklin and Dyrness (1973), and barely extends into  
25 southwest Washington around Vancouver (Figure 3-3). It consists mostly of the  
26 Willamette River valley bottom and adjacent lowlands, enclosed by the Cascade Range on  
27 the east and the Coast Ranges on the west. It is generally warmer and drier than the  
28 western hemlock zone and consists of a vegetational mosaic resulting from a long history  
29 of human influences. The mosaic includes Oregon white oak (*Quercus garryana*)  
30 woodlands, coniferous forests, grasslands, shrub communities, and riparian forests.  
31 Approximately 28,000 acres of covered lands (less than 1 percent of all covered lands) lie  
32 within the Willamette Valley zone.

33 **Subalpine Forests** – Subalpine forests in Washington include the Pacific silver fir,  
34 mountain hemlock, and subalpine fir zones (Figure 3-3). The Pacific silver fir zone is the  
35 lowest of the three zones and occupies the western slopes of the Cascade Range and all but  
36 the drier northeastern slopes of the Olympic Mountains at elevations ranging from 2,000 to  
37 4,300 feet. The mountain hemlock zone is the highest forest zone along the western slopes  
38 and crest of the Cascades and Olympics and generally lies between 4,100 and 6,000 feet  
39 elevation. This zone extends varying distances east across the Cascade crest until it is  
40 gradually replaced by the subalpine fir zone. The subalpine fir zone is common on  
41 secondary ranges that extend east from the Cascade crest, in the Okanogan Highlands of  
42 north-central Washington, and in the Blue Mountains of southeastern Washington. Its





Figure 3-3. Forest Vegetation Zones (Franklin and Dyness 1973) and Covered Lands in Washington







1 lower elevation boundary is generally 4,900 feet in the Cascades and 4,200 to 5,600 feet in  
2 other areas. Because covered lands generally lie at lower elevations, the subalpine forest  
3 region encompasses a relatively small proportion of covered lands (626,000 acres or  
4 6 percent of all covered lands).

5 Subalpine forests in Washington have wet, cool climatic regimes. Annual precipitation  
6 averages between 55 and 110 inches and is strongly influenced by elevation. Much of the  
7 precipitation falls as snow during the fall and winter months. Summers are cool and  
8 winters are cold with snow cover persisting for up to 6 months, particularly in the  
9 mountain hemlock zone.

10 Typical tree species in the Pacific silver fir zone include Pacific silver fir, western  
11 hemlock, noble fir (*Abies procera*), Douglas-fir, western red-cedar, and western white pine.  
12 The mountain hemlock zone is dominated by mountain hemlock, subalpine fir, and  
13 lodgepole pine (*Pinus contorta*), while subalpine fir, Engelmann spruce (*Picea*  
14 *engelmannii*), and lodgepole pine comprise the major tree species in the subalpine fir zone.

15 **Douglas-Fir/Grand Fir Zones** – The Douglas-fir and grand fir zones are found in eastern  
16 Washington and generally lie above the drier ponderosa pine zone and below the subalpine  
17 forests. These zones extend north from the Oregon-Washington border along the eastern  
18 slopes of the Cascades and across north-central and northeastern Washington (Figure 3-3).  
19 The Douglas-fir zone is typically found between 2,000 and 4,300 feet in northeastern  
20 Washington. This zone is more mesic than the lower elevation ponderosa pine zone, with  
21 cooler temperatures and higher annual precipitation. Douglas-fir, ponderosa pine,  
22 lodgepole pine, and western larch (*Larix occidentalis*) comprise the major tree species in  
23 the zone. Together, the Douglas-fir and grand fir zones encompass 2.1 million acres of  
24 covered lands (19 percent of all covered lands).

25 The grand fir zone usually lies above the Douglas-fir zone and has the most moderate  
26 environmental regime of any eastern Washington forest zone. Neither moisture nor  
27 temperature conditions are extreme. Precipitation is generally higher and temperatures  
28 lower than in lower elevation forest zones. Major tree species in the grand fir zone include  
29 grand fir, ponderosa pine, lodgepole pine, western larch, and Douglas-fir.

30 **Ponderosa Pine Zone** – The ponderosa pine zone occupies three areas in Washington: 1)  
31 a narrow band (10 to 20 miles wide) along the eastern flanks of the Cascade Range, 2) the  
32 Blue Mountains, and 3) the Okanogan Highlands (Figure 3-3). The zone lies between  
33 2,000 and 4,000 feet in elevation along the eastern flanks of the Cascades and between  
34 3,000 and 5,000 feet in the Blue Mountains. Covered lands occur sporadically throughout  
35 the ponderosa pine zone, encompassing 795,000 acres or 7 percent of all covered lands.

36 The ponderosa pine zone is characterized by a short growing season and minimal summer  
37 precipitation. Average annual precipitation ranges from 14 to 30 inches, much of which  
38 falls as snow during the winter months. Diurnal summer temperatures fluctuate widely,  
39 with hot days and cold nights. Winter temperatures are generally low and snow often  
40 accumulates to considerable depths.



## Chapter 3

---

1 Ponderosa pine is the climax species and is commonly associated with quaking aspen  
2 (*Populus tremuloides*) and lodgepole pine throughout the zone. In the Okanogan  
3 Highlands, grand fir, Douglas-fir, western larch, and western white pine are common  
4 associates, while in south-central Washington, Oregon white oak is often present.

### 5 **3.6.2 Riparian Vegetation**

6 The vegetative communities that are commonly associated with riparian areas can be  
7 divided into three general areas of Washington: forested areas in western Washington,  
8 forested areas in eastern Washington, and the non-forested shrub-steppe region in eastern  
9 Washington (Knutson and Naef 1997). For the purpose of this DEIS only forested riparian  
10 areas are generally described. The species, sizes, and density of vegetation occupying a  
11 riparian site are dependent upon soil moisture conditions and disturbance history.

12 Western Washington riparian habitats are associated with wet environmental conditions.  
13 Although considerable site-specific variability occurs, general vegetative characteristics  
14 include the presence of a mixture of conifer and hardwood trees (hardwoods are more  
15 abundant where natural and human disturbance is frequent). Common conifers found in  
16 riparian areas are western hemlock, western red-cedar, and Sitka spruce – species that are  
17 tolerant of shade and periodically saturated soils. Red alder is nearly always found in  
18 young and disturbed stands. Upland conifers (e.g., Douglas-fir) and hardwoods (e.g.,  
19 bigleaf maple) are dominant along small streams, which have narrow riparian areas.  
20 Lowland rivers and forested swamps with frequent flooding or gravelly soils often include  
21 black cottonwood, willow (*Salix* spp.), and red alder. Swampy areas may also have vine  
22 maple (*Acer circinatum*), cascara (*Rhamnus purshiana*), willow, western red-cedar, Sitka  
23 spruce, and western hemlock (Knutson and Naef 1997). Some of these species also are  
24 common components of riparian areas on drier sites.

25 Forested riparian areas of eastern Washington are typically located in deeply incised  
26 ravines in mountainous terrain (Carlson et al. 1990 in Knutson and Naef 1997). Lower  
27 elevations with moist soils and temperate microclimates support cedar, western hemlock,  
28 bigleaf maple, quaking aspen, and other hardwood trees. Large trees, snags, and downed  
29 wood can be abundant in areas depending on disturbance history. These relatively moist  
30 riparian areas also include a variety of understory shrubs and herbs including willow, red-  
31 osier dogwood (*Cornus stolonifera*), mountain alder (*Alnus incana tenuifolia*), devil's club  
32 (*Oplopanax horridum*), and other species. Drier sites are characterized by ponderosa pine  
33 in the uplands while trees in riparian areas include Douglas-fir, paper birch (*Betula*  
34 *papyrifera*), black cottonwood, and quaking aspen. High elevation riparian sites often have  
35 saturated soils that are dominated by understory species rather than by tree species. Where  
36 trees exist, they are commonly subalpine fir or Engelmann spruce and down wood is  
37 abundant because cold temperatures slow decomposition. Shrubs and herbs at high  
38 elevations are relatively diverse, but generally stunted due to the more severe  
39 environmental conditions (Knutson and Naef 1997).



### 1 **3.6.3 Disturbance Agents**

2 Stand disturbance has long been a factor in Pacific Northwest forests. Disturbance factors  
3 include fire, wind, insects, and pathogens.

#### 4 **3.6.3.1 Fire**

5 Fire has been an integral part of the forest environment for thousands of years. The  
6 likelihood of a fire initiation and the extent and severity of the resulting fire are affected by  
7 the vegetation and other fuels on the site. The most common natural cause of forest fires is  
8 lightning. Areas east of the Cascade crest average 10 to 15 thunderstorms per year, while  
9 areas west of the Cascades average five per year. Most of the forested areas of Washington  
10 experience between one and six lightning fires per 100,000 acres each year. Lightning  
11 fires, which usually start as the result of lightning strikes in large trees or snags, account  
12 for approximately 37 percent of the forest fires in Washington. Less than 1 percent is  
13 caused by spontaneous combustion or other natural causes (Agee 1993). The remaining  
14 fires are caused by humans, and are due to campfire escapes, industrial activity, other  
15 accidents, or are intentionally set. When conditions are dry and fuel is abundant, fires can  
16 burn large areas. One lightning fire in Chelan County, the 1994 Tye Fire, burned more  
17 than 140,000 acres.

18 In the cool, moist climate of western Washington, climatic conditions, fuel accumulation,  
19 and lightning ignition historically have combined to result in infrequent extensive stand-  
20 replacement fires (fires that kill virtually all vegetation). These fires were generally more  
21 intense than fires in eastern Washington and often resulted in a 50- to 100-year time span  
22 before the burn area became fully restocked with native conifers (Franklin et al. 1981).  
23 The historical natural fire return interval for cedar, spruce, and hemlock stands of western  
24 Washington is about 937 years and about 217 years for Douglas-fir stands (Agee 1993).  
25 Higher moisture levels found in riparian areas can increase fire return intervals in those  
26 areas (Agee 1993). The return interval (cycle or turnover time) is the mean time between  
27 disturbances on a given site.

28 Historically, fires in eastern Washington have been more frequent and less intense than  
29 fires in western Washington, with ponderosa pine forests experiencing extensive fires  
30 every 15 years on average, mixed conifer forests an average of every 50 years, and the  
31 moister, high elevation forests experiencing fire only about once every 500 years (Agee  
32 1993). Often the more frequent fires in eastern Washington represented understory burns  
33 that maintained the canopy, or at least a portion of the canopy and were not stand  
34 replacement fires where all vegetation would have been killed. These frequent lighter fires  
35 in the ponderosa pine and Douglas-fir zones were instrumental in maintaining the open  
36 pine forests noted by 19<sup>th</sup> century settlers. Natural disturbance from fire on the eastside is  
37 an important factor defining stand seral stage characteristics under unmanaged condition  
38 (Washington Forest Practices Board 2001a).

39 Fire control practices during the last century, in all areas of Washington, have increased the  
40 fire intervals throughout the State (i.e., fires occur less frequently) (Agee 1993). These  
41 increased fire intervals cause buildup of unnatural levels of dead and dying materials that  
42 can fuel fires. This fuel buildup results in more intense and frequent stand-replacement



## Chapter 3

---

1 fires, particularly in eastern Washington. Currently, fuel levels are high in many parts of  
2 eastern Washington because the frequent understory fires that once burned these areas and  
3 kept fuel levels in check have been aggressively suppressed. These frequent understory  
4 fires that historically occurred in eastern Washington helped maintain natural ponderosa  
5 pine forests that were open and park-like because ponderosa pine is a species that relies on  
6 high heat for natural regeneration of the species. The heat of forest fires facilitates the  
7 opening of ponderosa pine cones allowing the seeds to disperse and germinate. The  
8 reduction in forest fires in eastern Washington could be linked to a reduction in the  
9 quantity of ponderosa pine forests. Management of a disturbance agent such as fire, can  
10 itself cause a disturbance effect by altering types of vegetation present (Washington Forest  
11 Practices Board 2001a).

### 12 **3.6.3.2 Wind**

13 Historically, windstorms have had a larger effect on westside forests than on eastside  
14 forests, and fire has had a larger effect on eastside forests than on westside forests. Wind  
15 has been especially potent in coastal Sitka spruce and high elevation Pacific silver fir and  
16 alpine forests, where the moist conditions generally limited fire spread (Agee 1993). The  
17 historic return intervals for windstorms in western Washington are approximately 119 to  
18 384 years for small-scale to large-scale storms (Washington Forest Practices Board 2001a;  
19 Harcombe 1986 as cited in Agee 1993). The 1921 windstorm and the Columbus Day 1962  
20 windstorm blew down thousands of acres of mature trees in the western Olympic Peninsula  
21 and other areas of western Washington (Washington DNR 2003). In addition to major  
22 regional wind storms, locally intense winds are an important cause of small-scale  
23 disturbance, which occurs across the landscape on an annual basis.

### 24 **3.6.3.3 Insects and Pathogens**

25 Insects and pathogens (fungi, viruses, dwarf mistletoe) are an important part of forest  
26 ecosystems. They have many, complex biological roles and interactions. Insects and  
27 pathogens may weaken or kill trees; stimulate altered tree growth forms such as forked  
28 tops, galls, and bushy growths; affect light and heat reaching streams or the forest floor;  
29 and cause a fertilization effect with feces. Insects act as pollinators, but also consume  
30 seeds and fruit. Insects, fungi, and dwarf mistletoe plants are prey or food resources for  
31 fish (Dodge 2001) and wildlife. The presence or absence of some insect species can give  
32 indications of water quality characteristics. Mycorrhizal fungi enhance the uptake of  
33 water and nutrients by roots. Saprophytic insects and fungi consume dead plant and animal  
34 material, affecting the softness and longevity of snags, increasing soil quality, releasing  
35 nutrients.

36 The effects of native insects and pathogens are considered to be part of how ecosystems  
37 function. Damaged trees become nest sites for birds and salamanders and make space for  
38 other trees and plants to grow. A root rot pocket may not grow Douglas-fir for the next  
39 60 years, but the openings, scattered mortality, and increased volume of LWD may provide  
40 big game forage, small animal habitat sources, increased diversity in plant communities,  
41 and enhanced visual quality at the landscape scale (Theis and Sturrock 1995). Humans  
42 may judge the activity of insects and pathogens in relation to their long- and short-term



1 management objectives for the land. Humans intervene or affect these interactions by  
2 conducting management activities that control the trees species, structure, and vigor on a  
3 site. Non-native insects and diseases, introduced by human activities, have the potential to  
4 disrupt ecosystem balance and processes. Some examples of insect/pathogen and forest  
5 interactions are described below.

### 6 **Exotic Pest Species and Native Hosts Natural Defenses**

7 The white pine blister rust fungus was introduced accidentally in the early 20<sup>th</sup> century. It  
8 has killed vast numbers of native western white pine that produce heavy seed crops for  
9 wildlife, once abundant and widespread in eastern and western Washington. This fungus  
10 has now reached alpine areas where high losses to whitebark pine (*Pinus albicaulis*), a  
11 critical food and habitat tree in this harsh environment, are expected (Baskin 1998; Goheen  
12 et al. 2002). The balsam woolly adelgid, an introduced insect, is making grand fir more  
13 susceptible to drought and insect defoliators, and will likely reduce the range of subalpine  
14 fir (which die at low elevations, but survive in higher sites where it is too cold for the  
15 insect to complete its lifecycle) (Mitchell 2001). The exotic defoliator, larch casebearer,  
16 causes some damage to western larch, but has come somewhat into balance because  
17 intentionally-introduced exotic predatory insects control it (Ryan 1997). The exotic  
18 cinnabar moth was intentionally released in Washington and has successfully suppressed  
19 populations of the tansy ragwort, a noxious weed that invades grassland habitats and is  
20 toxic to animals that eat it (Washington State Noxious Weed Control Board 2004).

### 21 **Fire Suppression and Insects/Pathogens**

22 Humans have succeeded at suppressing almost all but the most severe and extreme forest  
23 fires. Therefore, particularly in eastern Washington where light fires were more common,  
24 without the thinning, clearing and diversity enhancing effects of fire, forests are more  
25 dense, dominated by shade-tolerant species (Douglas-fir and grand fir replacing pine and  
26 larch), and more uniform in structure than in the past. Weak, crowded trees are more  
27 susceptible to being killed by bark beetles, particularly during times of drought  
28 (Edmonds et al. 2000). Crowded trees in dense stands are excellent habitat for defoliating  
29 caterpillars such as the western spruce budworm and Douglas-fir tussock moth (that feed  
30 on Douglas-fir and true firs). Insect defoliator outbreaks are more frequent, last longer,  
31 and cause more damage to affected trees than prior to European settlement (Hessburg et al.  
32 1994). The dense forests are also more susceptible to root diseases and dwarf mistletoe  
33 infection because close proximity of weak trees facilitates transmission of the pathogens.  
34 Alternatively, the insect defoliators of mature pine trees have been less numerous, and  
35 insect and wildlife species (such as the mardon skipper butterfly, an endangered butterfly  
36 in Washington) that use the grasses and openings beneath pine trees have been reduced  
37 (Hessburg et al. 1994; Potter et al. 1999).

### 38 **Forest Fragmentation**

39 Fire suppression and logging have affected the continuity of forests. While fire  
40 suppression has made forests more uniform; logging has tended to break forest stands into  
41 discrete units (although the uniformity within a stand may be increased as it is planted to  
42 only one species with a relatively uniform spacing). The native defoliators, western spruce



## Chapter 3

---

1 budworm and Douglas-fir tussock moth, have benefited by the increase in forest density  
2 and continuity that fire suppression has brought (Hessburg et al. 1994).

### 3 **Forest Management and Pathogens**

4 Native tent caterpillars benefit from edge habitat and hardwood-favoring disturbances that  
5 logging and settlement have brought to Puget Sound. The white pine weevil thrives when  
6 Sitka spruce are grown in large openings or plantations, in contrast to the partial shade they  
7 probably grew in following fire or windthrow disturbances (Oregon Department of  
8 Forestry 1999b; Hamid et al. 1995). Moreover, creation of stumps enhances many root  
9 diseases. Root disease fungi persist in decaying underground root material and stumps  
10 long after the infected tree has died or been removed. If the tree is cut, the root rot fungi  
11 can persist in the buried portions of the stump for decades. If the tree tips over, then most  
12 of the roots and stump are levered out of the ground and dry out, reducing fungus  
13 inoculums on the site. Annosum root and butt rot invades trees when spores land on  
14 wounds or cut stumps. Levels of Annosum root and butt rot were reduced when forest  
15 managers initially removed very old, infected trees, but will increase with increasing  
16 management activities that create stumps and basal wounds and maintain trees for longer  
17 rotation ages (Schmitt et al. 2000).

### 18 **Species Composition**

19 Insects and pathogens have generally evolved to feed on a single or few closely-related  
20 host tree species. Their activity levels are affected by the abundance of their host tree. For  
21 example, the native foliar pathogen Swiss needle cast causes premature needle loss in  
22 Douglas-fir (Southwest Oregon Forest Insect and Disease Service Center 2004). It is most  
23 severe near coastal areas because the spores are protected by moist, foggy weather. Where  
24 humans have replaced coastal hemlock forests with Douglas-fir plantations, the intensity of  
25 Swiss needle cast disease has risen (Southwest Oregon Forest Insect and Disease Service  
26 Center 2004). In contrast, the hemlock looper (caterpillar) feeds on the foliage of western  
27 hemlock and associated conifers. It reaches high outbreak populations in old growth  
28 hemlock forests with abundant moss and lichen for egg laying sites.

### 29 **Snag and Downed Log Abundance**

30 Snags and dead portions of trees are important nesting sites for woodpeckers and carpenter  
31 ants, predators of bark beetles, and defoliating caterpillars. In many managed forests,  
32 timber harvest, salvage logging (i.e., commercial harvest of dead, downed logs) and  
33 firewood gathering have reduced the availability of snags and large logs, thus reducing  
34 insect predators. The natural population cycles of the herbivorous insects may become  
35 more intense, erratic, and damaging (Bull et al. 2001). Ants are an important food for  
36 black bears when berries are scarce (Schowalter et al. 2001). Logs may also protect soil  
37 organisms from disturbances like fire, summer drought, and timber harvest. Without  
38 sufficient numbers of logs, recolonization of disturbed soils could be slowed. These soil  
39 organisms provide critical functions of nutrient cycling and soil aeration (Schowalter et al.  
40 2001).





### 1 **3.6.3.4 Special Problems in Eastern Washington Riparian Areas**

2 Disturbance following European settlement has occurred primarily through timber harvest,  
3 land clearing, agriculture, and, in some areas, fire. Much of Washington forestlands have  
4 been logged at least once in the past 100 years (Washington DNR 2003). Generally,  
5 timber was selectively logged (i.e., the removal of specific trees only) in eastern  
6 Washington. As a result, eastern Washington riparian areas include dense understories,  
7 dense reproduction, and more fire-intolerant species resulting in high fuel accumulation  
8 and more intense and destructive fires as compared to natural conditions (Wissmar et al.  
9 1994; Washington Forest Practices Board 2001a).

10 Riparian habitat disturbance factors include all of the above mentioned factors plus a few  
11 additional factors in eastern Washington that can have significant impacts. These  
12 disturbance factors include grazing, mining, and irrigation (DEIS Appendix A). Studies  
13 have shown that livestock grazing within riparian areas eliminates or reduces streamside  
14 vegetation, destabilizes streambanks, causes channel sedimentation and aggradation,  
15 widens channels, increases stream temperature extremes, lowers the water table, reduces  
16 bank undercut, and reduces pool frequency and depth (Armour et al. 1991; Chaney et al.  
17 1993; Kauffman and Krueger 1984; Kovalchik and Elmore 1992; Meehan 1991; Platts  
18 1991). Grazing pressure usually is higher in the riparian zone because there typically is  
19 more shade, surface water for drinking, and more succulent vegetation than outside of  
20 these zones (Platts 1991).

### 21 **3.6.4 Threatened and Endangered Plants**

22 The Washington DNR Natural Heritage Program maintains a list of threatened,  
23 endangered, and sensitive plant species known to occur in each county. The list is derived  
24 from a comprehensive Geographic Information System database of known occurrences of  
25 threatened, endangered, and sensitive plants in the State ([http://www.dnr.wa.gov/nhp/  
26 refdesk/plants.html](http://www.dnr.wa.gov/nhp/refdesk/plants.html)). Many threatened, endangered, and sensitive plant habitats, such as  
27 alpine, beach, exposed rock, or exposed grassy bluff, are not likely to be affected by  
28 harvest or harvest-related activities. Other habitats such as meadows, prairies, or forest  
29 openings may not support trees for harvest but may be adjacent to harvest areas and could  
30 potentially be affected by harvest activities. The species that occur in forested habitats,  
31 including microhabitats in forests such as forest openings, have a high likelihood of being  
32 affected by harvest or harvest-related activities. Table 3-17 shows federally listed and  
33 candidate plant species, their habitat, and distribution on covered lands by analysis region.  
34 None of these species are intended to be covered by a take authorization under the FPHCP.

### 35 **3.6.5 Invasive Plants**

36 The historical emphasis of Washington State noxious weed law has been to protect the  
37 economic interests of commercial agriculture in the State. While the effects of noxious  
38 weed species on agriculture are large, their effects on the natural resources and ecological  
39 diversity of the State are also large. These resources, once destroyed, may be irreplaceable.  
40 Noxious weeds threaten not only agriculture, but also rangelands, waterways, parks,



## Chapter 3

1 **Table 3-17.** Federally Listed and Candidate Plant Species in Washington, their Habitat,  
2 and their Current and Potential Distribution

Species	Federal Status <sup>1</sup>	Habitat	Current Distribution <sup>2</sup>	Potential Distribution <sup>3</sup>
Marsh sandwort ( <i>Arenaria paludicola</i> )	E	Marshes, bogs	N/A <sup>4</sup>	NP, WP, SP, OC, SW
Showy stickseed ( <i>Hackelia venusta</i> )	E	Sand, rocks	UCDS	UCDS
Bradshaw's desert-parsley ( <i>Lomatium bradshawii</i> )	E	Wetlands, prairies, streams	LC	LC, SW
Wenatchee Mountains checker-mallow ( <i>Sidalcea oregana</i> var. <i>calva</i> )	E	Wet meadows, open conifer stands	UCDS	UCDS
Golden paintbrush ( <i>Castilleja levisecta</i> )	T	Open grasslands, some partial shade	SW, IS	LC, SW, SP, NP, WP, IS
Water howellia ( <i>Howellia aquatilis</i> )	T	Wetlands, vernal pools	SP, SR, UCUS	UCUS, SR, SPS, SW, LC
Nelson's checker-mallow ( <i>Sidalcea nelsoniana</i> )	T	Streams, meadows, open areas	SW, LC	SW, LC, SPS
Kincaid's lupine ( <i>Lupinus sulphureus</i> spp. <i>kincaidii</i> )	T	Prairies, open oak woods	SW	SPS, LC, SW
Spalding's catchfly ( <i>Silene spaldingii</i> )	T	Open grasslands, scattered conifers	SR, UCUS, CB	SR, UCUS, CB
Slender moonwort ( <i>Botrychium lineare</i> )	C	Wide range: shaded and open	UCUS	UCUS, UCDS

<sup>1/</sup> E = endangered; T = threatened; C = candidate.

<sup>2/</sup> Analysis region(s) in which the species is currently found: NP = North Puget; WP = West Puget; SP = South Puget; OC = Olympic Coast; SW = Southwest; LC = Lower Columbia; MC = Middle Columbia; UCDS = Upper Columbia downstream of Grand Coulee Dam; UCUS = Upper Columbia upstream of Grand Coulee Dam; SR = Snake River.

<sup>3/</sup> Analysis region(s) in which the species could potentially occur: NP = North Puget; WP = West Puget; SP = South Puget; OC = Olympic Coast; SW = Southwest; LC = Lower Columbia; MC = Middle Columbia; UCDS = Upper Columbia downstream of Grand Coulee Dam; UCUS = Upper Columbia upstream of Grand Coulee Dam; SR = Snake River.

<sup>4/</sup> Species Is Extirpated In Washington.

Sources: NatureServe 2003; Washington DNR 2000.

3 wildlife, property values, public health and safety, and general ecological health and  
4 diversity of native ecosystems (<http://www.nwcb.wa.gov/weedlist/overview.html>).  
5 Washington State Noxious Weed Control Board 2003). In recognition of these multiple  
6 impacts, Washington's Weed Law (RCW Chapter 17.10) was updated in 1987 to include  
7 limiting economic loss and adverse effects to Washington's agricultural, natural, and  
8 human resources.

9 The State Noxious Weed Control Board systematically classifies noxious weeds based on  
10 the stage of invasion of each species. The classification system is designed to: 1) prevent  
11 small infestations from becoming large infestations, 2) contain already established  
12 infestations to regions of the state where they occur and prevent their movement to  
13 uninfected areas of Washington, and 3) allow flexibility at the local level to include  
14 widespread weeds for landowner management programs (Washington State Noxious Weed  
15 Control Board 2003). A complete noxious weed list is published annually in Chapter 16-  
16 750 of the WAC (<http://www.nwcb.wa.gov/>).



## 1 **3.7 RIPARIAN AND WETLAND PROCESSES**

### 2 **3.7.1 Riparian Areas**

3 Riparian habitat is the area adjacent to streams, rivers, lakes, and wetlands and includes  
4 floodplains and stream-associated seeps and springs. A wide variety of hydrologic,  
5 geomorphic, and biotic processes determine the character of riparian areas. Raedeke  
6 (1988) describes riparian systems as having long, linear shapes with high edge-to-area  
7 ratios and microclimates distinct from those of adjacent upland areas. Portions of riparian  
8 areas are disturbed from periodic inundation, and water is present at or near the soil surface  
9 during all or part of the year. These unique characteristics result in variable soil moisture  
10 conditions and distinct plant, animal, and invertebrate communities that are often more  
11 diverse than surrounding upland areas.

12 Riparian areas have distinct resource values and characteristics that make them important  
13 zones of interaction between terrestrial and aquatic ecosystems (Knutson and Naef 1997).  
14 These areas are especially dynamic segments of a watershed. Disturbance processes in  
15 upland areas (e.g., fire, mass wasting, and windthrow), as well as disturbance processes  
16 unique to stream systems (e.g., bank erosion, peak flows, floods) all affect riparian areas  
17 (Benda et al. 1998; Montgomery and Wohl 2004; Spence et al. 1996; Reeves et al. 1995).  
18 Functional riparian areas along streams and rivers in Washington are generally composed  
19 of large conifers, or a mixture of large conifers and hardwoods and a diverse understory  
20 plant community. Riparian vegetation is important for maintaining streambank and  
21 floodplain integrity. Vegetation slows water velocity on the floodplain, and plant roots  
22 inhibit erosion along streambanks, reducing sediment deposition in streams (subsection  
23 3.4, Geology, Soils, and Erosion). Riparian vegetation also helps to provide shade, leaf  
24 and needle litter important to aquatic food chains and nutrient cycling, and LWD, which is  
25 an important component of instream fish habitat (subsection 3.8, Fish and Fish Habitat).  
26 Riparian ecosystems are also important for a variety of plant and wildlife species. They  
27 provide the primary habitat for many of the State's amphibians (subsection 3.9,  
28 Amphibians and Amphibian Habitat) and provide important reproductive and foraging  
29 areas and/or dispersal/movement corridors for a wide variety of other wildlife (subsection  
30 3.10, Birds, Mammals, Other Wildlife and Their Habitats). Clearing or harvesting trees  
31 along streambanks and road construction near or adjacent to streams can negatively affect  
32 riparian ecosystem functions (Chamberlin et al. 1991; FEMAT 1993; Spence et al. 1996).

33 This section summarizes the primary functions of the riparian area. In addition, it provides  
34 a general description of the history of riparian management/riparian protection on private  
35 and State lands in Washington and the current riparian conditions in Washington State.  
36 Riparian vegetation was described in subsection 3.6.2 (Riparian Vegetation), and special  
37 disturbance problems in eastern Washington riparian areas were described in subsection  
38 3.6.3.4 (Special Problems in Eastern Washington Riparian Areas).

#### 39 **3.7.1.1 Riparian Functions**

40 To understand the effects of various management actions, it is important to understand the  
41 function of riparian areas, which have been described by many authors (e.g., Karr and



## Chapter 3

---

1 Schlosser 1977; Meehan et al. 1977; Raedeke 1988; Bilby 1988; Murphy and Meehan  
2 1991; Beschta 1991; Castelle et al. 1991). The most important recognized functions of  
3 stream riparian areas include LWD recruitment, leaf and needle litter production, stream  
4 shade, microclimate, streambank stability, groundwater recharge, stream energy  
5 dissipation, and the routing and trapping of sediment. Streambank stability and sediment  
6 filtration are introduced and evaluated in subsection 3.4 (Geology, Soils, and Erosion).  
7 The other riparian functions (LWD recruitment, leaf and needle litter production, stream  
8 shade, and microclimate) are briefly summarized below.

### 9 **3.7.1.2 LWD Recruitment**

10 Riparian areas are an important source of LWD that enters, or is recruited to, the stream  
11 channel. LWD includes entire trees, rootwads, and larger branches. Numerous studies  
12 have shown that LWD is an important component of fish habitat (Swanson et al. 1976;  
13 Bisson et al. 1987; Naiman et al. 1992). Trees that fall into streams are critical for  
14 sediment retention (Keller and Swanson 1979; Sedell et al. 1988), gradient modification  
15 (Bilby 1979), structural diversity (Ralph et al. 1994), nutrient production (Cummins 1974),  
16 and protective cover from predators. LWD also creates storage sites for sediment in all  
17 sizes of streams. In small headwater streams, wood controls sediment movement  
18 downstream minimizing the risk of debris flows. In large streams, accumulation of coarse  
19 sediment behind LWD often provides spawning gravels (Bilby and Bisson 1998;  
20 Montgomery et al. 2003). LWD plays an important role in stream nutrient dynamics by  
21 providing inputs of leaf litter and needles.

22 Large wood recruitment originates from a variety of processes including tree mortality,  
23 windthrow, undercutting of streambanks, debris avalanches, deep-seated mass soil  
24 movements, and redistribution from upstream (Swanson and Lienkamper 1978). First and  
25 second-order headwater streams can provide wood to larger higher order channels  
26 downstream (Potts and Anderson 1990; Prichard et al. 1998; Coho and Burges 1991).  
27 Wood can be transported from upstream during high flow events and from debris torrents,  
28 which includes dam-break floods and debris flows (Swanson and Lienkaemper 1978).  
29 However, high flow events are more common in third- to fifth-order streams (See  
30 Glossary) because much of the wood that falls into streams is too large to float in smaller  
31 streams (Swanson and Lienkaemper 1978). Although less frequent than high flow events,  
32 debris torrents can introduce large amounts of LWD (Lamberti et al. 1991). Debris flows  
33 originating in managed forests (albeit, under older less protective rules) occur at a rate  
34 much higher than that of unmanaged forests (Swanston and Swanson 1976; Morrison  
35 1975). The majority of debris flows and dam-break floods are initiated in low order  
36 streams, primarily second-order streams (Coho and Burges 1991). Debris flows can travel  
37 upwards of 2.5 miles into higher order, low gradient valley floors, and cause substantial  
38 damage to riparian vegetation and aquatic habitat during and after the event (Coho and  
39 Burges 1991). The destructive impacts of debris torrents can be reduced by maintaining  
40 contiguous riparian zones of mature conifers and minimizing deposition of logging slash  
41 into and along low order channels (Coho and Burges 1991). Wood and coarse sediment  
42 deposits from debris torrents can also form new habitat at tributary junctions by supplying



1 LWD that forms pools, cover, channel complexity, and by supplying spawning gravel  
2 (Benda et al. 2003).

3 The potential size distribution of LWD is also an important factor when considering the  
4 appropriate activities in riparian buffers relative to LWD recruitment. LWD that is large  
5 enough to form a pool is referred to as “functional LWD.” In contrast, “key piece LWD,”  
6 a subset of “functional LWD,” is considered to be a better measure of the important wood  
7 recruitment sizes. Key pieces are large and effective in trapping other smaller, more  
8 mobile of LWD (i.e., forming logjams). Key pieces of LWD are also more likely to have  
9 long-term stability (Bilby and Ward 1989; Collins et al. 2002; Hyatt and Naiman 2001).

10 Minimum functional LWD size increases with channel width (Bilby and Ward 1989; Bilby  
11 and Wasserman 1989; Beechie and Sibley 1997; Beechie 1998; Washington Forest  
12 Practices Board 1995). For example, mean LWD diameter increased from 11.7 inches in  
13 westside channels, which were 5 feet wide, to 21.7 inches in channels 44 feet wide (Bilby  
14 and Ward 1989). Key piece size is also related to stream size and is about 15 percent  
15 larger in diameter than functional piece size for a 40-foot wide stream (Washington Forest  
16 Practices Board 1995; Bilby and Ward 1989). As a result, riparian management zones  
17 need to ensure not only an appropriate amount or volume of wood, but wood of sufficient  
18 size to serve as both functional pieces and key pieces (Murphy 1995). Consequently, the  
19 length of time needed for riparian areas to produce LWD after harvest depends upon the  
20 size of the stream, stand composition, and site potential. Measurable contributions of  
21 wood from second-growth riparian areas are documented to take anywhere from 60 to 250  
22 years, depending on region and size of stream (Grette 1985; Bilby and Wasserman 1989;  
23 Murphy and Koski 1989). Therefore, large streams that are deficient in LWD and have  
24 adjacent and upstream riparian areas bordered by early seral stage riparian stands are likely  
25 to remain deficient in LWD longer than smaller streams because of their requirement to  
26 have large key pieces to retain LWD (MacDonald et al. 1991; Abbe and Montgomery  
27 2003). However, if numerous key piece size LWD were available in these wide stream  
28 reaches, and because LWD transport distance increases with stream size, these large  
29 streams may be able to increase LWD locally by capturing downstream-transported LWD  
30 in jams developed by key pieces (Abbe and Montgomery 2003; Martin and Benda 2001;  
31 Collins et al. 2002). However, the development of key pieces of LWD is infrequent in  
32 most early seral stage forests.

### 33 **3.7.1.3 Leaf and Needle Litter Production**

34 In aquatic systems, vegetative organic materials originate within the stream, such as algae  
35 production, or from sources outside the stream, such as leaf and needle litter. Stream  
36 benthic communities (e.g., aquatic insects) are highly dependent upon algal production and  
37 detrital (i.e., organic debris) inputs. The abundance and diversity of aquatic species can  
38 vary substantially depending upon the total and relative amounts of algae and leaf litter  
39 inputs to a stream (Independent Multidisciplinary Science Team 1999). For example,  
40 grazing insects are more commonly found in stream reaches with algae production, while  
41 shredding insects are more commonly found in areas rich in leaf and needle input  
42 (Independent Multidisciplinary Science Team 1999). Detrital input is the primary source  
43 of organic productivity in heavily-shaded, small and medium size streams (Gregory et al.



## Chapter 3

---

1 1991; Richardson 1992). In contrast, wide, high order streams with high levels of direct  
2 sunlight, or low order streams with an open riparian canopy, have more algal production.  
3 As a riparian stand ages, the amount of litter-fall increases (Independent Multidisciplinary  
4 Science Team 1999). The importance of detrital input varies among streams, but can  
5 provide up to 60 percent of the total energy input into stream communities (Richardson  
6 1992). Small (low-order) streams are important sources of nutrients and contribute  
7 substantially to the productivity of larger streams in the lower reaches of a watershed  
8 (Independent Multidisciplinary Science Team 1999).

### 9 **3.7.1.4 Stream Shade**

10 There are several factors that make up the heat balance of water (subsection 3.5.1, Surface  
11 Water Quality) including: air temperature, solar radiation, evaporation, convection,  
12 conduction, and advection (Brown 1983; Adams and Sullivan 1989). Stream temperatures  
13 have a natural tendency to warm from upstream to downstream in the watershed (Sullivan  
14 et al. 1990; Zwieniecki and Newton 1999). Seasonal and diurnal variability also exists  
15 with stream temperatures. Other site-specific factors such as latitude, regional  
16 climate/weather, stream size, groundwater inflow, and distance from watershed divides all  
17 can affect stream temperature (Beschta et al. 1987; Sullivan et al. 1990). During the  
18 summer when stream temperatures are the highest, the combination of warmer air  
19 temperatures, increased direct solar radiation and decreased stream flows are the major  
20 factors affecting stream temperature (Beschta et al. 1987). Of these three factors, forest  
21 management can have the greatest effect on direct solar radiation by reducing shade.  
22 Shade alone does not physically cool the stream, but it prevents further heating of the  
23 stream and maintains the cool water temperatures from groundwater inputs or tributaries  
24 (Oregon Forest Practices Advisory Committee on Salmon and Watersheds 2000). Shade  
25 from riparian vegetation has been shown to minimize or eliminate increases in stream  
26 temperature associated with timber harvest (Brazier and Brown 1973; Lynch et al. 1985).  
27 Other factors that affect shading include local topography, stream size and aspect, stand  
28 age, composition, and stand density.

### 29 **3.7.1.5 Microclimate**

30 Microclimate is a collection of variables that are highly dependent on local conditions;  
31 hence, microclimates tend to vary greatly across the landscape. Important components of  
32 microclimate include solar radiation, soil temperature, soil moisture, air temperature, wind  
33 velocity, and air moisture or humidity (Chen 1991; Chen et al. 1992; Cadenasso et al.  
34 1997). Changes in microclimatic conditions within the riparian zone resulting from  
35 removal of adjacent vegetation can influence a variety of ecological processes that may  
36 affect the long-term integrity of riparian ecosystems (Spence et al. 1996). For example,  
37 many of the variables considered in microclimate studies (air temperature, humidity, wind  
38 velocity) are also variables that affect water temperature (Sullivan et al. 1990), an  
39 important component of fish habitat. Additionally, microclimate is known to be important  
40 for aquatic/riparian species other than fish, such as amphibians (subsection 3.9,  
41 Amphibians and Amphibian Habitat). In general, due to the low-lying position on the  
42 landscape, riparian areas tend to be cooler than the surrounding hillslopes, especially  
43 during the night. Because riparian areas are adjacent to water bodies, they often have a



1 higher relative humidity under the canopy than upslope areas. This increase in humidity,  
2 combined with shading effects, can result in forested riparian areas creating a moderating  
3 effect on microclimate (Beschta 1991).

### 4 **3.7.1.6 Historic Protection of Riparian Areas**

5 The protection of riparian areas is considered critical to the long-term health of aquatic  
6 ecosystems and salmonid conservation efforts (FEMAT 1993; Cederholm 1994; Murphy  
7 1995). Riparian areas are protected by restricting management activities within the  
8 riparian management zone (RMZ). Management within RMZs usually involves two main  
9 features: 1) establishment of a buffer width, and 2) restrictions on allowable activities (e.g.,  
10 timber harvest prescriptions) within the buffer. Protection of water quality and fish habitat  
11 is often given the highest management priority; however, buffers may also be designed to  
12 benefit wildlife and other non-fish aquatic species.

13 Washington Forest Practices Rules have consistently been reviewed and revised in light of  
14 new scientific information, changing public awareness, and evolving demands for forest  
15 and water resources. The Washington State Forest Practices Act of 1974 created a Forest  
16 Practices Board, which promulgates Washington Forest Practices Rules. Early riparian  
17 rules only considered changes in stream temperature and bank stability for the aquatic  
18 ecosystem. All riparian trees could be cut, sparing only the understory on certain  
19 temperature-sensitive streams. Since then the Washington Forest Practices Rules have  
20 undergone numerous revisions. Sweeping changes occurred in 1987 with the TFW  
21 Agreement and rule changes resulting from that agreement. Significant revisions were also  
22 made in 1992, 1996, and the most recent revisions, based on the FFR, in 2001 (subsection  
23 1.3.1, Forest Practices Program, and subsection 2.3, Alternatives Analyzed in Detail).

### 24 **3.7.1.7 Current Condition of Riparian Areas**

25 Current riparian conditions on State and private lands are mostly a function of past forest  
26 management practices, but natural phenomenon such as wildfire, blowdown, non-  
27 management related landslides, and disease have also contributed to conditions in many  
28 areas. Because riparian protection rules are a relatively recent phenomenon in Washington  
29 State (1982), the majority of riparian forests on State, private, and some Federal lands have  
30 been logged at least once. Therefore, long-term changes to the riparian habitat character  
31 have resulted from multiple forest practices over time. These changes to riparian habitat  
32 structure include simplification of the plant community, both in species composition and  
33 structure (Knutson and Naef 1997). Today it is believed that red alder dominates more  
34 riparian sites in western Washington than was “typical” under natural disturbance regimes  
35 (Washington Forest Practices Board 2001a; McHenry et al. 1998).

36 Two studies, described below, classified streamside vegetation into three seral stages to  
37 present a picture of current riparian vegetation conditions. Seral stage provides a general  
38 picture of riparian condition and quality. The two studies cannot be directly compared to  
39 each other because each study used its own definitions, study area, and data  
40 collection/analysis methods. One study sampled lands subject to Washington Forest  
41 Practices Rules throughout Washington, while the other study area was specific to western



## Chapter 3

1 Washington riparian areas. Both studies, however, provide information that indicates the  
2 current general condition of Washington's riparian areas.

3 The first study was conducted to provide information for the recent Forest Practices  
4 Alternatives SEPA EIS (Washington Forest Practices Board 2001a). Table 3-18 presents  
5 the percentage of streamside vegetation on private lands subject to Washington Forest  
6 Practices Rules that existed in the three seral stages (based on a random sample of lands).  
7 Seral stages were based on tree sizes as follows:

- 8 • Early Seral – Reproduction, conifer pole, hardwood pole, and mixed pole; less than 12  
9 inches in diameter at breast height (dbh).
- 10 • Mid-Seral – Conifer sawtimber, hardwood sawtimber, and mixed sawtimber; 12 to 24  
11 inches in dbh.
- 12 • Late Seral – Large conifer sawtimber, large hardwood sawtimber, and large mixed  
13 sawtimber; greater than 24 inches in dbh.

14 The Forest Practices Board study concluded that unnaturally high levels of early seral stage  
15 vegetation existed in riparian zones on private forestland, primarily as a result of timber  
16 management activities and, to a lesser extent, from fire, blowdown, and other natural  
17 processes in riparian areas (Washington Forest Practices Board 2001a). In general, early  
18 seral stages produce riparian vegetation that provides lower riparian values for aquatic and  
19 terrestrial biota. In contrast, later seral stages that are typically more diverse in species  
20 composition and stand structure can more fully provide for riparian functions (e.g., shade  
21 and LWD recruitment for aquatic biota) (subsection 3.8, Fish and Fish Habitat) and more  
22 complex vegetative structure (e.g., downed logs and snags for terrestrial biota associated  
23 with riparian habitat) (subsection 3.9, Amphibians and Amphibian Habitat, and subsection  
24 3.10, Birds, Mammals, Other Wildlife and Their Habitats).

25 These data indicate that within the lands subject to Washington Forest Practices Rules up  
26 to 2001, approximately 78 percent of western Washington stream miles and 61 percent of  
27 eastern Washington stream miles flow through early seral stage riparian areas, while about

28 **Table 3-18.** Estimated Percent of Each Seral Stage along Forested Streams on  
29 Private Lands. <sup>1/</sup>

Water Type	Seral Stage - Percent by Water Type (%)		
	Early	Mid	Late
<b>Westside-Private Lands</b>			
Types 1-3	64	33	2
Types 4-5	81	18	1
All Streams	78	21	1
<b>Eastside- Private Lands</b>			
Types 1-3	60	36	4
Types 4-5	61	33	6
All Streams	61	34	5

Source: Washington Forest Practices Board 2001a, Appendix C, Table 6. Based on a random sample of private lands (See DEIS Appendix B for a description of the sampling). See text for seral stage definitions.





1 1 percent of western Washington miles and 5 percent of eastern Washington miles are late  
2 seral. Though natural variability is expected in riparian areas, the level of alteration due to  
3 timber harvest and road building is apparent.

4 The second study (Lunetta et al. 1997) used digital elevation modeling of stream channels  
5 to determine channel gradient, and then characterized riparian seral stage by gradient  
6 category. The analysis looked at forest vegetation in 179 watersheds across western  
7 Washington. Stream channels were classified into three categories based on channel slope:  
8 response reach, transport reach, and source reach. The response reach seral stage data are  
9 the only raw data still available from this study and is, thus, the only data reported here.  
10 Response reaches were defined as channel reaches with less than 4 percent slope, and were  
11 considered the area where most anadromous fish production occurs.

12 Riparian response reaches were broken into three seral stages defined as follows:

- 13 • Early Seral – Hardwood dominated, shrub, or recent clearcut. Also includes forests  
14 with greater than or equal to 10 percent and less than 70 percent conifer crown cover,  
15 and less than 75 percent of total crown cover in hardwood tree/shrub cover.
- 16 • Mid-Seral – Forests with greater than 70 percent conifer crown cover, less than 10  
17 percent crown cover in trees greater than or equal to 21 inches dbh.
- 18 • Late Seral – Forests with greater than 70 percent conifer crown cover, more than 10  
19 percent of the crown cover must be in trees greater than 21 inches dbh.

20 Response reach seral stage data are summarized by analysis region in Table 3-19. Again,  
21 as of 1997, the most common seral stage was early seral, ranging from 52 percent in the  
22 West Puget Sound Region to 72 percent in the Lower Columbia Region. Late seral made  
23 up the lowest percentage, ranging from 5 percent in the Southwest and South Puget Sound  
24 Regions to 19 percent in the North Puget Sound Region.

25 Riparian condition and function (e.g., floodplain condition, bank stability, LWD, shade and  
26 stream temperature, water quality) vary from good to poor within each region, depending  
27 on site scale and location. For a more detailed analysis, see the Regional Summaries in  
28 DEIS Appendix A.

### 29 **3.7.2 Wetlands**

30 Wetlands are defined in terms of their physical, chemical, and biological characteristics,  
31 such as hydrologic regime, soil type, and plant species. Wetlands are formally defined as  
32 those areas that are inundated or saturated with surface or groundwater at a frequency and  
33 duration sufficient to support, and that under normal circumstances, do support a  
34 prevalence of vegetation typically adapted for life in saturated soil conditions (40 CFR  
35 230.41(a)(1); WAC 222-16-11). This definition includes forested swamps, marshes, bogs,  
36 and other similar areas. Wetlands are subject to regulation through the Clean Water Act by  
37 the U.S. Army Corps of Engineers and EPA. Sections 404 and 401 of the Clean Water Act



## Chapter 3

1

2 **Table 3-19.** Estimated Percent of Each Seral Stage along Response Reaches of  
3 Forested Streams for Western Washington Analysis Regions.

		Seral Stage		
		Early	Mid	Late
Analysis Region (Acres/Percent)		Conifer Crown Cover Between 10 and 70%	Greater than 70% Conifer Canopy; 10% or Less of the Canopy in Conifers Greater than 21"dbh	Greater than 70% Conifer Canopy; More than 10% of the Canopy in Conifers Greater than 21"dbh
North Puget Sound	Acres	55,704	17,323	16,808
	Percent	62	19	19
South Puget Sound	Acres	34,829	18,613	2,938
	Percent	62	33	5
West Puget Sound	Acres	27,416	17,994	7,157
	Percent	52	34	14
Olympic Coast	Acres	47,897	26,226	14,869
	Percent	54	29	17
Southwest	Acres	103,810	46,893	8,582
	Percent	65	29	5
Lower Columbia	Acres	69,833	17,421	10,080
	Percent	72	18	10

Source: Personal Communication, Brian Cosentino, Washington Department of Fish and Wildlife, October 2003.

4

5 were created specifically with the intent “to restore and maintain the chemical, physical,  
6 and biological integrity of our nation’s waters.”

7 Wetland ecosystems provide a variety of physical and biological functions. Additionally,  
8 they provide many values to society including recreation, water quality enhancement, and  
9 flood attenuation. The National Wetland Policy Forum (Conservation Foundation 1988)  
10 identified eight natural functions that wetlands may perform at a landscape level. These  
11 eight functions are: 1) nutrient removal and transformation, 2) sediment and toxicant  
12 retention, 3) shoreline and bank stabilization, 4) flood flow alteration, 5) groundwater  
13 recharge, 6) nutrient export, 7) aquatic diversity and abundance, and 8) wildlife diversity  
14 and abundance. Values to society include recreation, water quality enhancement, and flood  
15 control.



### 1 **3.7.2.1 Wetland Functions**

2 As noted above, wetlands provide a variety of functions and values. The key wetland  
3 functions that are the focus of this DEIS include fish and wildlife habitat, water quality,  
4 and hydrology. These functions were chosen because they may be most directly impacted  
5 by timber harvest related activities. The functions are briefly discussed below.

#### 6 **Fish and Wildlife Habitat**

7 Wetland and riparian habitats are considered to be among the richest zones for aquatic and  
8 terrestrial organisms (Dodd 1978; Brinson et al. 1981; Kauffman and Krueger 1984).  
9 Because wetland and riparian habitats exhibit an “edge effect” due to overlapping types of  
10 habitats, these areas provide more niches than are provided by any other habitat types.  
11 Eighty-six percent (359 out of 414) of the terrestrial vertebrate species in western  
12 Washington, and 85 percent (320 out of 378) of terrestrial vertebrate species in eastern  
13 Washington utilize wetland and associated riparian habitats for portions of their life needs  
14 (Brown 1985; Thomas 1979).

15 Wetlands provide habitat or perform functions that contribute to the health of ecosystems  
16 of many anadromous and resident fish species within Washington. Wetlands are known to  
17 help maintain cool water temperatures, retain sediments, store and desynchronize flood  
18 flows, maintain stream base flows, and provide food and cover for fish (Cederholm and  
19 Scarlett 1981; Beechie et al. 1994; Mitsch and Gosselink 1993; Washington Department of  
20 Ecology 1993b).

#### 21 **Water Quality**

22 Wetlands can improve water quality through nutrient removal and transformation  
23 (Hammer 1989). For example, wetlands can remove nitrate and phosphorus from  
24 agricultural runoff. Nutrient-rich sediments may also become trapped and removed from  
25 the water. Wetlands can also remove toxic chemicals, such as pesticides, heavy metals, or  
26 excess nutrients from water (Mitsch and Gosselink 1993). Wetlands can reduce shoreline  
27 and bank erosion by binding soil substrates in wetland plant roots. Thus, wetlands protect  
28 upland habitats along streams and rivers from erosion, and protect downstream habitats  
29 from sedimentation and pollution. Wetlands, which discharge cool groundwater, can help  
30 maintain desirable stream temperatures in the summer. Forested riparian and wetland areas  
31 serve an important role in shading streams from direct solar heating. Other wetlands,  
32 without cool groundwater discharge, may be a source of warmer water to stream  
33 temperatures, but they also provide fish habitat and a source of nutrients.

#### 34 **Hydrology**

35 Headwater riverine and depressional wetlands can delay discharge of peak run-off into  
36 streams and impede passage of overbank flow downstream during storm events, thus  
37 reducing the potential for downstream flooding (Winter 1988; Roth et al. 1993).  
38 Depressional wetlands can help maintain minimum stream base flow by naturally  
39 regulating the release of groundwater discharge into streams and by recharging aquifers  
40 that discharge groundwater to streams (Dinicola 1990; Hidaka 1973; O’Brien 1988; Mitsch  
41 and Gosselink 1993).



## Chapter 3

---

### 3.7.2.2 Historic/Current Wetland Protection

Wetlands are subject to regulation under Sections 401 and 404 of the Clean Water Act. Discharge into wetlands may also be regulated under Section 402 of the Clean Water Act. Exemptions granted under Section 404(f)(1) allow for normal agricultural, ranching, and silvicultural activities, as well as maintenance of existing drains, farm ponds, and roads. The construction or maintenance of forest roads for silvicultural purposes is exempt from regulation when such roads are constructed and maintained in accordance with BMPs.

On State and private lands in Washington, the Washington Forest Practices Rules provide wetland resource protection from timber harvest-related activities. For management purposes, the Washington Forest Practices Rules recognize two major categories of wetlands: forested and non-forested. Non-forested wetlands are divided further into two classes: Type A (greater than 0.5 acre, with open water) and Type B (other non-forested wetlands). The Washington Forest Practices Rules require buffers, termed Wetland Management Zones, on all Type A wetlands and most Type B wetlands. Harvest may occur in forested wetlands; however, harvest methods are limited to low impact harvest or cable systems.

For Type A wetlands greater than 5 acres in size, an average Wetland Management Zone of 100 feet is required. For Type A wetlands between 0.5 and 5 acres, a 50-foot average Wetland Management Zone is required. For Type B wetlands greater than 5 acres, a 50-foot average Wetland Management Zone is required. For other wetlands between 0.5 and 5 acres, a 25-foot Wetland Management Zone is required. Wetlands less than 0.5 acre have no buffer requirement.

In addition to leaving Wetland Management Zones, there are several other harvest restrictions around non-forested wetlands. For example, individual trees and small patches of forested wetlands (0.5 acre) cannot be harvested if surrounded by a Type A or Type B wetland. Harvest of upland areas or large forested wetlands require a plan approved by Washington DNR if they are surrounded by Type A or Type B wetlands. Additionally, timber cannot be felled into or cable-yarded across a Type A or Type B wetland without prior approval by Washington DNR.

### 3.7.2.3 Current Condition of Wetlands

Since the time of colonization, Washington State has lost between 30 to 50 percent of its wetlands (USFWS 1999). Additionally, the functions of existing wetlands have been reduced. Various factors have contributed to wetland loss and wetland function reduction including agriculture development, urbanization, timber harvest, road construction, and other land management activities. It is difficult to assess the current condition of wetlands in forested lands across the entire State of Washington. However, some wetlands on lands subject to Washington Forest Practices Rules have been altered in the past due to timber harvest and road building. These actions can impact wetland sites directly through vegetation alteration, soil compaction, changes in hydrologic regime, and degradation of water quality; or indirectly through sedimentation from adjacent land management practices. Additionally, harvest of trees in or adjacent to wetland sites can impact the

---

## Chapter 3



1 associated microclimate (Brosofske et al. 1997; Chen et al. 1995, 1999). Other impacts to  
2 wetlands have likely occurred from fires and other natural disturbances.

3 Washington DNR wetland GIS coverage was used to generate the data reported in Table 3-  
4 20. Overall, approximately 4.4 percent of the forestland base subject to the Washington  
5 Forest Practices Rules evaluated is comprised of wetland habitats, based on this mapping.  
6 Wetland areas comprise approximately 2 percent of the land base in eastern Washington  
7 and approximately 6 percent in western Washington.

8 **Table 3-20.** Approximate Wetland Area as a Percentage of Forested  
9 Ownership, by Region and Wetland Type.

Region	Ownership	Type A Wetland (%)	Type B Wetland (%)	Other Open Water (%)	Forested Wetland (%)
Westside	Private Lands	0.7	<0.1	<0.1	5.7
Eastside	Private & State Lands	0.4	<0.1	<0.1	1.2
Statewide	Private and State on Sampled lands	<b>0.6</b>	<b>&lt;0.1</b>	<b>&lt;0.1</b>	<b>3.8</b>


Note: < = less than

Source: Washington Forest Practices Board 2001a, Section 3.5.2.3, Table 3.5-1. Based on a random  
sample of lands subject to Washington Forest Practices Rules in each area/ownership category.

---

10

11



## **Chapter 3**

---

1 This page is intentionally left blank.

2



## 1 **3.8 FISH AND FISH HABITAT**

### 2 **3.8.1 Introduction**

3 Fish are an important natural resource with both biological and economic significance in  
4 the State of Washington. In particular, Pacific salmon and trout, as well as other fish  
5 species, are indicators of a properly functioning aquatic ecosystem because they require  
6 cool, clean water, complex channel structures and substrates, and low levels of silt (Bjornn  
7 and Reiser 1991). In addition, Pacific salmon and trout support economically important  
8 commercial and sport fishing industries, as well as subsistence fishing by many  
9 Washington Indian Tribes.

10 This section discusses the fish species in Washington and their habitats. A complete list of  
11 all species that are intended to be “covered” by the FPHCP is provided in Table 1-1.  
12 Chapter 3 of the FPHCP presents life history and status information for all of these species.  
13 Those fish species with the more critical Federal status of “endangered” or “threatened” are  
14 given the most attention within this section; however, other fish species with less critical  
15 Federal or State status are also described. Further, this section describes important  
16 components of the aquatic environment that fish require and that forest practices may  
17 effect. These components include water quality, water quantity, channel conditions, LWD,  
18 channel morphology, and fish passage.

19 The following subsections emphasize the affected environment for fish species on State  
20 and private lands within Washington State, which are regulated by Washington Forest  
21 Practices Rules. The discussion contains a review of fish distribution and status within the  
22 12 analysis regions defined in subsection 3.1 (Introduction). This section also contains a  
23 review of important components of the aquatic ecosystem upon which fish rely for  
24 sustaining healthy, well-dispersed populations.

### 25 **3.8.2 Fish Status in Washington**

26 More than 70 species of freshwater fish are present in the more than 8,000 lakes and  
27 50,000 miles of streams within Washington (Wydoski and Whitney 2003). Generally, at  
28 least one fish species is found in perennial streams with gradients less than 20 percent  
29 (Fransen et al. 1997). Occasionally, fish are found in streams with steeper gradients, but  
30 these circumstances are less common. Land-use practices upstream of fish-bearing waters  
31 can affect downstream fish habitat. Consequently, the affected environment for fish  
32 includes both fish-bearing and non-fish-bearing streams.

33 Two of the four goals of the Forest Practices Board for the Washington Statewide Salmon  
34 Recovery Strategy (Washington Forest Practices Board 1999) have special reference to fish  
35 and forestry interactions. One of the goals is to provide compliance with the Endangered  
36 Species Act (ESA) for aquatic and riparian-dependant species on all lands subject to the  
37 Forest Practices Act. A second goal is to restore and maintain riparian habitat on these  
38 forestlands to support a harvestable supply of fish.

39 Notably, NMFS has not listed any Pacific salmon or trout species as threatened or  
40 endangered throughout their entire range, because many populations within the entire range



## Chapter 3

---

1 are considered healthy or at least stable. Instead, NMFS has listed specific salmon and  
2 trout stocks based upon distinct populations that are “substantially reproductively isolated”  
3 and “represent an important component in the evolutionary legacy of the species” (Waples  
4 1991). NMFS has termed these populations “Evolutionarily Significant Units.”  
5 Analogously, the USFWS has chosen to use the term “Distinct Population Segments” for  
6 freshwater fish species under its regulatory authority.

7 Beginning in 1991 with the listing of Snake River sockeye salmon by NMFS, the ESA has  
8 increasingly affected the way government agencies and public and private landowners  
9 conduct business in or near the streams and rivers found in the State. The rate of new  
10 listings escalated in the late 1990s such that at least one Evolutionarily Significant Unit of  
11 all of the Pacific salmon species, with the exception of pink salmon, have been listed as  
12 threatened or endangered within one or more areas of Washington State (Table 3-21). In  
13 addition to the Pacific salmon and trout listed by NMFS, the USFWS has listed bull trout  
14 throughout its range in the contiguous United States. Consequently, there are few areas  
15 within Washington State that do not have at least one listed fish species (Figures 3-4  
16 through 3-9).

### 17 **3.8.3 Life History of Covered and Affected Fish Species**

18 A basic understanding of the life history and habitat requirements of the various covered  
19 fish species is important for recognizing the type and level of effects that may result from  
20 forest practices. The life history characteristics can vary substantially in different locations  
21 depending on climate, food supply, stream flow, and other factors (Flosi and Reynolds  
22 1994).

#### 23 **3.8.3.1 Pacific Salmon and Trout - General**

24 The life cycle of anadromous Pacific salmon and trout can be divided into seven distinct  
25 phases or life stages: upstream migration, spawning, egg incubation, fry emergence,  
26 juvenile rearing, smolt outmigration, and marine rearing. Two important common  
27 denominators in the life history of Pacific salmon and trout is they all construct redds  
28 (nests) in gravel beds for spawning, and they all include life history forms that exhibit  
29 anadromy. In other words, spawning occurs in freshwater, followed by migration to the  
30 ocean for feeding and maturation, and finally fish return to their natal sites for completion  
31 of the life cycle. Four of the salmon and trout species also have life history forms that live  
32 their entire lives in freshwater (i.e., sockeye/kokanee, steelhead/rainbow, cutthroat, and  
33 bull trout). Several anadromous species demonstrate extremely complex variations in  
34 length of freshwater rearing, use of lake systems, run timing, degree of anadromy, and age  
35 structure. These variations, in conjunction with geographically separate spawning  
36 populations, have led to the stock concept of salmon management (Larkin 1972). It is the  
37 demonstration of unique behavioral patterns, physical characteristics, and ultimately  
38 genetic makeup that has made it possible to list any salmon stock within the framework of  
39 the ESA (Nehlsen et al. 1991; Waples 1991).



# Chapter 3



**Table 3-21.** Covered Fish Species with Federal or State Listed Status in Washington State.

Species	Population <sup>1/</sup>	Federal Status	State Status	Distribution <sup>2/</sup>
Chum Salmon	Hood Canal Summer-run	Threatened	Candidate	5
<i>Oncorhynchus keta</i>	Columbia River	Threatened	Candidate	3
	Puget Sound—Strait of Georgia	Species of Concern	None	1, 4-7
Coho Salmon <i>O. kisutch</i>	Lower Columbia River	Candidate	None	3
	Southwest Washington	Species of Concern	None	2
Sockeye Salmon <i>O. nerka</i>	Snake River	Endangered	Candidate	12
	Ozette Lake	Threatened	Candidate	1
Chinook Salmon <i>O. tshawytscha</i>	Snake R. Fall-run	Threatened	Candidate	12
	Snake R. Spring/Summer-run	Threatened	Candidate	12
	Puget Sound	Threatened	Candidate	4, 5, 7
	Lower Columbia River	Threatened	Candidate	3, 9
	Upper Willamette R.	Threatened	Candidate	9
Steelhead Trout <i>O. mykiss</i>	Upper Columbia River Spring-run	Endangered	Candidate	8
	Upper Columbia River	Endangered	Candidate	8
	Snake River	Threatened	Candidate	12
	Lower Columbia River	Threatened	Candidate	3,9
	Upper Willamette	Threatened	Candidate	9
Bull Trout <i>Salvelinus confluentus</i>	Middle Columbia River	Threatened	Candidate	9
	Columbia River	Threatened	Candidate	3, 8-10, 12
Coastal Cutthroat Trout <i>O. clarki clarki</i>	Coastal - Puget Sound	Threatened	Candidate	1, 2, 4, 5, 7
		Species of Concern	None	1-7
Westslope Cutthroat Trout <i>O. clarki clarki</i>		Species of Concern	None	8-10, 12
Interior Redband Trout		Species of Concern	None	8-10, 12
Green Sturgeon <i>Acipenser medirostris</i>		Species of Concern	None	2-5, 7
Pacific Lamprey <i>Lampetra tridentate</i>		Species of Concern	None	1-5, 8, 9, 12
River Lamprey <i>L. ayresi</i>		Species of Concern	Candidate	3-5, 7, 9, 12



## Chapter 3

**Table 3-21.** Covered Fish Species with Federal or State Listed Status in Washington State (continued).


Species	Population <sup>1/</sup>	Federal Status	State Status	Distribution <sup>2/</sup>
Margined Sculpin <i>Cottus marginatus</i>		Species of Concern	Sensitive	12
Pygmy Whitefish <i>Prosopium coulteri</i>		None	Sensitive	1, 4, 8-10
Olympic Mudminnow <i>Novumbra hubbsi</i>		None	Sensitive	1, 2, 4, 5
Umatilla Dace <i>Rhinichthys Umatilla</i>		None	Candidate	8-10
Leopard Dace <i>R. falcatus</i>		None	Candidate	3, 8, 9
Lake Chub <i>Couesius plumbeus</i>		None	Candidate	10
Eulachon <i>Thaleichthys pacificus</i>		None	Candidate	1-3, 7
Mountain Sucker <i>Catostomus platyrhynchus</i>		None	Candidate	3, 8, 9, 12
Salish Sucker <i>C. carli</i>		None	Monitor	4, 5, 7
Sandroller <i>Percopsis transmontana</i>		None	Monitor	3, 8-10, 12

<sup>1/</sup> Populations of Pacific salmon are designated as Evolutionarily Significant Units (ESU) by NMFS. The USFWS designates threatened and endangered population segments as Distinct Population Segments (DPS).


<sup>2/</sup> Numbers indicate EIS Regions where species occurs. Region: 1 = Olympic Coast; 2 = Southwest; 3 = Lower Columbia; 4 = South Puget Sound; 5 = West Puget Sound; 6 = Islands; 7 = North Puget Sound; 8 = Upper Columbia (downstream Grand Coulee); 9 = Mid Columbia; 10 = Upper Columbia (upstream Grand Coulee); 11 = Columbia Basin; 12 = Snake River.



1 **Figure 3-4.** Distribution and ESA Status of Chinook Salmon within Washington  
2 State (Source: Streamnet Version 99.1; NMFS -  
3 <http://www.nwr.noaa.gov/1salmon/Salmesa>).



4 **Figure 3-5.** Distribution and ESA Status of Chum Salmon within Washington  
5 State (Source: Streamnet Version 99.1; NMFS -  
6 <http://www.nwr.noaa.gov/1salmon/Salmesa>).






# Chapter 3

1 **Figure 3-6.** Distribution and ESA Status of Coho Salmon within Washington  
2 State (Source: Streamnet Version 99.1; NMFS -  
3 <http://www.nwr.noaa.gov/1salmon/Salmesa>).




4 **Figure 3-7.** Distribution of ESA Status of Sockeye Salmon within Washington  
5 State (Source: Streamnet Version 99.1; NMFS -  
6 <http://www.nwr.noaa.gov/1salmon/Salmesa>).  
7






1  
2  
3  
4

**Figure 3-8.** Distribution and ESA Status of Steelhead within Washington State  
(Source: Streamnet Version 99.1; NMFS - <http://www.nwr.noaa.gov/1salmon/Salmesa>).



5  
6  
7

**Figure 3-9.** Distribution and ESA Status of Bull Trout within Washington State  
(Source: Streamnet 2002; USFWS Draft Recovery Plan 2004b).





## Chapter 3

---

1 One commonly recognized variation in life history traits for Pacific salmon and steelhead  
2 is run timing. The seasonal stock distinctions are based upon the date individual stocks of  
3 maturing adults enter freshwater. For example, chinook salmon are often divided into  
4 “spring,” “summer,” and “fall” runs, while steelhead stocks are divided into “winter” and  
5 “summer” runs. Sockeye usually do not have multiple distinct runs and the seasonal  
6 descriptor is often omitted. Most pink salmon in the Puget Sound Region enter freshwater  
7 during the fall while sockeye salmon runs peak in early July. Chum salmon exhibit  
8 summer, fall, and winter run timing.

9 Additional stock and species-specific variability is demonstrated in the duration of  
10 freshwater rearing and the type of habitat that is utilized. Spring chinook salmon, coho  
11 salmon, and steelhead juveniles typically spend 1 or 2 years rearing in streams prior to  
12 outmigration. Similarly, sockeye salmon usually spend a year rearing in a lake prior to  
13 outmigration. In contrast, ocean-type chinook and chum, and pink salmon outmigrate to  
14 the ocean as fry. Chum salmon usually complete their outmigration shortly after  
15 emergence (Wydoski and Whitney 2003), while fall chinook may have a protracted  
16 outmigration period that occurs throughout the summer (Dawley et al. 1986). While most  
17 summer/fall chinook outmigrate during their first year, a small proportion overwinter in  
18 freshwater and then migrate as yearlings the following spring.

19 Bull trout and coastal cutthroat trout also express high variability in migratory behavior  
20 and habitat use. They have four different migratory forms: anadromous, adfluvial, fluvial,  
21 and resident. Adfluvial stocks rear in lake systems, but migrate to tributary streams for  
22 spawning. Fluvial stocks rear entirely in larger streams or rivers, but have lengthy  
23 migrations between headwater spawning and rearing areas. Other trout subspecies such as  
24 westslope cutthroat and redband trout are solely resident and do not exhibit anadromy.  
25 Though resident stocks demonstrate little “migratory” behavior (compared with  
26 anadromous fish), connectivity is critical throughout their specific habitat ranges in order  
27 to fulfill all of their life history needs (i.e., spawning, rearing, foraging, escaping extreme  
28 environmental fluctuations, and recolonization after habitat disturbance).

29 For freshwater spawning and rearing, salmon and trout (as well as other freshwater fish  
30 species) have life-stage and species-specific habitat requirements. Important components  
31 to spawning habitat include substrate size, water depth, and water velocity (Bjornn and  
32 Reiser 1991). In general, the larger species utilize larger substrates and deeper and faster  
33 water (0.5 to 4 in. [1.3 to 10.2 cm], greater than 9 in. [24 cm] depth, 1.1 to 3.6 ft/sec [32 to  
34 109 cm/sec] velocity) (Bjornn and Reiser 1991). Tail-outs to pools (the downstream end  
35 where the pool changes to a riffle) that meet criteria for these features are generally  
36 considered optimal spawning areas for salmonids because stream morphology maximizes  
37 the passage of oxygenated water through redds. Chums commonly utilize spring or  
38 groundwater upwelling areas for spawning (Salo 1991). However, runs, riffles, and  
39 groundwater or spring upwelling zones along lakeshores and intertidal sloughs are also  
40 utilized during spawning. Redd building is important for three principle reasons (Chapman  
41 1988): 1) redds provide physical protection to eggs during periods when they are extremely  
42 fragile; 2) redd digging removes a portion of the fines and sands deleterious to egg



1 survival; and 3) redd construction and morphology enhances the passage of water through  
2 the egg pockets.

3 Following emergence from the redd, salmon and trout fry typically utilize shallow and  
4 slow moving areas of a stream (Bjornn and Reiser 1991). Optimal depths and velocities  
5 increase as the fish grow, but preferred areas are usually associated with some form of  
6 cover, usually pools with LWD or boulders, or with faster water areas. Differences among  
7 the species are apparent in the degree of flexibility for utilizing riffles, runs, and other  
8 habitat features. Stream dwelling juvenile salmonids are typically territorial and exhibit a  
9 dominance hierarchy among individuals and species. Drifting insect larvae and benthic  
10 macroinvertebrates account for the majority of food items eaten by juvenile salmon and  
11 trout within streams. In contrast to the typical stream dweller, sockeye fry migrate to a  
12 lake shortly after emergence where shallow nearshore (or littoral) areas are preferred  
13 habitat. As sockeye fry grow, they begin to move offshore and have a characteristic  
14 diurnal vertical migration timed for utilization of zooplankton food sources (Bjornn and  
15 Reiser 1991; Burgner 1968; Groot and Margolis 1991; Wydoski and Whitney 2003).

16 Riparian areas have distinctive resource values and characteristics that are critical to  
17 salmonid production (FEMAT 1993). Riparian vegetation is important for maintaining  
18 streambank and floodplain integrity. The vegetation slows water velocity on the floodplain  
19 and roots inhibit erosion along streams and riverbanks, which reduces sediment deposition  
20 in streams. Riparian vegetation also helps to provide shade (important for the maintenance  
21 of stream temperatures), leaf, and needle litter important to aquatic food chains and LWD  
22 (FEMAT 1993). Riparian vegetation and LWD also traps and retains salmon carcasses, an  
23 important source of marine-derived nitrogen (Bilby et al. 1996).

24 In general, the marine phase of salmonid life history is not well understood. Only recently  
25 have ocean environmental conditions been considered an important factor in the  
26 management of salmon resources (Bisbal and McConnaha 1999). Historically, the ocean  
27 was assumed an unlimited resource for salmon production, but this assumption is now  
28 being widely questioned. Forest practices are likely to have very minimal direct effects on  
29 anadromous fish once they are in the marine environment, since many other factors such as  
30 tides, currents, nutrients, marine food sources, marine predators, become paramount in  
31 their growth and survival. However, forest practices may affect woody debris and  
32 sediment composition and supply in estuarine and nearshore environments, which have the  
33 potential to affect rearing and migratory life stages (Simenstad et al 2003; Maser and  
34 Sedell 1994; Washington Department of Fish and Wildlife 1997; Collins et al. 2002).

35 The following subsections provide a life history for each of the covered species considered  
36 in this DEIS, beginning with those species having federally listed status. See FPHCP  
37 Chapter 4 (Life History, Status, Distribution and Factors Affecting Covered Species) for  
38 additional descriptions of species characteristics.

39 **Chum Salmon**

40 Within Washington State, two populations of chum salmon are federally listed as  
41 threatened (i.e., Hood Canal summer chum and Columbia River chum). Chum salmon,



## Chapter 3

---

1 also known as dog salmon and/or calico salmon, are distinguished by the reddish purple  
2 vertical markings along the sides of spawning adults. In the Pacific Northwest, freshwater  
3 migration is typically short (less than 50 miles). Chum salmon utilize the low gradient (0  
4 to 8 percent) reaches of the stream, sometimes tidally influenced, for spawning, but the  
5 nearshore and estuarine areas are known to be early rearing areas for many salmonid  
6 species (Groot and Margolis 1991; Wydoski and Whitney 2003). The estuarine area is a  
7 transition zone for acclimation from fresh to marine water and commonly a feeding area  
8 for early stages of many fish. These regions, estuaries and nearshore areas, are also  
9 spawning and rearing areas for many of the baitfish species that are often food sources for  
10 these early and later stages of salmonids. Chum fry typically spend less than 30 days in  
11 freshwater after emergence but remain in the estuary and nearshore environments as  
12 juveniles, feeding primarily on copepods, tunicates, and euphausiids prior to migrating out  
13 to the ocean (Lichatowich 1993). Chum return to freshwater in 3 to 5 years to spawn and  
14 tend to be group spawners with each female accompanied by one or more males. Chum  
15 salmon carcasses, as well as the other salmon species, provide a food source for juvenile  
16 salmonids and numerous wildlife species. The abundance of chum salmon tends to  
17 fluctuate during even/odd cycles, suggesting a possible competitive interaction with pink  
18 salmon, where present in estuary or nearshore habitats (Salo 1991).

19 Chum salmon have three distinct run times: summer, fall, and winter (late fall). Summer  
20 chum begin their upstream migration and spawning during low summer flows in mid-  
21 August through mid-October with fry emergence ranging from the beginning of February  
22 through mid-April, depending on water temperatures (Washington Department of Fish and  
23 Wildlife (WDFW) and Point No Point Treaty Tribes 2000). Fall chum adults enter the  
24 rivers in late October through November, and spawn in November and December. Winter  
25 chum adults migrate upstream from December through January and spawn from January  
26 through February. Fall and winter chum fry emerge from the gravels in March and April  
27 and quickly outmigrate to the estuary for rearing (Smith 1999).

### 28 **Coho Salmon**

29 One population of coho (Lower Columbia River) is a Federal candidate species within  
30 Washington State. Coho, also known as silver salmon, are distinguished by black spots on  
31 the upper part of the caudal fin and a white mouth. Coho begin their upstream migration  
32 between September and December, penetrate deep into the upper watersheds, generally  
33 spawn from October through February, and emerge in early March to late July. Most  
34 juvenile coho remain at least 1 year in freshwater; although recent studies have shown that  
35 some populations spend time in estuaries prior to smoltification. Those that remain in  
36 freshwater typically spend the summer months rearing in pools or other low-velocity  
37 habitats. Many juvenile coho migrate to off channel habitats such as wetlands or side  
38 channels, during winter, a strategy that provides protection from severe winter flows. They  
39 school for a brief period after emerging from gravel, but later disperse and become  
40 aggressive and territorial (Smith 1999).

41 Streams with more structure (logs/rootwads, boulders, undercut-banks) support more coho,  
42 not only because they provide more territories/usable habitat, but they also provide more  
43 food and cover (Scrivener and Andersen 1982). There is a positive correlation between





1 their primary diet of insect material and the extent the stream is overgrown with vegetation  
2 (Chapman 1965). During the winter, coho often feed on adult salmonid carcasses (Bilby et  
3 al. 1996). As coho juveniles grow into yearlings, they become more predatory on other  
4 salmonids. Coho go through physiological changes preparing for life in salt water, and  
5 migrate to sea in spring (Lichatowich 1993). They typically spend 2 years at sea and return  
6 as 3-year-old adults. Coho use estuaries primarily for interim food while they adjust  
7 physiologically to saltwater and then move offshore to deeper waters (Smith 1999).

### 8 **Sockeye Salmon**

9 Within Washington State, two populations of sockeye salmon are federally listed: Snake  
10 River Sockeye are listed as endangered (they spawn in Redfish Lake, Idaho and migrate  
11 within the Columbia River) and Ozette Lake Sockeye are listed as threatened. Sockeye,  
12 also known as red salmon, are distinguished as spawning adults by their red bodies and  
13 green heads as well as lack of spots on the back or caudal fin. Sockeye enter freshwater for  
14 upstream migration during the summer months, spend time resting in deep pools or lakes,  
15 and enter the spawning grounds when ready to spawn. Sockeye are unique in that they  
16 exhibit three life history strategies: one type spawns in rivers but rears in lakes for 1 to  
17 3 years to complete their freshwater life cycle prior to migrating out to sea (lacustrine-  
18 adfluvial); one type spawns along lake shores and rears in lakes for 1 to 3 years prior to  
19 migrating out to sea (lacustrine); and one less common type spawns and rears in rivers and  
20 streams (fluvial).

21 Incubation time varies from 50 days to 5 months, depending on water temperature. After  
22 emerging, lake-rearing fry find their way to a nursery lake, where they feed on insects,  
23 larvae, and copepods. Juvenile sockeye spend up to 3 years in freshwater prior to  
24 smoltification in spring; although some populations outmigrate immediately upon  
25 emergence and others may remain in freshwater for their entire lives (e.g., kokanee).  
26 Migrating sockeye juveniles remain within the estuarine/nearshore environment throughout  
27 the summer, feeding on insects, crustaceans, and small fish and their larvae. Sockeye grow  
28 and develop for 2 to 4 years in the ocean prior to returning to their natal stream to spawn  
29 (Wydoski and Whitney 2003).

### 30 **Chinook Salmon**

31 Within Washington State, Upper Columbia spring chinook salmon are federally listed as  
32 endangered. Chinook in the Snake River, Puget Sound, Lower Columbia River, and upper  
33 Willamette River are listed as threatened. Chinook salmon, also known as king salmon,  
34 are distinguished as adults by black spots on both lobes of the caudal fin and black gums  
35 along the lower jaw. At maturity (4 to 5 years average), chinook seek out spawning  
36 grounds, which can extend from just above tidal influence to as far as 1,200 miles upstream  
37 (Wydoski and Whitney 2003). Spawning habitat preferences include deeper water and  
38 larger gravels than for most other salmon (Healey 1991).

39 Run timing, when adults return to the freshwater to spawn, can occur during spring,  
40 summer, or fall, depending upon particular stock and river system. Spring chinook adults  
41 begin river entry in May or early June and spawn from July through September, typically  
42 in headwater areas with higher gradient habitat. Incubation continues through autumn and



## Chapter 3

---

1 winter, generally requiring additional development time due to the colder water  
2 temperatures of the headwaters. Adult summer chinook begin river entry as early as June  
3 and spawn from September through October. Fall chinook stocks range in spawn timing  
4 from late September through December. Chinook juveniles incubate in the gravel until  
5 January through early March. After emerging from the gravel, juveniles rear in fresh water  
6 from a couple of months to a couple of years. Outmigration of smolts to the marine  
7 environment occurs over a broad period, typically January through August (Smith 1999).  
8 Time is spent within estuarine and nearshore environments prior to entering the ocean.

9 While some emerging chinook fry outmigrate quickly, most inhabit the shallow side  
10 margins and side sloughs for up to two months. At this time, some gradually move into the  
11 faster water areas of the stream and/or disperse into tributaries for rearing, while others  
12 outmigrate to the estuary. Most summer and fall chinook outmigrate within their first year  
13 of life, but a portion of some stocks remain in the river an additional year. Spring chinook  
14 stocks, which rear in colder water and often further upstream, typically rear at least a year  
15 in fresh water (Marshall et al. 1995). However, for Washington populations, Columbia  
16 River Chinook stocks, other than Hanford Reach fall Chinook salmon and a few others, are  
17 primarily river type (with extended freshwater residence), while coastal and Puget Sound  
18 populations exhibit a variable river and ocean life history (with short-term freshwater  
19 rearing).

### 20 **Steelhead**

21 Within Washington State, Upper Columbia Steelhead are federally listed as endangered.  
22 Steelhead within the Snake River and Lower and Middle Columbia River, and Upper  
23 Willamette River steelhead are listed as threatened. Steelhead are distinguished by their  
24 uniform silvery color until darkening toward spawning time. They are the anadromous  
25 form of rainbow trout. Unlike Pacific salmon, steelhead may return to sea after spawning  
26 and migrate again to freshwater to spawn again another year. There are two races of  
27 steelhead: summer and winter. While there is some overlap, winter run steelhead typically  
28 enter streams for spawning between November and April, while stream-maturing summer  
29 steelhead enter streams between May and October (Wydoski and Whitney 2003). Summer  
30 steelhead usually spawn further upstream than winter stocks and dominate inland areas  
31 such as the Columbia Basin. The coastal streams in western Washington typically support  
32 more winter steelhead populations (Smith 1999).

33 Steelhead juveniles typically spend 1 to 2 years (rarely 3 years) in fresh water, preferring  
34 riffle areas in the summer and occupying pools during the rest of the year (Wydoski and  
35 Whitney 2003). During the winter, they often feed on the carcasses of adult salmonids  
36 (Bilby et al. 1996). Steelhead migrate to sea in the spring, spending up to 4 years in the  
37 open ocean (Wydoski and Whitney 2003), feeding on crustaceans, squid, herring, and other  
38 fish (Lichatowich 1993).

### 39 **Bull Trout/Dolly Varden**

40 Bull trout are federally listed as threatened throughout their range. Until 1978, bull trout  
41 and Dolly Varden, both native char, were considered the same species. They were  
42 eventually separated based on morphometrics, osteological features, and embryological



1 development. Bull trout inhabit both eastern and western Washington State, while Dolly  
2 Varden inhabit only Puget Sound and coastal rivers west of the Cascades. Bull trout and  
3 Dolly Varden exhibit four life history strategies: anadromous, adfluvial, fluvial, and  
4 resident.

5 Anadromous forms move upstream in late summer and early fall to spawn in September  
6 and October, or in November at higher elevations (Wydoski and Whitney 2003). All types  
7 of bull trout and Dolly Varden prefer clean, cold water (50°F [10 °C]) for spawning  
8 (Oregon Department of Environmental Quality 1995) and even colder water (36 to 39°F [2  
9 to 4 °C]) for incubation (Rieman and McIntyre 1993). Extended incubation periods (up to  
10 220 days) make bull trout eggs and fry particularly susceptible to increases in fine  
11 sediments (USFWS 1998). Fry are typically found in shallow, backwater side channels  
12 and eddies in proximity to instream cover (Pratt 1984), juveniles in interstitial spaces in the  
13 substrate, and subadults in deeper pools in streams or in the deep water of lakes with  
14 temperatures less than 59°F (15 °C) (Pratt 1992). Bull trout mature at approximately 5  
15 years, typically reproduce in alternate years (Armstrong and Morrow 1980; USFWS 1998)  
16 and live for 12 or more years. Migratory forms of bull trout (anadromous and fluvial) are  
17 known to move between fresh water and marine water and between natal (spawning and  
18 rearing fresh water) and non-natal (habitat outside of their spawning and rearing habitat)  
19 waters, particularly to find forage and overwinter (USFWS 2004b).

20 Resident bull trout and Dolly Varden exhibit three life history strategies, each with unique  
21 habitat requirements: adfluvial, fluvial, and resident. Adfluvial bull trout rear as juveniles  
22 in tributaries, migrate to lakes where most of their growth occurs, then return to the  
23 tributaries as adults to spawn. Fluvial bull trout spawning occurs in smaller tributaries with  
24 major growth and maturation occurring in river mainstems. Resident bull trout complete  
25 all life stages (spawning, rearing, overwintering) in small headwater streams, often  
26 upstream of barriers to other salmonids (USFWS 2004b).

### 27 **Cutthroat Trout**

28 The coastal sea-run cutthroat trout and the westslope cutthroat trout are two subspecies of  
29 cutthroat trout native to Washington State. Both subspecies are Federal species of concern.  
30 Only the coastal sea-run cutthroat trout is anadromous, but it also exhibits fluvial,  
31 adfluvial, and resident life history forms. The westslope cutthroat, which predominately  
32 occurs in major watersheds of eastern Washington, has only fluvial, adfluvial, and resident  
33 life history forms.

34 Coastal cutthroat trout spawn in the cooler waters of the smallest headwater streams and  
35 tributaries used by any salmonid species, and the young usually remain in these streams  
36 about a year before moving down into larger streams. They live in these larger streams for  
37 another 2 to 5 years (usually 3) before migrating to the Pacific Ocean (Wydoski and  
38 Whitney 2003). Some stocks remain as residents of small headwater tributaries, or migrate  
39 only into rivers or lakes (Scott and Crossman 1973). Sea-run cutthroat do not migrate to  
40 the open ocean; rather, they stay in estuarine habitats near the mouths of their natal streams  
41 for 5 to 8 months of the year. Upstream migration to freshwater feeding/spawning areas  
42 occurs from late June through March. Re-entry timing is consistent from year to year



## Chapter 3

---

1 within streams, but varies widely between streams. Spawning generally occurs between  
2 December and May in pool tailouts located in streams with low gradient and low flows or  
3 in shallow riffles (Wydoski and Whitney 1979).

4 Historically within Washington State, westslope cutthroat were distributed in two adjacent  
5 river basins (Lake Chelan and Methow) in the mid-Columbia River drainage and the  
6 headwaters of the Pend Oreille River in northeastern Washington (Williams 1999 as cited  
7 within Wydoski and Whitney 2003). However, over the years, hatchery westslope  
8 cutthroat have been introduced throughout eastern Washington as well as a few streams on  
9 the westside of the Cascades. Genetically pure populations have been identified in various  
10 pristine headwater streams and alpine lakes within eastern Washington; however,  
11 inventories are not complete. Adfluvial westslope cutthroat spend 1 to 4 years as juveniles  
12 in tributary streams before moving into lakes to rear. These adult cutthroat move back into  
13 tributaries during high stream flows and spawn between March and July; however, they  
14 return to the lake habitat shortly after spawning. Fluvial westslope cutthroat move from  
15 tributary streams to mainstem habitat. Overwintering occurs in deeper pools and beaver  
16 ponds. The resident life history of westslope cutthroat is similar to that of resident coastal  
17 cutthroat (Wydoski and Whitney 2003).

### 18 **Interior Redband Trout**

19 Interior redband trout are a Federal species of concern. Native redband trout are the non-  
20 anadromous inland subspecies of rainbow trout. Although redband trout appear to be  
21 widely distributed within the Columbia River Basin, their status is clouded by the  
22 uncertainty over taxonomic classification within the species and by more than a century of  
23 stocking hatchery rainbow trout and steelhead. Little published information exists for  
24 redband trout in Washington State. Oregon status reports have described some life history  
25 traits. In some basins, fluvial and adfluvial redband trout migrate upstream in the spring  
26 and spawn in their respective basins from April to July depending upon elevation. Most  
27 resident fish spawn in the spring and summer (Oregon Department of Fish and Wildlife  
28 1999).

### 29 **3.8.3.2 Green Sturgeon**

30 The green sturgeon is a Federal species of concern. It is known to spawn in only three  
31 river basins outside of Washington State (Klamath, Rogue, and Sacramento), leaving it  
32 vulnerable to a catastrophic event. Freshwater habitats used by sturgeon are generally  
33 large and deep. With the exception of a few landlocked populations, green sturgeon spend  
34 most of their time in marine waters, moving into fresh water only to spawn (Scott and  
35 Crossman 1973; Setter and Brannon 1992). Physical characteristics of green sturgeon eggs  
36 suggest that the species probably requires cold, clean water for spawning and probably do  
37 not spawn in Washington waters (Moyle et al. 1995). Green sturgeon have been reported  
38 as far as 140 miles inland in the Columbia River, but are presently restricted to areas below  
39 Bonneville Dam, and are found almost exclusively in the lower 40 miles of the river  
40 (Moyle et al. 1995; Wydoski and Whitney 2003). The Columbia River estuary and other  
41 coastal Washington estuaries appear to attract concentrations of green sturgeon during late  
42 summer and early fall. Neither feeding nor spawning occurs in association with these



1 concentrations, and there is no information about how much of the population is in these  
2 concentrations each year, or whether this varies (Adams et al. 2002).

### 3 **3.8.3.3 Pacific and River Lamprey**

4 Pacific and river lamprey are both Federal species of concern. Pacific lamprey populations  
5 in the upper Columbia and Snake River basins have declined dramatically, likely as a result  
6 of elevated water temperatures, sedimentation of spawning gravels, and barriers to  
7 migration (Close et al. 1995). Population declines of the Pacific lamprey have prompted  
8 concern for the river lamprey. In Washington, the Pacific lamprey is found in most large  
9 coastal and Puget Sound rivers and occurs long distances inland in the Columbia, Snake,  
10 and Yakima River systems (Lee et al. 1980; Wydoski and Whitney 2003). River lamprey  
11 are found in coastal streams and estuaries and inland in the Columbia River to the  
12 Columbia Gorge (Kostow 2002; Wydoski and Whitney 2003). Both lamprey are  
13 anadromous and parasitic. Pacific lamprey enter freshwater between July and October.  
14 They gradually move upstream to spawn the following spring (Hart 1973). Eggs hatch in 2  
15 to 4 weeks (19 days at 59°F [15 °C]); newly hatched ammocoetes (larvae) remain in their  
16 nests for 2 to 3 weeks before drifting downstream and burying themselves in mud at the  
17 bottom of pools, or other areas of soft mud and sand (Hart 1973; Moyle 1976). Increased  
18 water flows during runoff can also encourage outmigration by washing away the sand and  
19 silt that the larvae require for anchoring themselves to the bottom (Hardisty and Potter  
20 1971).

21 Little is known regarding the habitat requirements of the river lamprey. Adults migrate  
22 into deep freshwater habitats in the fall. They spawn in the winter and spring in clean  
23 gravel areas of small tributaries and die after spawning (Moyle et al. 1995). Based on  
24 comparisons with other lamprey species, Hart (1973) surmised that river lamprey  
25 ammocoetes (larvae) remain in their natal streams for several years, usually in silt-sand  
26 backwaters and eddies near the bank.

### 27 **3.8.3.4 Margined Sculpin**

28 Margined sculpin have no Federal listing, but are a Washington State sensitive species.  
29 They are found in the Walla Walla, Touchet, and Tucannon Rivers. Margined sculpin are  
30 predominantly found in pools and glides but are also observed in riffles. They prefer small  
31 gravels and silts, avoiding larger gravels and cobble. They reach an average length of  
32 about 2.5 inches and probably live to be 4 years old. Spawning occurs in spring under  
33 rocks, rootwads, or logs (Wydoski and Whitney 2003).

### 34 **3.8.3.5 Pygmy Whitefish**

35 Pygmy whitefish have no Federal listing, but are a Washington State sensitive species.  
36 Pygmy whitefish are members of the trout and salmon family (Salmonidae) and are  
37 typically around 5 to 6 inches in length when mature, reaching a maximum length of about  
38 11 inches. The pygmy whitefish is a remnant species from the last ice age with a spotty  
39 distribution across northern North America and in the Columbia River drainage in  
40 Washington. The pygmy whitefish inhabits cold lakes and streams. Streams inhabited by  
41 pygmy whitefish may be of moderate to swift current and may be silty or clear (Hallock  
42 and Mongillo 1998). The pygmy whitefish has been eliminated from a minimum of



## Chapter 3

---

1 40 percent of its range in Washington. Historically, pygmy whitefish were known to have  
2 occupied 15 lakes in Washington; however, today they are currently found in only 9  
3 (Hallock and Mongillo 1998). The future of pygmy whitefish populations in Washington  
4 is dependent upon our ability to maintain water quality, spawning habitat, and prevent  
5 introduction of new predator species. Additionally, pygmy whitefish populations are  
6 especially vulnerable to local extinction because natural reintroduction of new fish is  
7 usually impossible among isolated lake systems.

### 8 **3.8.3.6 Olympic Mudminnow**

9 Olympic mudminnows have no Federal listing, but are a Washington State sensitive  
10 species. They are found only in slow-moving streams, wetlands, and ponds with soft mud-  
11 bottom substrate, little or no water flow, and abundant aquatic vegetation (Harris 1974;  
12 Mongillo and Hallock 1999; Wydoski and Whitney 2003). Species distribution is limited  
13 to low gradient, low elevations in the coastal lowlands of the Olympic Peninsula, the rivers  
14 of the Chehalis and lower Deschutes drainages, and south Puget Sound lowlands west of  
15 the Nisqually River (Mongillo and Hallock 1999; Wydoski and Whitney 2003). The  
16 species is considered vulnerable due to its limited distribution and dependence on healthy  
17 wetland habitat (Mongillo and Hallock 1999). Wydoski and Whitney (2003) observe that  
18 mudminnows are usually found under overhanging banks or shore vegetation, preferring  
19 areas with low light and brownish water of bogs and swamps. Meldrim (1968) found a  
20 wide tolerance for temperature extremes and low oxygen levels, but a restricted tolerance  
21 range for salinity and water current. Adults spawn between November and June (peaking  
22 in April and May) and females deposit eggs amidst clumps of vegetation, to which fry  
23 remain firmly attached for approximately one week after hatching (Meldrim 1968 and  
24 Hagen et al. 1972 in Mongillo and Hallock 1999).

### 25 **3.8.3.7 Umatilla Dace and Leopard Dace**

26 Neither the Umatilla nor leopard dace have Federal listing status; however, they are both  
27 classified as Washington State candidate species. Concern for these species is prompted  
28 by their restricted distribution as well as their preference for shallow water, which may  
29 increase their vulnerability to activities that affect water levels (Royal British Columbia  
30 Museum 1995). Two other species of dace (Longnose and speckled dace) also occur in  
31 Washington State and share many key life history characteristics, such as breeding habitat,  
32 spawning period, and habitat associations of juveniles. Dace typically occur in shallow  
33 waters with cobble or gravel substrate (Scott and Crossman 1973; Troffe 1999; Wydoski  
34 and Whitney 2003). Spawning generally occurs from late spring through summer  
35 (Wydoski and Whitney 2003). Currently, relatively little information is available about the  
36 Umatilla dace. At one time, this species was considered to be a stable hybrid between  
37 leopard dace and speckled dace and, to date, is not a “recognized species in the United  
38 States by the American Fishery Society. The taxonomic status is still open to debate”  
39 (Personal Communication, Molly Hallock, Washington Department of Fish and Wildlife,  
40 October 2003). The Umatilla dace is endemic to the Columbia River basin (Troffe 1999).  
41 In Washington, it is known only from the larger main rivers of the basin, including the  
42 Columbia River at Rock Island Dam and elsewhere in the Columbia River east of the  
43 Cascades (Royal British Columbia Museum 1995).



1 The leopard dace is a Columbia River system fish, found west of the Cascade Mountains in  
2 the Lower Columbia and the Cowlitz River system (Personal Communication, Molly  
3 Hallock, Washington Department of Fish and Wildlife, October 2003) as well as in the  
4 Columbia River system east of the Cascade Mountains (Scott and Crossman 1973;  
5 Wydoski and Whitney 2003).

### 6 **3.8.3.8 Lake Chub**

7 Lake chub have no Federal status, but are a Washington State candidate species. In  
8 Washington, they are known only from the northeastern part of the State (Wydoski and  
9 Whitney 2003). Lake chub exhibit a variety of habitat preferences across their range,  
10 living in large rivers at northern latitudes, but using lake habitat when it is available (Isaak  
11 et al. 2003; Scott and Crossman 1973). In the southern portion of their range, the lake  
12 chub is uncommon in lakes; but whether this represents a habitat preference or is simply  
13 because fewer lakes are available is unknown (Isaak et al. 2003). The only known  
14 Washington populations are in Cedar Lake in Stevens County (1998 observation) and in  
15 North Fork Beaver Creek in Okanogan County (1999 observation; Personal  
16 Communication, Molly Hallock, Washington Department of Fish and Wildlife, October  
17 2003). Throughout their range, lake chub are found in clear, cool water with clean cobble  
18 or gravel substrates (Isaak et al. 2003). Spawning lake chub move into shallow areas along  
19 the margins of streams or lake shores during the spring in water between 55 and 65 °F  
20 (12.8 and 18.3 °C).

### 21 **3.8.3.9 Eulachon**

22 Eulachon have no Federal status, but are classified as Washington State candidate species.  
23 Eulachon, also known as candlefish, are anadromous with large spawning runs in the  
24 Columbia, Cowlitz, Kalama, Lewis, Sandy, and Nooksack Rivers during late winter. They  
25 spend most of their lives in the nearshore waters of the Pacific Ocean, migrating only a  
26 short distance upstream to spawn. They spawn at night in gravel with males preceding the  
27 females in the migration. Larvae emerge in about 1 month and generally move  
28 immediately out to sea (Wydoski and Whitney 2003).

### 29 **3.8.3.10 Mountain Sucker and Salish Sucker**

30 Mountain suckers are classified as Washington State candidate species, and Salish suckers  
31 are classified as Washington State monitor species. Neither species have a Federal listing  
32 status. In British Columbia, the Salish sucker's restricted distribution, along with the  
33 effects of habitat degradation and loss, have led to considerable concern that the species  
34 may face extirpation in that part of its range. Populations in Washington appear to be more  
35 stable, and face fewer threats (British Columbia Ministry of Environmental, Lands, and  
36 Parks 1993). The mountain sucker is a State candidate species due to its sensitivity to high  
37 stream temperatures, sedimentation of spawning habitat, and/or lack of preferred food  
38 items (Washington Department of Fisheries 1991). Suckers comprise a family of bottom-  
39 dwelling fish in lakes and streams. The Salish sucker has been documented at Lake  
40 Cushman on the Olympic Peninsula, and in the river systems along east/northeast Puget  
41 Sound (Personal Communication, Molly Hallock, Washington Department of Fish and  
42 Wildlife, October 2003). The mountain sucker occurs in the Columbia River and its



## Chapter 3

---

1 tributaries east of the Cascades (Wydoski and Whitney 2003), as well as in the Toutle and  
2 Cowlitz River systems west of the Cascades (Personal Communication, Molly Hallock,  
3 Washington Department of Fish and Wildlife, October 2003). Suckers spawn over gravel  
4 substrates in riffles during spring and early summer. Mountain suckers appear to prefer  
5 riffles at the downstream ends of pools. Fingerling mountain suckers have been found in  
6 small intermittent streams with little discharge and abundant vegetation, and large young  
7 have been associated with weedy side channels or deep pools (Scott and Crossman 1973;  
8 Wydoski and Whitney 2003).

### 9 **3.8.3.11 Sandroller**

10 The sandroller has no Federal listed status. However, the sandroller is classified as a  
11 Washington State monitor species, presumably due to its restricted range and the paucity of  
12 information about its life history needs. The sandroller is a small fish, generally less than 5  
13 inches long, and is one of only two species in the trout-perch family in North America.  
14 These fish appear to be very secretive in behavior, remaining well spaced from other  
15 individuals and are rarely collected in abundance. In Washington, the sandroller is found  
16 only in the Columbia River system and some of its tributaries, including the Yakima and  
17 Cowlitz Rivers, and up the Snake River into Idaho (Wydoski and Whitney 2003). This  
18 species has been found in quiet backwaters with cover such as undercut banks, submerged  
19 tree roots, and debris in small streams, but is also found as deep as 71 feet in the Columbia  
20 River (Wydoski and Whitney 2003). Habitat associations appear to vary with age; young-  
21 of-the-year occur primarily in weed bays or waterways adjacent to the main river, while  
22 adults may be associated with eddies behind large boulders, logs, and bridge supports  
23 (Katula 1992). Page and Burr (1991) describe the sandroller as usually found in sand and  
24 near vegetation. Spawning may occur on rocky substrates in shallow streams or along the  
25 shallow shores of rivers (Katula 1992).

### 26 **3.8.4 The Freshwater Aquatic Ecosystem**

27 Natural channels are complex and contain a mixture of habitats differing in depth, velocity,  
28 and cover (Bisson et al. 1987). They are often formed during storm events with associated  
29 flows that mobilize sediment in the channel bed (Murphy 1995). The hydrologic regime of  
30 a watershed, combined with its geology, hillslope characteristics, and riparian vegetation  
31 determines the nature of stream channel morphology (e.g., number and spacing of pools  
32 and width-to-depth ratio) (Beschta et al. 1995; Sullivan et al. 1987). Therefore, activities  
33 in these areas would be expected to affect the shape and form of the stream channel. For  
34 example, substantial increases in volume and frequency of peak flows can cause streambed  
35 scour and bank erosion. A large sediment supply may cause aggradation (i.e., filling and  
36 raising the streambed level by sediment deposition) and widening of the stream channel,  
37 pool filling, and a reduction in gravel quality (Madej 1982). Upslope activities (e.g.,  
38 timber harvest, land clearing, and road development) can change channel morphology by  
39 altering the amount of sediment or water contributed to the streams. This, in turn, can  
40 disrupt the balance of sediment input and removal in a stream (Sullivan et al. 1987).

41 Stream habitat conditions in Washington are affected by a wide range of factors, including  
42 geophysical changes (e.g., volcanic eruptions, earthquakes, and associated uplifting),





1 extremes of flow (e.g., flooding and low flow), existing geological conditions (e.g.,  
2 erodible soils), and land-use practices (e.g., timber harvest, grazing, urban development,  
3 road construction and operation, and gravel mining). The effects of these combined factors  
4 result in the existing stream habitat conditions.

5 Streams that lack a balance between pools and riffles are often less productive than streams  
6 that have more complex structure. Pools are used as holding and resting areas for adult  
7 fish prior to spawning, deep water cover for protection, and cool water refugia during low  
8 flow summer months. Riffles are important for re-oxygenation of water, habitat for food  
9 organisms such as aquatic macroinvertebrates, and as rearing areas for fish (Gregory and  
10 Bisson 1997). Intensive timber harvest has been reported to decrease pool depth, surface  
11 area, and the general diversity of pool character (Ralph et al. 1994). Possible mechanisms  
12 include decreased occurrence of LWD (which can help form and stabilize pools) and filling  
13 of remaining pools with bed material.

14 Attempts that have been made to define a range of optimum pool-to-riffle ratios for a  
15 properly functioning system have been described in the literature (NMFS 1996b; USFWS  
16 1998). However, applying these values to field conditions would require considering site-  
17 specific characteristics such as existing LWD, stream gradient, bank characteristics,  
18 sediment load, bed material (e.g., bedrock and boulders), and other watershed factors such  
19 as hydrologic conditions (Murphy 1995).

20 The following describes components of the freshwater aquatic ecosystem that are  
21 influenced by forest practices. These include coarse sediment, fine sediment, hydrology,  
22 LWD, leaf/needle litter recruitment, floodplains and off-channel features, water  
23 temperature, dissolved oxygen, forest chemicals (contaminants), and fish passage.

### 24 **3.8.4.1 Coarse Sediment**

25 A certain amount of bedload material is necessary to provide substrate for cover and  
26 spawning habitat for fish. For example, anadromous salmon typically use gravels ranging  
27 from 0.5 to 4 inches (12.7 to 101.6 mm), whereas steelhead and resident trout may use  
28 smaller substrates ranging from 0.25 to 4 inches (6.4 to 101.6 mm) (Bjornn and Reiser  
29 1991). Increased levels of coarse sediment bedload above background levels can,  
30 however, lead to streambank instability, pool filling, and changes in the water transport  
31 capacity of the channel (Spence et al. 1996). The larger the sediment size, the higher the  
32 flow that is required to mobilize sediment. Consequently, the recovery periods for streams  
33 with severe coarse sediment aggradation could range from decades to 100 years or more.  
34 The major factors influencing the excessive delivery of sediment to a stream include the  
35 intensity and location of streambank erosion, mass-wasting events, and road and culvert  
36 failures.

### 37 **3.8.4.2 Fine Sediment**

38 Fine sediment, including sand- and silt-sized particles can reduce stream habitat quality,  
39 restrict sunlight penetration, and fill pores between the gravel, thus preventing the flow of  
40 oxygen-rich water to fish eggs that may be deposited in the gravel. In laboratory studies, a  
41 substrate containing 20 percent fines was found to reduce emergence success of young



## Chapter 3

---

1 salmon and trout by 30 to 40 percent (Phillips et al. 1975; MacDonald et al. 1991).  
2 According to study results and summaries from Peterson et al. (1992) and Chapman  
3 (1988), survival of salmonid eggs is highest in substrates that contain less than 11 to  
4 16 percent particles within the fine sediment category.

5 Fine sediments may clog pores or breathing surfaces of aquatic insects, physically smother  
6 them, or decrease available habitat (Spence et al. 1996; Nuttall and Bilby 1973; Bjornm et  
7 al. 1974; Cederholm et al. 1978; Rand and Petrocelli 1985). In general, the larger the  
8 gravel size and more complex the substrate, the more diverse the invertebrate fauna (Hynes  
9 1970). Important factors influencing the excessive delivery of fine sediment to a stream  
10 include the presence of adequate streamside vegetation to filter fine sediment derived from  
11 hillslopes and road surface erosion (subsection 3.4, Geology, Soils, and Erosion;  
12 subsection 3.5, Water Resources; and subsection 3.7, Riparian and Wetland Processes).  
13 Also, fine sediment is usually present with coarse sediment delivery processes described  
14 above.

15 Biological effects of increased turbidity due to suspended sediments may include a  
16 decrease in primary productivity of algae and periphyton due to the decrease in light  
17 penetration. Declines in primary productivity can adversely affect the productivity of  
18 higher trophic levels such as macroinvertebrates and fish (Gregory et al. 1987). Suspended  
19 sediments can also interfere with feeding behavior or cause gill damage in fish (Hicks et al.  
20 1991). While most effects of elevated suspended sediment (or turbidity) are negative, in  
21 some cases turbidity enhances juvenile fish cover from predatory fish (Gregory and  
22 Levings 1998).

### 23 **3.8.4.3 Hydrology**

24 The amount of water provided to aquatic ecosystems at critical times is important for  
25 sustaining fish and other aquatic species. Fish adapt to natural flow cycles for feeding,  
26 spawning, migration, and survival needs. The timing, magnitude, and duration of peak and  
27 low flows must be sufficient to create and maintain riparian and aquatic habitat. Flows can  
28 be influenced by management activities such as timber harvest and roads (subsection 3.5.2,  
29 Surface Water Quantity). In general, low- or base-level stream flows that occur during the  
30 late summer often limit habitat and survival for rearing juvenile salmon and trout. They  
31 can also negatively affect migration and access to habitat and food resources, as well as  
32 disrupting spawning behavior. Such conditions can occur naturally during this period due  
33 to lack of precipitation. However, low flows can be exacerbated by water withdrawals,  
34 silting (which can decrease pool depth), and stream widening resulting from unstable  
35 banks.

36 High winter flows and floods that scour the streambed can be detrimental to eggs or young  
37 fish that may be incubating in the stream gravels (Thorne and Ames 1987). Both extreme  
38 high and low flow conditions may occur in different regions of the State. Rain-on-snow  
39 events are a common reason for flooding and streambed scour on the west of the Cascade  
40 Mountains (Harr 1986; Washington Department of Fish and Wildlife 2000;  
41 <http://wdfw.wa.gov/fish/chum/chum-10.htm>). In contrast, the eastside of the State lies in  
42 the rain shadow of the Cascade Mountains (subsection 3.5.2, Surface Water Quantity).



1 Consequently, extreme low flows and high water temperatures can be detrimental during  
2 the summertime (Talayco 2002; Northwest Power Planning Council 1986; Haring 2001).

### 3 **3.8.4.4 Large Woody Debris**

4 LWD is one of the most important components of high quality fish habitat (Marcus et al.  
5 1990). LWD is the primary channel-forming element in some channel types and affects  
6 many aspects of channel morphology including stream roughness, sediment storage, water  
7 retention, energy dissipation, and fish habitat (Marcus et al. 1990; Lisle 1986; Swanson et  
8 al. 1987; Martin et al. 1998). Pools formed by stable accumulations of LWD provide  
9 important habitat for rearing salmonids, particularly in winter (Heifetz et al. 1986; Murphy  
10 et al. 1986). LWD has also been shown to affect macroinvertebrate populations (Naiman  
11 and Bilby 1998). Macroinvertebrates readily colonize LWD using it as a stable substrate,  
12 and in some cases, as a food resource (Anderson and Sedell 1979). The value of LWD in  
13 providing aquatic habitat depends on stream size, tree species, and numerous other factors  
14 (subsection 3.7, Riparian and Wetland Processes). Field studies in old-growth, Douglas-fir  
15 forest streams in coastal Oregon and Washington have shown that the number of woody  
16 debris pieces varies by channel width and size of debris under undisturbed conditions. For  
17 example, studies by Bilby and Ward (1989) and the Forest Practices Board (1995) show  
18 that the number of LWD pieces decreased with increasing width of a stream; however, the  
19 average diameter, length, and volume of LWD increased. The type of wood is an  
20 important factor (subsection 3.7.1.1, Riparian Functions). For example, coniferous wood  
21 (e.g., Douglas-fir or cedar) is more resistant to decay than deciduous wood (e.g., alder).  
22 Therefore, coniferous wood has a greater longevity in a stream (Cummins et al. 1994 as  
23 quoted in Spence et al. 1996).

24 Past forest management practices often included splash dams and stream cleaning efforts  
25 (Maser and Sedell 1994). During the last century, splash dams were built to aid in floating  
26 and transporting harvested trees to the mill. From the 1950s through the 1970s, removal of  
27 LWD from streams was based on the belief that it was detrimental to salmon migration.  
28 Much of the LWD removed from streams was logging slash and debris; however, naturally  
29 occurring LWD was also removed (Personal Communication, Steve Keller, NMFS  
30 [formerly], November 17, 2004). Both practices, splash dams and stream cleaning,  
31 contributed to major changes in the amount of cover habitat available and often changed  
32 stream habitats to a single, cobble-bed channel lacking pools and LWD, or to bedrock  
33 channels lacking gravel, woody debris, and other channel features (Murphy 1995; Maser  
34 and Sedell 1994). This decrease in LWD corresponded to a reduction in salmonid use  
35 (House and Boehne 1987). Due to the time required for streamside trees to grow and  
36 mature to potential LWD, there may be a considerable lag period (e.g., greater than about  
37 50 years and up to 300 years) before additional LWD is contributed to a cleared stream  
38 (Gregory and Bisson 1997).

39 In general, information on LWD must be viewed from the perspective of the timber harvest  
40 activity in the area, historic floods that have removed or redistributed LWD, and the  
41 activities that were performed to actively remove LWD (subsection 3.7.1.1, Riparian  
42 Functions). Potential LWD recruitment from existing mature or old growth riparian zones  
43 would be anticipated to be higher than younger or recently clearcut areas (subsection



## Chapter 3

---

1 3.7.1.1, Riparian Functions). There may be no potential for LWD recruitment in currently  
2 open areas such as prairies and grasslands, which may not develop into forested areas in  
3 the foreseeable future.

4 LWD enhancement has recently become a more common method for improving stream  
5 reaches lacking wood. The methods for placing LWD are fairly advanced (Oregon  
6 Department of Forestry and Oregon Department of Fish and Wildlife 1995). LWD  
7 placement would provide short-term benefits to stream systems providing a more complex  
8 habitat structure, nutrient input, and substrate for invertebrate colonization, all of which  
9 would benefit fish habitat. These benefits may improve current conditions in many areas  
10 until the natural riparian corridor can regenerate and provide consistent inputs of LWD.

### 11 **3.8.4.5 The Aquatic Food Chain**

12 The base of the aquatic food chain is supported by the combination of dissolved chemical  
13 nutrients and detrital materials. The chemical constituents such as nitrogen (usually in the  
14 form of nitrates and nitrites), phosphorus, and carbon can be derived from the breakdown  
15 of detritus and through leaching and runoff from surrounding soils (Gregory et al. 1987).  
16 Many bacterial and macroinvertebrate species rely directly on detrital material from leaf  
17 and needle litter, branches, and stems from the surrounding riparian zone vegetation.  
18 Some estimates indicate that leaf and needle recruitment may provide up to 60 percent of  
19 the total energy input to stream communities (Richardson 1992). Other macroinvertebrate  
20 species rely on aquatic algae that primarily use dissolved chemical nutrients and require  
21 solar radiation. In streams containing spawning habitat for Pacific salmon, substantial  
22 influxes of nutrients from the marine environment occur during the decomposition of  
23 carcasses (Bilby et al. 1996).

24 The abundance and diversity of macroinvertebrate food sources to salmonids is dependent  
25 upon the primary algae and detrital food sources. Forest harvest activities affect the food  
26 chain by changing the relative macroinvertebrate production between herbivores and  
27 detritivores (Gregory et al. 1987). The magnitude and duration of the change is dependent  
28 upon a variety of factors, including stream size, gradient, location (headwater versus  
29 mainstem), and the type of riparian vegetation and management prescriptions. Gregory et  
30 al. (1987) suggest that tree harvest in riparian areas initially leads to higher total  
31 invertebrate abundance, but fewer invertebrate species, and that recovery of the  
32 macroinvertebrate community occurs over periods similar to recovery of riparian zones.

33 Bilby and Bisson (1992) observed higher summer production of coho fry in a stream  
34 flowing through a clearcut area relative to a nearby stream reach in an old-growth riparian  
35 stand. However, no differences in coho production were present during fall censuses, and  
36 the high summer fish production was attributed to high algae production (Bilby and Bisson  
37 1992). Bilby and Bisson (1992) and Spence et al. (1996) have noted that other changes in  
38 habitat features (e.g., numbers of pools) required by yearling and adult fish could likely  
39 offset any increases in sub-yearling production. Gregory et al. (1987) argued that short-  
40 term higher fish productivity might occur downstream of timber harvest units in some  
41 areas, but at the expense of long-term stability in the overall abundance and diversity of the  
42 aquatic community.



### 1 **3.8.4.6 Floodplains, Off-Channel Habitats, and Hyporheic Zones**

2 Floodplains and off-channel areas are an important component of aquatic habitat that  
3 includes side channels, backwater alcoves, ponds, and wetlands. They provide important  
4 habitat seasonally to particular life stages as well as input of organic matter and LWD.  
5 Seasonally flooded channels and ponds are particularly important for rearing coho salmon  
6 and other fish species during winter months. Large floodplains can also function as filters  
7 for subsurface flows and maintenance of water quality (Gregory and Bisson 1997). Some  
8 backwater alcoves and ponds result from groundwater seeps and may have shade levels  
9 higher than the main channel. These areas provide cool water refugia during high  
10 summertime temperatures. Major floodplains in the planning area generally are located in  
11 the lowest reaches of major rivers. Beavers can play a substantial role in the development  
12 of ponds and wetlands important as habitat for salmon and trout, particularly for juvenile  
13 coho salmon (Cederholm et al. 2001; Pollock et al. 2001).

14 Hyporheic zones (the saturated sediment region under and along streams) are often the  
15 connections between groundwater and surface water in these habitat areas and often supply  
16 substantial habitat for hyporheic organisms such as insects, bacteria, and fungi (Edwards  
17 1998; Naiman et al. 2000). The presence of a hyporheic zone is most often associated with  
18 the alluvium below the stream and in the floodplain adjacent to streams. Nutrients and  
19 organic matter is processed in this zone by bacteria and other organisms (e.g.,  
20 invertebrates, specialize insects, and crustaceans) (Boulton et al. 1998). Where this water  
21 surfaces it may be high in nutrients producing locally high primary production areas  
22 (Edwards 1998; Boulton et al. 1998). The overall exchange of organisms and effects on  
23 stream production is not well known, and in most systems may be relatively minor;  
24 however, it could be more important in some floodplain habitats (Boulton et al. 1998). In  
25 dry summer months or during floods this zone may provide a refuge for some aquatic  
26 organisms (Boulton et al. 1998). Substrate porosity may affect its function and size, but  
27 the relationship is not clear (Boulton et al. 1998).

### 28 **3.8.4.7 Water Temperature**

29 Water temperature plays an integral role in the biological productivity of streams. Water  
30 temperature fluctuations and their relationship to dissolved oxygen can affect all aspects of  
31 salmon and trout life histories in fresh water including:

- 32 • incubation and egg survival in stream gravel;
- 33 • emergence, feeding, and growth of fry and juvenile fish;
- 34 • outmigration of young fish;
- 35 • adult migration, holding and resting; and
- 36 • prespawning and spawning activities.

37 A rise in temperature increases the metabolic rate of aquatic species. Consequently, more  
38 energy is required, even during periods of low activity. In addition, dissolved oxygen  
39 decreases as water temperature increases, potentially increasing stress on fish. Water  
40 temperatures in the range of 70°F (21°C) or greater can cause death in cold-water species  
41 such as salmon and trout within hours or days (Oregon Department of Environmental



## Chapter 3

---

1 Quality 1995). In general, water temperatures of 54 to 59°F (12 to 15°C) have been found  
2 to provide a properly functioning condition for juvenile salmon and trout. However, bull  
3 trout require much lower temperatures during spawning 39 to 50°F (4 to 10°C) and egg  
4 incubation 34 to 43°F (1 to 6°C) (Oregon Department of Environmental Quality 1995).

5 Increases in water temperature in forest streams can often be traced to reduction of shade-  
6 producing riparian vegetation along fish-bearing and tributary streams that supply water to  
7 other fish-bearing streams. However, streams also naturally tend to become warmer as  
8 water flows from headwaters to the sea (Sullivan et al. 1990; Zwieniecki and Newton  
9 1999). This warming occurs as water equilibrates to local environmental conditions  
10 including air temperature, which in turn is highly correlated with elevation. In addition,  
11 water temperatures can be affected by stream widening, sedimentation/stream depth,  
12 microclimate, groundwater, and other upstream inputs (subsection 3.5.1, Surface Water  
13 Quality). Long-term sublethal temperature effects as well as short-term acute effects of  
14 warm water temperatures can be detrimental to the overall health of aquatic species. Heat  
15 stress may accumulate such that increased exposure for juvenile fish increases their  
16 susceptibility to disease (Oregon Department of Environmental Quality 1995).

17 Shade does not always maximize aquatic productivity. The availability of instream algae  
18 can be a limiting factor in some streams. Algae and other sources of organic matter are at  
19 the lowest level of the food chain and important to higher trophic level production such as  
20 fish. Along with light, nutrients (e.g., nitrogen and phosphorus) are key factors that result  
21 in algae production. High levels of shade can result in low levels of algae production even  
22 if adequate nutrient sources are present (Gregory et al. 1984). Under unmanaged  
23 conditions, forested lands generally have low light and low primary productivity in low  
24 order streams with high canopy cover. In contrast, primary productivity in wide, high  
25 order streams is generally unaffected by riparian management because adequate light  
26 penetration occurs even under mature riparian conditions (Gregory et al. 1984).

### 27 **3.8.4.8 Dissolved Oxygen**

28 Adequate dissolved oxygen levels are important for supporting fish, invertebrates, and  
29 other aquatic life. Salmonids are particularly sensitive to reduced dissolved oxygen  
30 (Oregon Department of Environmental Quality 1995). Intergravel dissolved oxygen has  
31 been recognized as crucial to the survival of salmonid embryos. Intergravel dissolved  
32 oxygen depends on several interrelated factors such as water temperature, surface-water  
33 concentrations, percentage of fine sediment and gravel in pores, and the oxygen demand of  
34 the eggs. Management-induced depletion of dissolved oxygen in stream water can occur  
35 from harvest activities, such as excessive amounts of logging debris left in a stream that  
36 can result in decreased dissolved oxygen (MacDonald et al. 1991). Critical levels of  
37 dissolved oxygen also depend on the velocity of the water passing over the eggs, as oxygen  
38 consumption would rapidly reduce oxygen supply to the egg without replenishment  
39 through adequate intergravel flow (velocity). Therefore, at lower velocities, higher initial  
40 oxygen concentrations are needed for proper egg development (Oregon Department of  
41 Environmental Quality 1995). Forest management activities can exacerbate any



1 intergravel dissolved oxygen problems through increases in fine sediment, which reduce  
2 intergravel water velocity (Bjornn and Reiser 1991; Ringler and Hall 1975; Moring 1975).

### 3 **3.8.4.9 Forest Chemicals**

4 Water quality contaminants (e.g., petroleum products, chemicals, sewage, heavy metals)  
5 can severely impair aquatic ecosystems either by sublethal (e.g., reduced growth) or lethal  
6 effects (e.g., fish kills). The water quality contaminants considered here are pesticides  
7 used to prevent tree diseases and to deter other plant species that compete with harvestable  
8 trees for nutrients, space, and light (subsection 3.5.1, Surface Water Quality) (Washington  
9 Forest Practices Board 2001a, Appendix J).

### 10 **3.8.4.10 Fish Passage**

11 Upstream migration of adult salmon, steelhead, and trout to spawning areas or  
12 redistribution of rearing fish to potential habitat in upstream areas can be impeded or  
13 blocked by a number of different mechanisms. These mechanisms can include the  
14 following:

- 15 • Water Temperature—Elevated water temperatures (e.g., greater than 60°F and 68°F  
16 [15.6°C and 20°C] for coho salmon and fall chinook salmon, respectively) are known  
17 to stop the migration of fish (Bjornn and Reiser 1979).
- 18 • Dissolved Oxygen—At least 5 milligrams per liter (mg/l) of dissolved oxygen is  
19 recommended to provide oxygen needs for migrating fish (Bjornn and Reiser 1979).  
20 Decreased oxygen can occur as a result of high water temperatures and oxygen  
21 consumption created by decay of organic debris, chemicals, and respiration.
- 22 • Turbidity—High levels of sediment (e.g., 4,000 mg/l) have been reported (Bjornn and  
23 Reiser 1979) as ceasing upstream migration.
- 24 • Physical Barriers—High waterfalls or cascades that are beyond the jumping or physical  
25 capabilities of fish can prevent upstream migration. Similarly, excessive water  
26 velocities that result in conditions beyond the physical capabilities of a given fish  
27 species can also restrict or prevent upstream migration. The maximum velocity  
28 beyond which coho and chinook salmon cannot successfully move upstream is about 8  
29 feet per second (2.44 meters per second) (Bjornn and Reiser 1979). Shallow water  
30 depths from conditions such as low flow can impede or prevent passage (e.g., chinook  
31 and coho salmon are generally not successful at upstream migration at water depths  
32 less than about 0.8 feet (0.24 meters) or 0.6 feet (0.18 meters), respectively (Bjornn  
33 and Reiser 1979). Such conditions can occur during low flow periods where riffles  
34 between pools can become completely dry or lack sufficient depths for fish passage.
- 35 • Man-made Barriers—Man-made barriers include features such as dams and stream  
36 crossings (usually culverts, but sometimes bridges as well).
- 37 • Streambed Aggradation and Subsurface Flow—Debris flow deposits in fish-bearing  
38 streams have been found to cause fish blockages (Pearce and Watson 1983; Bryant  
39 1983). High stream bedload accumulations have been found to result in subsurface  
40 flow and isolating stream reaches either inhibiting or delaying passage (Furniss et al.  
41 1991; Hartmen et al. 1995).



## Chapter 3

---

1 Stream crossings by forest roads are the most common passage barrier influenced by  
2 Washington Forest Practices Rules. A Hydraulic Project Approval is needed for the  
3 construction of stream crossings, which are regulated by the WDFW under the Hydraulic  
4 Code (WAC 220-110-070). Barriers such as culverts used at stream crossings can prevent  
5 passage due to high water velocities, restricted depths, excessive elevation for successful  
6 entry, size and length of the culvert, and other factors. Similarly, debris jams can prevent  
7 or delay upstream passage (Bjornn and Reiser 1979).

### 8 **3.8.4.11 Lakes, Reservoirs, and Nearshore Marine Habitats**

9 Lakes and reservoirs provide areas for spawning, early rearing, and growth for many fish  
10 species including salmonids. They also function as migratory pathways, especially for  
11 adult and juvenile salmon in many lake and reservoirs systems in the State. The nearshore  
12 environment of lakes and reservoirs depends on riparian areas in many ways that are  
13 similar to streams, relying at least partly on input of LWD from shoreline areas (Harmon et  
14 al. 1986; Christensen et al. 1996) and on terrestrial nutrient input and leaf and litter input as  
15 an aquatic food base (Wetzel 1975). However, much of the production is based on  
16 autochthonous growth (e.g., algae) within the system, typically much more so than most  
17 flowing water systems (Wetzel 1975). Nearshore areas are often a sink for nutrients and  
18 organic matter entering from streams and rivers that enter these systems (Washington  
19 Department of Fish and Wildlife 1997). Temperature and flow conditions in lakes and  
20 reservoirs influence suitability of these environments for various fish species (Washington  
21 Department of Fish and Wildlife 1997). Habitat structure along shorelines influence  
22 species use, including species that may be predatory to juvenile salmonids (Wedge and  
23 Anderson 1979; Savino and Stein 1989).

24 Nearshore marine areas have many similarities to that of lakes and reservoirs as they  
25 provide important rearing and migratory areas for many salmonids. This habitat is also  
26 important for many other marine species includes some of importance as prey for  
27 salmonids (Washington Department of Fish and Wildlife 1997). The special nearshore  
28 marine habitats, estuaries, are often considered very important for development and growth  
29 of many salmonid species during outmigration (Simenstad et al. 1982, Groot and Margolis  
30 1991). Some juvenile salmonids may spend more than a month rearing in estuaries prior to  
31 migrating to the open ocean (Washington Department of Fish and Wildlife 1997). Like  
32 lakes and reservoirs, estuaries are often considered a sink for a variety of upland and  
33 riverine processes. River flow and transported nutrients, sediment, and other organic  
34 matter including LWD are important to estuaries. LWD provides structure and nutrients  
35 for marine habitats (Maser and Sedell 1994), although its relative importance and use by  
36 salmonids in these habitats is not clear (Simenstad et al. 2003). Historically, LWD was  
37 found in great abundance in some Pacific Northwest estuaries, influencing the formation  
38 and distribution of estuarine channels (Collins et al. 2002).

### 39 **3.8.5 Fish and Fish Habitat by Analysis Region**

40 For the purposes of this analysis, the State has been divided into 12 analysis regions. The  
41 12 regions are groupings of smaller units known as Washington Resource Inventory Areas  
42 (WRIAs). The State includes 62 WRIAs, which are primarily divided along major





1 drainage areas or combinations of smaller drainage areas and have State surface water  
2 regulatory status (See <http://salmon.scc.wa.gov/> for a map of WRIAs). A map showing the  
3 12 regions along with the land areas covered by Washington Forest Practices Rules is  
4 provided in Figure 3-1.

5 The analysis regions are listed below, and their fish and fish habitats are summarized in the  
6 following subsections. They are described in greater detail in DEIS Appendix A.

7 Western Washington

- 8 • North Puget Sound
- 9 • South Puget Sound
- 10 • West Puget Sound
- 11 • Islands
- 12 • Olympic Coast
- 13 • Southwest
- 14 • Lower Columbia

15 Eastern Washington

- 16 • Middle Columbia
- 17 • Upper Columbia below Grand Coulee Dam
- 18 • Upper Columbia above Grand Coulee Dam
- 19 • Snake River
- 20 • Columbia Basin

21 The distribution of the covered fish species and other affected fish species, as well as the  
22 and State and private forestlands that are subject to Washington Forest Practices Rules,  
23 varies within each of the regions (DEIS Appendix A). In addition, the number and type of  
24 factors that influence the current conditions of the aquatic system and status of the covered  
25 species in each of the regions may be very different. NMFS sometimes refers to general  
26 factors affecting listed salmonid species as “the four Hs,” which include habitat, harvest,  
27 hatcheries, and hydropower (Federal Caucus 1999; see also  
28 [http://www.salmonrecovery.gov/Archive\\_chronological.shtml](http://www.salmonrecovery.gov/Archive_chronological.shtml)). The Washington Forest  
29 Practices Rules are generally considered to affect only the habitat part of these complex  
30 issues. In addition, other land-use practices such as agriculture and urbanization can also  
31 have a substantial effect on habitat.

32 One of the regions, Islands, does not weigh heavily in the analysis for fisheries because  
33 only a relatively small number of streams exist in forested portions of these regions, or  
34 they contain low numbers of fish species. The following is a short synopsis of the  
35 remaining 11 regions in regards to the fish species present and the components of the four  
36 Hs affecting their ESA status. Table 3-22 shows the distribution of streams among

37



## Chapter 3

1 **Table 3-22.** Percentage of Stream Miles among Forested Ownership Categories  
2 and Non-forest by Analysis Region in Washington State.

Analysis Region	Percent of Stream Miles within Analysis Region (%)					Total Miles
	Federal Forested	Tribal Forested	State Fish and Wildlife and Parks Forested	Forestlands Subject to Washington Forest Practices	Non-Forested	
North Puget Sound	35	0	0	39	25	28,653
South Puget Sound	17	0	0	62	20	13,832
West Puget Sound	29	1	1	54	16	9,114
Islands	2	0	4	49	46	1,009
Olympic Coast	30	11	0	50	9	14,959
Southwest	3	0	0	86	10	28,607
Lower Columbia	23	0	0	63	14	29,645
Middle Columbia	19	8	1	20	53	32,878
Upper Columbia (Downstream of Grand Coulee Dam)	36	2	1	11	50	38,869
Upper Columbia (Upstream of Grand Coulee Dam)	24	19	0	31	25	33,917
Snake River	10	0	1	5	84	19,488
Columbia Basin	0	0	0	0	99	14,157
<b>Total State</b>	<b>22</b>	<b>4</b>	<b>0</b>	<b>37</b>	<b>36</b>	<b>265,129</b>

Source: Washington DNR Hydrography layer and Major Public Lands layer, 2004; U.S. Geological Survey National Land Cover Data layer, 2004.

3  
4 different forest ownership and non-forested categories, including the percentage of all  
5 streams that are subject to Washington Forest Practices Rules in each region. This  
6 information provides an indication of the type of management approach that might be  
7 prevalent on the waters of each region. For example, in regions in western Washington  
8 (North, South, and West Puget Sound; Olympic Coast; Southwest; and Lower Columbia),  
9 streams within Federal ownership are managed based upon the Aquatic Conservation  
10 Strategy outlined in the Northwest Forest Plan (USDA Forest Service and USDI Bureau of  
11 Land Management 1994).

12 In addition, the majority of the Federal lands occur further up the watershed, occupy higher  
13 elevations, and are steeper than the majority of other lands in Table 3-21. This can be  
14 important from the perspective of sediment production and delivery. High gradient, non-  
15 fish-bearing streams are generally source and transport reaches for sediment; low gradient,  
16 fish-bearing streams are areas of sediment accumulation. The information provides some  
17 insights on which regions might be most affected by the alternatives described in Chapter 2  
18 (Alternatives). The information presented below is a summary of the information  
19 presented in DEIS Appendix A. See these documents for more complete descriptions  
20 along with references for the observations presented.



### 1 **3.8.5.1 North Puget Sound**

2 The North Puget Sound Region includes the northeast portion of Puget Sound (WRIAs 1,  
3 3, 4, 5, and 7; see North Puget Sound Region map in DEIS Appendix A), south of the  
4 Canadian border, exclusive of the San Juan and Whidbey Islands (the Islands Region), and  
5 down to the southern border of the Snohomish River System. Other major river systems  
6 include the Nooksack, Skagit, Sauk, and Stillaguamish. The Region contains an estimated  
7 28,653 stream miles.

8 Many of the fish species are present in the North Puget Sound Region. Chinook and bull  
9 trout are listed as threatened in the Region. Coho salmon and green sturgeon are species of  
10 concern. Other fish species with Federal or State listing status include coastal cutthroat,  
11 Pacific lamprey, river lamprey, and Salish sucker. Each of the four Hs has been cited as  
12 contributing to the listing of one or more of the species. Hydropower dams and storage  
13 facilities have impacted stream flows, channel morphology, and fish habitat (e.g., Skagit  
14 River, Baker River, Tolt River) (Washington State Conservation Commission 2003).  
15 Various watershed analyses and limiting factors analyses have documented mass wasting  
16 as being one of the most substantial impacts associated with recent forest practices,  
17 primarily from clearcuts and roads (Washington DNR 1993, 1997a, 1997b; Washington  
18 State Conservation Commission 1999a, 2002). Many of the lowland areas of the Region  
19 are highly urbanized, resulting in loss or degradation of floodplain and off channel habitats,  
20 loss of wetlands, and overall reduction of riparian forests. This Region is one of the most  
21 heavily populated regions of the State.

22 About 39 percent of the streams occur on lands subject to the Washington Forest Practices  
23 Rules (Table 3-21). Federal forest management strategies occur along approximately 35  
24 percent of the streams in the Region. Almost all of the remaining streams flow through  
25 non-forested lands. All State lands within the range of the northern spotted owl have been  
26 operating under the State Trust Lands HCP ITP issued by the Services in 1997  
27 (Washington DNR 1997d).

### 28 **3.8.5.2 South Puget Sound**

29 The South Puget Sound Region wraps around the southeastern and southern edge of Puget  
30 Sound, and includes six WRIAs (8, 9, 10, 11, 12, and 13; see South Puget Sound Region  
31 map in DEIS Appendix A). Major stream systems include Lake Washington, Cedar River,  
32 Sammamish River, Green River, Duwamish River, Soos Creek, Puyallup River, White  
33 River, Carbon River, Nisqually River, Deschutes River, and South Sound tributaries. The  
34 Region contains an estimated 13,832 stream miles.

35 Chinook and bull trout are federally listed as threatened, and coho and green sturgeon are  
36 Federal species of concern. Other fish species with Federal or State listing status include  
37 coastal cutthroat, Pacific lamprey, river lamprey, pygmy whitefish, Olympic mudminnow,  
38 and Salish sucker. Each of the four Hs has been cited as contributing to the listing of one  
39 or more of the species. Hydropower dams and storage facilities have impacted stream  
40 flows, channel morphology and fish habitat (e.g., Cedar River, Green River, White River,  
41 Puyallup River, and Nisqually River) (Hunter 1992). This Region is the most developed  
42 and heavily populated region of the State. Managed forestlands are fragmented and sparse



## Chapter 3

---

1 in the floodplains and lower foothills. The impact of urban development on nearshore  
2 areas, estuaries, freshwater wetlands, and floodplains is moderate to severe. Some of the  
3 remaining managed forestlands in the Cascade Range and higher foothills are vulnerable to  
4 landslides (Washington State Conservation Commission 1999b, 2000a, 2001; Washington  
5 DNR 1996, 1998a, 1998b, 2002).

6 About 62 percent of streams occur on lands subject to the Washington Forest Practices  
7 Rules (Table 3-22). Federal forest management strategies occur along approximately 17  
8 percent of the streams. Almost all of the remaining streams flow through non-forested  
9 lands. A substantial portion of the State and private forestlands in this Region is currently  
10 managed under HCPs. All State lands within the range of the northern spotted owl have  
11 been operating under the State Trust Lands HCP since 1997 (Washington DNR 1997d).

### 12 **3.8.5.3 West Puget Sound**

13 The West Puget Sound Region includes five WRIs (14, 15, 16, 17, and 18; see West  
14 Puget Sound Region map in DEIS Appendix A). Major stream systems include the  
15 Skokomish, Duckabush, Dosewalips, Big Quilcene, Elwha, and Dungeness Rivers, as well  
16 as other South Sound and Hood Canal tributaries. The Region contains an estimated 9,114  
17 stream miles.

18 Fish species within the Region include bull trout, Hood Canal chum, and chinook, which  
19 are federally listed as threatened; coho, coastal cutthroat, and green sturgeon, which are  
20 Federal species of concern. Other fish species with Federal or State listing status include  
21 Pacific lamprey, river lamprey, Olympic mudminnow, Salish sucker. Each of the four Hs  
22 has been cited as contributing to the listing of one or more of the species. Notably, the two  
23 major hydroelectric dams on the Elwha River have blocked large portions of salmonid  
24 spawning habitat and are scheduled for demolition starting in 2008  
25 (<http://www.nps.gov/olym/elwha/home.htm>). Summer irrigation (Dungeness River) and  
26 groundwater withdrawals also contribute to reduced stream flows.

27 Managed forestlands throughout the Puget Lowlands are becoming increasingly  
28 fragmented by urban development, although some large commercial timber plantations  
29 remain on the western side of the Kitsap Peninsula and in eastern Jefferson County.  
30 However, most of these forestlands in Kitsap and eastern Jefferson Counties have not been  
31 designated under the Growth Management Act as forest resource lands. Agricultural uses  
32 are common in the floodplains of the area. Recreational, residential, and limited urban  
33 development has resulted in a substantial impact, especially along the marine shorelines.  
34 Most of the larger rivers drain from the Olympic National Park and U.S. Forest Service  
35 lands; thus, many of the upper watersheds are substantially protected. However, timber  
36 harvest and the associated forest road construction occurred in some of the high Olympics  
37 in the South Fork Skokomish and Dungeness Basins. These forest practices were followed  
38 by severe landslide episodes (Bountry et al. 2002; Washington DNR 1997c). Private and  
39 State commercial timber plantations are present around the fringes of this Federal land and  
40 occupy most of the foothills.



1 About 54 percent of streams occur on lands subject to the Washington Forest Practices  
2 Rules (Table 3-22). Federal forest management strategies occur along approximately 29  
3 percent of the streams. Almost all of the remaining streams flow through non-forested  
4 lands. A substantial portion of the State and private forestlands in this Region is currently  
5 managed under HCPs. All State lands within the range of the northern spotted owl have  
6 been operating under the State Trust Lands HCP since 1997 (Washington DNR 1997d)

### 7 **3.8.5.4 Olympic Coast**

8 The Olympic Coast Region (WRIAs 19, 20, and 21; see Olympic Coast Region map in  
9 DEIS Appendix A) includes coastal rivers and streams from the north of and including the  
10 Copalis River to the west of, but not including, the Elwha River. The Region contains an  
11 estimated 14,959 miles of stream.

12 Many of the covered fish species are present in the Olympic Coast Region. Bull trout are  
13 listed as threatened throughout the Region, and the Ozette Lake population of sockeye  
14 salmon is listed as threatened. Other fish species with Federal or State listing status  
15 include coastal cutthroat, Pacific lamprey, pygmy whitefish, and Olympic mudminnow. Of  
16 the four Hs, habitat appears to be the highest priority factor for bull trout. Historic timber  
17 harvest, road construction, and forest fires have had substantial impacts on fish habitat. No  
18 notable hydroelectric facilities are present in the Region, and no hatcheries are stocking  
19 bull trout or sockeye salmon. However, small diversion dams for agricultural purposes are  
20 present in some watersheds.

21 About 50 percent of streams occur on lands subject to the Washington Forest Practices  
22 Rules (Table 3-22). Federal forest management strategies occur along approximately 30  
23 percent of the streams, although a significant portion of these streams are in Olympic  
24 National Park or National Forest wildernesses. Tribal forest management (on the Makah  
25 and Quinault Indian Reservations) occurs along approximately 11 percent of the streams.  
26 Essentially all of the remaining streams flow through non-forested lands. A substantial  
27 portion of the State forestlands in this Region is currently managed under an HCP. All  
28 State lands within the range of the northern spotted owl have been operating under the  
29 State Trust Lands HCP since 1997 (Washington DNR 1997d).

### 30 **3.8.5.5 Southwest**

31 The Southwest Region (WRIAs 22, 23, and 24; see Southwest Region map in DEIS  
32 Appendix A) includes coastal rivers and streams north of the Columbia River to the Grays  
33 Harbor drainage. The Region contains an estimated 28,607 miles of stream.

34 Many covered species are present in this Region. Bull trout are listed as a threatened  
35 species. Other fish species with Federal or State listing status include coastal cutthroat,  
36 Pacific lamprey, and Olympic mudminnow. Similar to the Olympic Coast Region, habitat  
37 degradation appears to be the leading factor influencing listing of species in the Region.  
38 Fine sediment is the key limiting factor in much of the coastal hill drainages (DEIS  
39 Appendix A). Landslides and unpaved roads are both substantial contributors. The coastal  
40 foothills are one of the most landslide sensitive areas of the State. Urbanization and



## Chapter 3

---

1 agricultural development have impacted the Chehalis Valley and, to a lesser degree, the  
2 small coastal valleys (DEIS Appendix A).

3 Streams in the Region are substantially influenced by the Washington Forest Practices  
4 Rules; about 86 percent of them are on State, private, and other lands that are subject to  
5 Washington Forest Practices Rules. Federal management strategies have only a minor  
6 influence on streams in the Region with only 3 percent under Federal management.  
7 Essentially all the remaining lands flow through non-forested lands. State lands in the  
8 Region are covered by the State Trust Lands HCP (Washington DNR 1997d).

### 9 **3.8.5.6 Lower Columbia River**

10 The Lower Columbia Region (WRIAs 25, 26, 27, 28; see Lower Columbia River Region  
11 map in DEIS Appendix A) includes the Columbia River and rivers and streams that drain  
12 from Washington into the Columbia River from its mouth to streams west of (but exclusive  
13 of) Rock Creek. The Region contains an estimated 29,645 stream miles.

14 Many of the covered species are present in this Region. Sockeye do not spawn or rear in  
15 the Region, but use the mainstem Columbia River as a migration corridor, as do Oregon  
16 chinook and steelhead stocks. Chinook salmon, chum salmon, and steelhead are listed as  
17 threatened in the Region and found downstream of Mossyrock Dam and Merwin Dam on  
18 the Cowlitz River and Lewis River, respectively. Bull trout are listed as threatened  
19 throughout the Region. Coho salmon and green sturgeon are candidate species. Other fish  
20 species with Federal or State listing status include coastal cutthroat, Pacific lamprey, river  
21 lamprey, leopard dace, mountain sucker, sandroller. Each of the four Hs has been cited as  
22 contributing to the listing of one or more of the species. Hydropower development is one  
23 of the largest impacts to salmonid habitat in the Lower Columbia Region (DEIS Appendix  
24 A). Construction of dams on the Cowlitz and Lewis Rivers has removed hundreds of  
25 stream miles from anadromous fish access (Washington State Conservation Commission  
26 2000b, 2000c). The May 1980 eruption of Mount St. Helens had major impacts to riparian  
27 zones and channels of the Cowlitz and Toutle Rivers. Fine sediment is one of the key  
28 limiting factors in the coastal hills and many of the Cascade foothills and mountains,  
29 contributed substantially by landslides and unpaved roads (DEIS Appendix A).  
30 Urbanization and agricultural development has impacted most of the larger valleys,  
31 especially Cowlitz Valley and eastern Clark County (Washington State Conservative  
32 Commission 2000b, 2000c).

33 About 63 percent of streams occur on lands subject to the Washington Forest Practices  
34 Rules (Table 3-22). Federal forest management strategies occur along approximately 23  
35 percent of the streams. Almost all of the remaining streams flow through non-forested  
36 lands. State lands in the Region are covered by the State Trust Lands HCP (Washington  
37 DNR 1997d).

### 38 **3.8.5.7 Middle Columbia River**

39 The Middle Columbia Region (WRIAs 29, 30, 31, 37, 38, and 39; see Middle Columbia  
40 River Region map in DEIS Appendix A) includes rivers and streams that drain from  
41 Washington State to the Columbia River from Rock Creek through the Yakima River, not



1 including the Snake and Walla Walla Rivers. The Region contains an estimated 32,878  
2 miles of stream.

3 Many of the covered species are present in this Region. Sockeye do not spawn or rear in  
4 the Region, but use the mainstem Columbia River as a migration corridor. Chinook,  
5 steelhead, and bull trout are listed as threatened. Other fish species with Federal or State  
6 listing status include cutthroat, Pacific lamprey, river lamprey, pygmy whitefish, leopard  
7 dace, Umatilla dace, mountain sucker, and sandroller. Each of the four Hs has been cited  
8 as contributing to the listing of one or more of the species. Primary limiting factors for fish  
9 and associated habitats within the Region are generally a result of hydropower, water  
10 storage, logging, farming, grazing, urban and suburban development, irrigation,  
11 transportation, and industrial or nuclear development. Irrigation and hydropower  
12 development have had some of the largest effects within much of this Region (Harding  
13 2001; Washington State Conservation Commission 1999). Hydropower and irrigation  
14 storage reservoirs have reduced or eliminated up and downstream fish passage on the  
15 mainstem Columbia and headwater areas. Irrigation reservoirs and diversions have  
16 affected passage in tributaries by altering seasonal patterns or reducing instream flow,  
17 which affects habitat and influences water quality conditions in much of the Region (DEIS  
18 Appendix A).

19 About 20 percent of streams occur on lands subject to the Washington Forest Practices  
20 Rules (Table 3-22). Federal forest management strategies occur along approximately 19  
21 percent of the streams. Tribal management (primarily on the Yakama Indian Reservation)  
22 occurs along 8 percent of the streams. Almost all of the remaining streams flow through  
23 non-forested lands.

### 24 **3.8.5.8 Upper Columbia River - Downstream of Grand Coulee Dam**

25 The Upper Columbia River (downstream of Grand Coulee Dam) Region includes the  
26 mainstem of the Columbia River and its tributaries to Grand Coulee Dam (WRIAs 40 and  
27 44 through 50; see Upper Columbia River - Downstream Region map in DEIS Appendix  
28 A). The Region has 38,869 miles of mapped streams. The major tributaries include the  
29 Wenatchee River, Methow River, Okanogan River, and Lake Chelan and its tributaries.

30 Upper Columbia chinook and steelhead are federally listed as endangered; bull trout are  
31 listed as threatened. Other fish species with Federal or State listing status include  
32 cutthroat, Pacific lamprey, pygmy whitefish, leopard dace, Umatilla dace, mountain sucker,  
33 and sandroller. Each of the four Hs has been cited as contributing to the listing of one or  
34 more of the species. The predominant limiting factor of the Columbia mainstem has  
35 generally been the result of hydropower development and storage dams. Other activities  
36 have also contributed to habitat degradation associated with farming, irrigation, livestock  
37 grazing, logging, urban and suburban development, and transportation.

38 About 11 percent of the streams occur on lands subject to the Washington Forest Practices  
39 Rules (Table 3-22). Federal forest management strategies occur along approximately 36  
40 percent of the streams. Tribal management occurs along 2 percent of the streams, and  
41 about 50 percent of the stream miles are in non-forested lands.



## Chapter 3

---

### 1 **3.8.5.9 Upper Columbia River - Upstream of Grand Coulee Dam**

2 The Upper Columbia River (upstream of Grand Coulee Dam) Region includes all of the  
3 Columbia River mainstem and its tributaries upstream of Grand Coulee Dam within  
4 Washington (WRIAs 51 through 62; see Upper Columbia River - Upstream Region map in  
5 DEIS Appendix A). Major tributaries include the Sanpoil River, Spokane River, Kettle  
6 River, and Colville River. The Region contains an estimated 33,917 stream miles.

7 Grand Coulee Dam is a complete barrier to anadromous fish. Consequently, the only  
8 federally listed species present in this Region is bull trout, which are listed as threatened.  
9 Other fish species with Federal or State listing status include westslope cutthroat, redband  
10 trout, pygmy whitefish, Umatilla dace, lake chub, and sandroller. Hydroelectric and  
11 irrigation dams, which have fragmented bull trout distribution, plus habitat degradation,  
12 have been major factors leading to the listing in this Region (U.S. Federal Register, Vol.  
13 63, No. 111, June 10, 1998, pages 31647-31674). Timber harvesting, agriculture, and  
14 livestock grazing have degraded habitats in both upland and riparian areas.

15 About 31 percent of the streams occur on lands subject to the Washington Forest Practices  
16 Rules (Table 3-22). Federal forest management strategies occur along approximately 24  
17 percent of the streams. Tribal management (on the Colville Indian Reservation) occurs  
18 along 19 percent of the streams, and about 25 percent of the stream miles are in non-  
19 forested lands.

### 20 **3.8.5.10 Snake River**

21 The Snake River Region (WRIAs 32, 33, 34, and 35; see Snake River Region map in DEIS  
22 Appendix A) includes all portions of the Snake River and its tributaries that lie within  
23 Washington State. The Region also includes the Walla Walla River drainage. The Snake  
24 River Region is relatively arid, but contains an estimated 19,488 stream miles.

25 Sockeye salmon are federally listed as endangered. Chinook salmon, steelhead, and bull  
26 trout are listed as threatened. Sockeye salmon, however, do not spawn or rear in the  
27 Region, but use the mainstem Snake River as a migration corridor. Sockeye spawning and  
28 rearing occur within Idaho. Other fish species with Federal or State listing status include  
29 cutthroat, Pacific lamprey, mountain sucker, sandroller, and margined sculpin. Each of the  
30 four Hs has been cited as contributing to the listing of one or more of the species.  
31 Hydropower has altered the natural hydrograph of the Snake River and impacted fish  
32 habitat. Other habitat limiting factors within the Region include till crop production and  
33 irrigation withdrawals, livestock grazing, logging, urbanization, and transportation  
34 networks.

35 About 5 percent of the streams occur on lands subject to Washington Forest Practices  
36 Rules (Table 3-22). Federal forest management strategies occur along approximately 10  
37 percent of the streams in the Region. About 84 percent of the mapped streams in the  
38 Region flow through non-forested lands. Relative to other regions, the Snake River Region  
39 is relatively arid and does not include large amounts of commercial forestlands.





## 3.9 AMPHIBIANS AND AMPHIBIAN HABITAT

### 3.9.1 Introduction

This section describes the biology of the seven amphibian species that: 1) are considered sensitive to riparian forest practices, and 2) have been chosen for coverage under the FPHCP.

- Coastal (Pacific) tailed frog (*Ascaphus truei*)
- Rocky Mountain (inland) tailed frog (*Ascaphus montanus*)
- Van Dyke’s salamander (*Plethodon vandykei*)
- Dunn’s salamander (*Plethodon dunni*)
- Olympic torrent salamander (*Rhyacotriton olympicus*)
- Columbia torrent salamander (*Rhyacotriton kezeri*)
- Cascade torrent salamander (*Rhyacotriton cascadae*)

These seven species were selected because: 1) they are closely associated with aquatic and riparian habitats, 2) they are thought to have some sensitivity to forest practices, and 3) they lack Federal protection in some portion of their range (either through status or occurrence on Federal lands). Other aquatic or riparian-associated species with special status are addressed in subsection 3.10 (Birds, Mammals, Other Wildlife and Their Habitats).

The following subsections provide information about the distribution, regulatory status, and habitat associations of the seven selected species. Table 3-23 summarizes the distribution of these species among the 10 analysis regions with substantial forestlands. More detailed regional information is provided in DEIS Appendix A. Descriptions of known distributions may be conservative because no systematic surveys have been conducted in most areas.

**Table 3-23.** Distribution of the Selected Amphibian Species by Analysis Region.<sup>1/</sup>

Species	North Puget Sound	South Puget Sound	West Puget Sound	Olympic Coast	Southwest Washington	Lower Columbia River	Middle Columbia River	Snake River	Upper Columbia below Grand Coulee	Upper Columbia above Grand Coulee
Van Dyke’s salamander		X	X	X	X	X				
Dunn’s salamander					X	X				
Olympic torrent salamander			X	X	X					
Columbia torrent salamander					X	X				
Cascade torrent salamander			?		X	X	X			
Coastal (Pacific) tailed frog	X	X	X	X	X	X	X		X	
Rocky Mountain (inland) tailed frog								X		?

<sup>1/</sup> X = documented occurrences; ? = no known occurrences, but may occur; [Blank] = not expected to occur.

Source: DEIS Appendix A.



## Chapter 3

---

### 3.9.2 Amphibian Distribution, Status, and Habitat

#### 3.9.2.1 Van Dyke's Salamander

Endemic to Washington, the Van Dyke's salamander is known from three discrete regional centers: the Olympic Mountains, the southern Cascades (including populations in southeastern Thurston County), and the Willapa Hills (Leonard et al. 1993). Van Dyke's salamanders have been found at elevations ranging from sea level to 5,000 feet, in areas with an average annual precipitation of at least 59 inches (Jones 1999; Wilson et al. 1995). Most recorded locations come from the wetter, western slopes of these areas (Dvornich et al. 1997). Populations are thought to be patchily distributed and often appear to be in low densities with much potential habitat apparently unoccupied (Blaustein et al. 1995; Jones 1999), but focused surveys for this species have not been done over much of its range. The Regional Summaries (DEIS Appendix A) describe this species' distribution as intermittent throughout the middle and upper elevations of WRIAs 10, 11, 16, and 19 through 27.

The Van Dyke's salamander is a USFWS species of concern and a Washington State candidate species. Limited distribution and isolation of Van Dyke's salamander populations have prompted concern for this species' persistence (Holthausen et al. 1994; Nordstrom and Milner 1997). Lehmkuhl and Ruggiero (1991) assigned this species a high risk of local extinction, based on its habitat associations, frequency of occurrence, abundance, and dispersal ability. Similarly, Thomas et al. (1993) identified the Van Dyke's salamander as a high-risk species, closely associated with old-growth forest conditions. Two out of three regions where this species occurs are dominated by Federal ownership (Olympic National Park and Wilderness Area, Mount St. Helens National Monument, Gifford Pinchot National Forest), and the third is dominated by private commercial forestlands (southwest Washington). Unpublished studies by the U.S. Forest Service in the Olympic Coast Region indicate that this species occurs at low densities in landscapes where timber harvest has occurred (DEIS Appendix A).

Van Dyke's salamanders are most commonly associated with riparian habitats, or with cool, moist microsites within other habitat types (Jones 1999; Nordstrom and Milner 1997; Petranka 1998). Juveniles and adults have been found in the splash zones of streams where a thin film of water runs between or under rocks adjacent to the stream margin, upland forests, moist talus, cave entrances, seeps, and along lakeshores (Blaustein et al. 1995; Jones 1999). Of six Van Dyke's salamander nests that have been located and described to date, five were near small, headwater streams (Blessing et al. 1999; Jones 1989); no information is available on the location of the sixth (Noble 1925). Clutches of eggs, apparently laid during spring, have been found under rocks or inside large, moss-covered logs; eggs may require more than 4 months to hatch, nearly twice as long as the incubation period of other similar salamander species in this area (Blessing et al. 1999). Jones (1999) indicates that Van Dyke's salamanders may be found near streams and seeps that are perennial, or spatially or temporally intermittent (i.e., surface water may be absent during some periods or in some stretches).



### 1 **3.9.2.2 Dunn's Salamander**

2 Dunn's salamanders occur in rocky forest habitats from sea level to 3,300 feet, from  
3 northwestern California to extreme southwestern Washington (Leonard et al. 1993;  
4 Nussbaum et al. 1983; Petranka 1998). The Willapa Hills represent the northernmost limit  
5 of this species' range, which extend north to the south side of the Chehalis River floodplain  
6 (Leonard et al. 1993). Most of the recorded locations for this species in Washington State  
7 come from Pacific, Lewis, Wahkiakum, and Cowlitz Counties (Dvornich et al. 1997). The  
8 regional summaries indicate that Dunn's salamanders are found in only 2 of the 10 regions,  
9 with a widespread distribution south of the Chehalis River in WRIAs 22 through 25, and  
10 along the western edge of WRIA 26 (DEIS Appendix A).

11 The Dunn's salamander is a Washington State candidate species; concern for this species  
12 in the State is prompted by its distribution in small and fragmented populations, and by  
13 Washington's position at the northernmost end of the salamander's range (Nordstrom and  
14 Milner 1997). Dunn's salamanders are relatively common in the Oregon Coast Range,  
15 locally common along larger streams in Washington (Personal Communication, Marc  
16 Hayes, Washington Department of Fish and Wildlife, October 2003), but become  
17 uncommon to rare in headwater habitats in Washington (Jackson et al. 2003). Private  
18 commercial timberlands dominate most of this species' range in southwestern Washington.

19 Dunn's salamanders have been found inhabiting heavily shaded, wet, rocky substrates such  
20 as seeps, moist talus slopes, and stream edges in forested areas (Leonard et al. 1993;  
21 Nordstrom and Milner 1997). Information on the life history of Dunn's salamanders is  
22 scarce, based on two nesting records (Dumas 1955; Nauman et al. 1999) and comparisons  
23 to similar species (Petranka 1998). Eggs are probably laid underground in rocky areas or  
24 within woody debris during spring and hatch in late summer or fall. Juveniles, which may  
25 take 2 to 4 years to reach sexual maturity, have been found in the same habitats as adults  
26 (Petranka 1998). In the Oregon Coast Range, Corn and Bury (1991) found a strong  
27 association between the abundance of Dunn's salamanders and the percent cover of rock.  
28 They also found that Dunn's salamanders occurred more often on steep slopes where  
29 exposed talus was present. Wilkins and Peterson (2000) found an increased probability of  
30 Dunn's salamander occurrence along streams with steep maximum sideslopes; the species  
31 was present at all streams with sideslopes greater than 80 percent gradient, and absent from  
32 streams with a sideslope less than 50 percent. Though usually associated with rock,  
33 Dunn's salamanders also use downed logs and woody debris for cover and feeding  
34 (Corkran and Thoms 1996; Leonard et al. 1993).

35 Dunn's salamanders are not considered aquatic, but rather riparian associates (Corkran and  
36 Thoms 1996; Gomez and Anthony 1996). Approximately 90 percent of Dunn's  
37 salamanders observed by Bury et al. (1991) were found in streambank habitat as opposed  
38 to riffle or pool habitat. Wilkins and Peterson (2000) confirm the strong riparian  
39 association of this species, noting a marked decrease in captures at microsites greater than  
40 3 feet (1 meter) from perennial stream channels.



## Chapter 3

---

### 3.9.2.3 Olympic Torrent Salamander

Until recently, all torrent salamanders were considered to be a single species (*Rhyacotriton olympicus*) with two subspecies (Petranka 1998). Studies of protein variation, however, provided evidence of genetic differences among four geographically isolated subgroups, resulting in the recognition of four distinct species (Good and Wake 1992). Three of these species (Olympic, Columbia, and Cascade torrent salamanders) occur in Washington State, and are addressed in the EIS. Little has been written about the specific habitat needs of the different torrent salamander species. Most information comes from studies that did not distinguish among *Rhyacotriton* species, or that focused on other members of this species group. All four species are similar enough to have been considered as one species until very recently; therefore, the torrent salamanders of Washington State likely have similar habitat needs. Much of the following discussion is based on studies of the southern torrent salamander (*Rhyacotriton variegatus*).

Olympic torrent salamanders are known to occur only on the northern portion of the Olympic Peninsula (Leonard et al. 1993; Petranka 1998). Most recorded locations for this species come from Clallam, Jefferson, and Mason Counties (Dvornich et al. 1997). The Regional Summaries indicate that this species is widespread within WRIAs 14 through 24, although it is absent from the Puget Lowlands and from areas south of the Chehalis River (DEIS Appendix A).

The Olympic torrent salamander is a USFWS species of concern and a Washington State monitor species. As with the other torrent salamander species in Washington, concern for this species centers on its limited distribution, narrow range of tolerance for environmental conditions, and the associated risk of local extirpation following clearcut timber harvest (Blaustein et al. 1995; Bury and Corn 1988; Hallock and McAllister 2002). Most of the range of the Olympic torrent salamander occurs on Federal lands (mainly Olympic National Park and nearby wilderness areas). Unpublished studies by the U.S. Forest Service in the Olympic Coast Region indicate that this species occurs at low densities in landscapes where timber harvest has occurred (DEIS Appendix A).

Torrent salamanders inhabit cold, permanent, heavily shaded seeps and headwater streams in humid forests (Good and Wake 1992; Nussbaum et al. 1983). Available data indicate that torrent salamanders require at least 6 years to achieve adulthood, spending the bulk of that time in their natal streams either as eggs or aquatic larvae (Nussbaum and Tait 1977). Eggs are probably laid singly, unattached and scattered about in deep cracks and crevices; egg-laying may occur throughout the year, but peaks from May through July (Nussbaum and Tait 1977). Breeding habitat is generally considered to be forested permanent seeps, streams, and waterfalls with rocky substrates and cold temperatures (optimum 46 to 55 °F [8° to 13°C]); foraging occurs in moist areas in or near streams and seeps (Corn and Bury 1991; Diller and Wallace 1996; Leonard et al. 1993; Welsh and Lind 1996).

Torrent salamanders are strongly associated with non-fish-bearing waters high in the stream network (Jackson et al. 2003). When torrent salamanders occur along large fish-bearing streams, they are usually found in margins where they can find cover from predators (e.g., fish and Pacific giant salamanders) in the spaces between stones, or in



1 seeps on the valley wall rather than within the stream itself (Welsh 1993; Welsh and  
2 Ollivier 1992; Wilkins and Peterson 2000). Perhaps because they are very sensitive to  
3 drying out (Ray 1958), torrent salamanders are relatively sedentary, remaining near water  
4 and seldom moving more than 6 feet from one location (Nussbaum and Tait 1977).  
5 Individuals have been found more than 150 feet from permanent water, but such  
6 movements are thought to be rare (Good and Wake 1992).

### 7 **3.9.2.4 Columbia Torrent Salamander**

8 The Columbia torrent salamander is distributed in the Coast Ranges of Washington and  
9 Oregon from the Willapa Hills of Washington to the Grand Ronde River Valley in Oregon  
10 (Leonard et al. 1993). The valley of the Chehalis River separates the range of this species  
11 from that of the Olympic torrent salamander. Most recorded locations for this species in  
12 Washington come from Pacific, Lewis, and Wahkiakum Counties (Dvornich et al. 1997).  
13 The Regional Summaries (DEIS Appendix A) describe this species as widespread within  
14 its limited distribution south of the Chehalis River in WRIAs 22 through 25, and the  
15 western edge of WRIA 26. Although other torrent salamander species have substantial  
16 ranges in Federal lands with long-term management objectives for old-growth forest  
17 conditions (Tuchmann et al. 1996), over 95 percent of the known distribution of the  
18 Columbia torrent salamander is on private and State lands subject to intensive timber  
19 harvest (Good and Wake 1992; McAllister 1995). Few studies have addressed the ecology  
20 of Columbia torrent salamanders, but like other torrent salamanders, this species is  
21 presumed to require habitats best provided by old-growth forests (e.g., cold, shaded  
22 streams with clean gravel and cobble substrates; Bury et al. 1991; Corn and Bury 1989;  
23 Welsh and Lind 1996).

24 The Columbia torrent salamander is a USFWS species of concern and a Washington State  
25 candidate species, and its range consists primarily of privately owned commercial  
26 timberlands. Recent studies have revealed Columbia torrent salamanders to occur in  
27 extraordinarily high densities in headwater habitats on private timberlands in the Willapa  
28 Hills of southwestern Washington (Personal Communication, Marc Hayes, Washington  
29 Department of Fish and Wildlife, October 2003) and the Coast Ranges of northwestern  
30 Oregon (Russell et al. 2004), possibly reducing the levels of concern for this species. As  
31 stated above, habitat requirements of the Columbia torrent salamander are likely similar to  
32 those of the Olympic torrent salamander. However, a recent study on private timberlands  
33 may indicate that this torrent salamander responds differently than other torrent  
34 salamanders to the influence of forest practices (Russell et al. 2004).

35 Results from Russell et al. (2004) indicate that attributes of overstory or understory  
36 vegetation were not related to the distribution and density of Columbia torrent salamanders  
37 at either the landscape or stream-reach scales. Two factors were identified to explain the  
38 possible insensitivity of this species (i.e., wide distribution and high abundance) to  
39 variations in vegetative conditions. First, in the Oregon Coast Range, plant growth is so  
40 rapid that post-harvest vegetation often grows as high as the base of tree crowns in  
41 adjacent, unharvested stands in as little as 10 years (Hibbs and Bower 2001). Thus, direct  
42 solar radiation and airflows quickly become limited and moist, humid microclimates  
43 similar to those of unharvested areas are quickly reestablished (Hibbs and Bower 2001).



## Chapter 3

---

1 Deep (3 to 6 feet [1 to 2 meters]) accumulations of post-logging debris (slash) across small  
2 stream channels are common in recently harvested areas and may adequately buffer  
3 temperatures for stream salamanders (Jackson et al. 2003). Second, the cool, moist climate  
4 across their study area, particularly near the coast, may have mitigated the effects of  
5 periodic canopy removal on water temperatures (Diller and Wallace 1996; Welsh and Lind  
6 1996). The narrow range of water temperatures measured by Russell et al. (2004) in all  
7 streams during June and July (less than 54 °F [11°C]) would suggest that the existing  
8 vegetation and moderate climate of the area are sufficient to maintain favorable water  
9 temperatures for torrent salamanders (less than 61°F [16°C]) (Brattstrom 1963; Diller and  
10 Wallace 1996; Welsh and Lind 1996). However, within the more extreme climates  
11 associated with interior physiographic provinces (e.g., Cascade and Klamath-Siskiyou  
12 Ranges), removal of riparian vegetation may have more pronounced effects on the  
13 distribution and abundance of other torrent salamander species (Welsh and Lind 1996;  
14 Hunter 1998; Welsh et al. 2000).

### 15 **3.9.2.5 Cascade Torrent Salamander**

16 The Cascade torrent salamander occurs along the western slopes of the Cascade Range  
17 from northeastern Lane County, Oregon north to the vicinity of Mount St. Helens (Leonard  
18 et al. 1993). The valley of the Cowlitz River separates the range of this species from that  
19 of the Columbia torrent salamander. Most recorded locations for this species in  
20 Washington come from Skamania, Cowlitz, and Clark Counties (Dvornich et al. 1997).  
21 According to the Regional Summaries (DEIS Appendix A), this species occurs at the  
22 eastern edge of WRIA 23 and the mid-elevation areas of the Cascade slopes in WRIs 26  
23 through 28. It is also predicted to occur at the southern margin of WRIA 11, but no  
24 occurrences have been recorded there.

25 The Cascade torrent salamander is a Washington State candidate species. As with the  
26 other torrent salamander species in Washington, concern for this species centers on its  
27 limited distribution, narrow range of tolerance for environmental conditions, and the  
28 associated risk of local extirpation following clearcut timber harvest (Blaustein et al. 1995;  
29 Bury and Corn 1988; Hallock and McAllister 2002). The range of the Cascade torrent  
30 salamander falls mostly in areas of Federal land (Gifford Pinchot National Forest and  
31 Mount St. Helens National Volcanic Monument).

32 Habitat requirements of the Cascade torrent salamander are likely similar to those of the  
33 Olympic torrent salamander (See above).

### 34 **3.9.2.6 Pacific (coastal) and Rocky Mountain (inland) Tailed Frog**

35 Endemic to the Pacific Northwest, tailed frogs are the only species of the family  
36 Ascaphidae and among the most primitive living frogs. Males are distinguished by a tail-  
37 like appendage that is used for internal fertilization, an adaptation to their life in cold, swift  
38 streams. Based on an examination of genetic differences, Nielson et al. (2001)  
39 recommended that coastal and inland populations of tailed frogs be recognized as distinct  
40 species, *Ascaphus truei* (coastal) and *A. montanus* (inland). To date, no research has  
41 documented differences between the life histories and habitat associations of these species;  
42 therefore, they are treated together in this discussion.

## Chapter 3



1 Tailed frogs occur throughout the Pacific Northwest, with a range that extends from  
2 southwestern British Columbia south to northwestern California (Leonard et al. 1993). In  
3 Washington, Pacific tailed frogs have been found at elevations up to 5,250 feet in the  
4 Cascade and Olympic Mountains as well as the Willapa Hills (Dvornich et al. 1997;  
5 Leonard et al. 1993). The regional summaries show the Pacific tailed frog as having the  
6 widest distribution of any of the seven selected amphibians, occurring in 8 of the 12  
7 regions. This species is generally widespread wherever it occurs, although it is absent  
8 from the highest elevations of the North Puget Sound Region, the lowlands of the West  
9 Puget Sound Region, and the eastern portions of the Middle and Upper Columbia (below  
10 Grand Coulee) Regions.

11 Rocky Mountain tailed frogs have been found in the Blue Mountains in the extreme  
12 southeastern portion of the State (Dvornich et al. 1997, Leonard et al. 1993, United States  
13 Geological Survey 2003). According to the Regional Summaries (DEIS Appendix A), this  
14 species is known from only nine sites in WRAs 32 and 35. The Rocky Mountain tailed  
15 frog also occurs in British Columbia and Idaho immediately adjacent to the Upper  
16 Columbia (above Grand Coulee) Region; future systematic surveys may lead to the  
17 discovery of populations in that region.

18 Both tailed frogs are USFWS species of concern and Washington State monitor species.  
19 Compared to other stream-breeding amphibians, tailed frogs appear to be the most  
20 narrowly distributed in coastal areas of Washington and may be the most sensitive to short-  
21 and intermediate-term effects from timber harvest (Jackson et al. 2003). Tailed frogs have  
22 demonstrated sensitivity to increased levels of fine sediment, which may reduce refuge by  
23 filling instream interstitial spaces in the substrate (Bury and Corn 1988) and also reduce  
24 the availability of algae and other foods important to tadpoles (Welsh and Ollivier 1998).  
25 Local populations are susceptible to extirpation for several reasons, including narrow niche  
26 requirements combined with isolated population distribution, long generation time, and  
27 loss of mature forest along headwater stream habitats (Welsh 1990). Of seven Pacific  
28 Northwest frogs and toads associated with old-growth forest, the tailed frog is probably the  
29 species most likely to be affected by old-growth habitat loss and degradation (Blaustein et  
30 al. 1995). Unpublished studies by the U.S. Forest Service in the Olympic Coast Region  
31 indicate that coastal tailed frogs occur at lower densities in landscapes where timber  
32 harvest has occurred (DEIS Appendix A).

33 Tailed frogs are found almost exclusively in cold, rocky streams; the tadpole's sucker-like  
34 mouth (used for clinging to rocks and scraping away food) is a sign of this species'  
35 adaptation to life in fast-flowing water (Nussbaum et al. 1983; Leonard et al. 1993).  
36 Breeding and developmental habitat for the tailed frog generally consists of perennial, cool  
37 (usually less than 59°F [15°C]) streams with a cobble/boulder substrate and woody debris  
38 (de Vlaming and Bury 1970; Welsh et al. 1993). These conditions are typically associated  
39 with cold, clear, headwater to mid-order streams in mature forest ecosystems (Welsh et al.  
40 1993). Adults forage mainly on land along streambanks but also underwater, seeking  
41 cover under rocks and woody debris in streams (Zeiner et al. 1988). At night, adult tailed  
42 frogs emerge and may forage up to 1,300 feet into adjacent forested areas (McComb et al.  
43 1993). Older (greater than 200 years), multi-layer forests, downed woody material,



## Chapter 3

---

1 ground-level vegetation, ground cover, and canopy closure have been shown to be  
2 important in the occurrence of tailed frogs in northwestern California and southern  
3 Washington (Aubry and Hall 1991; Welsh et al. 1993). The presence of tailed frogs in  
4 younger-age stands indicates that suitable microhabitat conditions can be found in forests  
5 less than 200 years old (Corn and Bury 1989; Aubry and Hall 1991).

6 Tailed frogs typically mate during late August and September; females lay eggs in July of  
7 the year after mating, and larvae (tadpoles) remain in the water for 1 to 4 years after  
8 hatching (Leonard et al. 1993; Nussbaum et al. 1983; Welsh et al. 1993). The tailed frog's  
9 exceptionally long period of larval and pre-reproductive adult development (estimated 7 to  
10 9 years based on studies in Montana) increases the vulnerability of local populations to  
11 habitat disturbance (Brown 1975; Daugherty and Sheldon 1982; Jennings and Hayes  
12 1994). These factors may also increase the amount of time required for recovery following  
13 disturbance (Blaustein et al. 1995). Tailed frog larvae are likely particularly sensitive to  
14 sedimentation following clearcutting along headwater streams; they cannot adhere to rocks  
15 that are coated with fine sediment, and have difficulty moving to find suitable substrate  
16 (Jackson et al. 2003).

### 17 **3.9.3 Review of Timber Harvest Effects on Amphibians**

#### 18 **3.9.3.1 Van Dyke's Salamander**

19 Some studies have suggested that the distribution of Van Dyke's salamander has been  
20 limited by clearcutting (Wilson et al. 1995; Corn and Bury 1989). Welsh (1990) found an  
21 increased risk of local extirpation in intensively managed forests. On the other hand, the  
22 presence of this species in 30- to 60-year-old forests indicates that individuals may persist  
23 within or recolonize disturbed habitats (Nordstrom and Milner 1997; Personal  
24 Communication, Marc Hayes, Washington Department of Fish and Wildlife, October  
25 2003). At this time, more natural history information is needed before conclusions can be  
26 drawn about this species' relation to forest habitat conditions (Blaustein et al. 1995; Jones  
27 1999; Wilson et al. 1995). Currently, it is assumed that the maintenance of riparian buffers  
28 on headwater streams may protect existing populations by maintaining cool, moist  
29 microclimate conditions and LWD (Nordstrom and Milner 1997; Petranka 1998). Notably,  
30 three of the four nests described by Blessing et al. (1999) were in old-growth forest, and  
31 one was in a 300-foot wide (total width) riparian buffer of old-growth trees adjacent to a  
32 10-year-old logged stand.

#### 33 **3.9.3.2 Dunn's Salamander**

34 Nordstrom and Milner (1997) noted that Dunn's salamanders depend on moist, well-  
35 shaded substrates with stable microclimates. Timber harvest can remove canopy cover that  
36 maintains microclimatic conditions favored by this species, including cool substrate  
37 temperatures and high relative humidity (Chen et al. 1993, 1995; Ledwith 1996; Nordstrom  
38 and Milner 1997). Populations can persist in logged areas, but are more likely to do so  
39 when mature timber is present upstream than when stands upstream have been harvested  
40 (Corn and Bury 1989). Vesely and McComb (2002) found that Dunn's salamanders were  
41 sensitive to forest practices in riparian areas, and concluded that riparian buffers may cause  
42 local declines in abundance. Similarly, West and O'Connell (1998) observed that riparian





1 buffers can encourage persistence of amphibians following timber harvest. Several studies  
2 have demonstrated a direct relationship between buffer width and the maintenance of cool  
3 microclimate and high humidity (Brown and Krygier 1970; Ledwith 1996).

### 4 **3.9.3.3 Torrent Salamanders**

5 Much of the following discussion is based on studies of the southern torrent salamander  
6 (*Rhyacotriton variegatus*). Because all four species are similar enough to have been  
7 considered one species until recently, torrent salamanders of Washington State likely have  
8 similar habitat needs. Where available data differentiate among the species, these  
9 differences are noted in the discussion.

10 Several studies have documented greater abundance of torrent salamanders in old-growth  
11 forests compared to younger stands (Ashton 2002; Corn and Bury 1989; Corn and Bury  
12 1991; Vesely and McComb 2002; Welsh and Lind 1991). Reasons for reduced population  
13 density in logged areas may include loss of optimal microclimate conditions (e.g., cool  
14 temperatures and high relative humidity) due to canopy removal, and reduced habitat  
15 quality due to sedimentation (Nordstrom 1997). Timber harvest and associated road  
16 construction activities may increase the risk of debris torrents, causing scouring and  
17 increase the presence of fine sediments in headwaters and high-gradient streams (Morrison  
18 1975; Swanston and Swanson 1976). Jackson et al. (2003) documented dramatic increases  
19 in the proportion of fine sediments in stream channels following clearcut harvest of  
20 adjacent timber stands, but found no significant differences among treatments in response  
21 by torrent salamanders. The presence of fine sediments may reduce instream habitat  
22 quality for torrent salamanders by filling interstitial spaces critical for movement, egg  
23 deposition, and larval development (Corn and Bury 1989; Diller and Wallace 1996), but  
24 this has yet to be definitively demonstrated.

25 In managed landscapes, torrent salamander abundance appears to be closely tied to  
26 landform characteristics and parent geology. Working in second-growth forests in the  
27 range of the Columbia torrent salamander, Wilkins and Peterson (2000) found the highest  
28 rates of occupancy in high-gradient (greater than 20 percent) channels in small drainage  
29 basins (less than 35 acres). In managed forests of northern California, Diller and Wallace  
30 (1996) found that geology—namely, the presence of consolidated lithologies—was the  
31 only landscape variable that predicted presence of southern torrent salamanders. In  
32 contrast to the studies of second-growth forests, Adams and Bury (2002) studied streams  
33 within unmanaged forests in Olympic National Park, and found that all stream amphibians  
34 were common in waters with unconsolidated surface geology (i.e., exposed, fractured  
35 bedrock, soil, and vegetation overlying solid intact bedrock below).

36 These findings suggest that topography and parent geology may play a role in determining  
37 torrent salamander abundance in unmanaged landscapes, but may substantially mitigate the  
38 negative effects of sedimentation, particularly in areas where logging is practiced. Wilkins  
39 and Peterson (2000) speculated that landform features may have interacted with previous  
40 forest management to influence habitat occupancy. Torrent salamanders may have been  
41 more widespread in the study area prior to logging; fine sediment accumulation may have  
42 depressed salamander populations in low-gradient streams, while faster flushing of fine



## Chapter 3

---

1 sediments may have allowed populations in steep streams to persist or recover more  
2 quickly (Wilkins and Peterson 2000). However, although Wilkins and Peterson found that  
3 sand accumulation decreased with increasing channel gradient in marine sediment streams,  
4 and marine sediment streams had more than double the sand accumulation of basalt  
5 streams, they failed to find a difference in torrent salamander occupancy or abundance  
6 attributable directly to the distinction between these two stream types. This indicates that  
7 the interaction between forest practices and geology remains unclear in relation to  
8 salamander habitat.

9 Vesely and McComb (2002) concluded that riparian buffers may prevent or reduce torrent  
10 salamander population declines following logging, but that current buffer width  
11 requirements in many areas may be inadequate to prevent local declines in amphibian  
12 diversity. Similarly, Jackson et al. (2003) found that the post-harvest proportion of fine  
13 sediments showed little change in headwater streams that were protected by a riparian  
14 buffer. Corn and Bury (1989) also recommended riparian buffers as a means of protecting  
15 the streamside microhabitat variables required by torrent salamanders.

16 Notably, most of the studies that demonstrate negative effects of sedimentation are from  
17 the ranges of the southern species. Streams in the range of the southern torrent salamander  
18 (northwestern California and southwestern Oregon) are prone to carry heavier sediment  
19 loads than streams in the Olympics and Washington Cascades. The differences between  
20 the two areas may be due to the presence of unconsolidated marine sediments (i.e.,  
21 sediments that accumulated on the ocean floor in an unconsolidated form created and  
22 transported by weathering, erosion, volcanoes), heavier rainfall (in some cases), and  
23 warmer climate. Thus, torrent salamanders in Washington may face a lower risk of  
24 negative effects due to logging-related sedimentation than the southern species. This is  
25 corroborated by recent studies of the Columbia torrent salamander in southwestern  
26 Washington (Personal Communication, Marc Hayes, Washington Department of Fish and  
27 Wildlife, October 2003) and northwestern Oregon (Russell et al. 2004), which found this  
28 species not only to be widespread in the timber managed landscape, but at densities higher  
29 than ever reported for a stream-associated salamander. This also may be particularly true  
30 of the Olympic torrent salamander, whose range is characterized by consolidated parent  
31 geology and large Federal reserves.

### 32 **3.9.3.4 Tailed Frogs**

33 Nussbaum et al. (1983) reported that tailed frogs disappeared from streams when areas  
34 were logged, speculating that increased water temperature and siltation were the cause.  
35 Jackson et al. (2003) compared pre- and post-logging populations of tailed frogs at five  
36 streams in southwestern Washington. In the three streams that were clearcut harvested, no  
37 tailed frogs were detected immediately following harvest; 2 years later, tailed frogs were  
38 still absent from two of the three streams. Corn and Bury (1989) and Dupuis and  
39 Steventon (1999) also found that logging had substantially negative effects on densities of  
40 tailed frogs. Such population effects may last for decades. In the redwood forests of  
41 northern California, Ashton (2002) documented substantially greater numbers of tailed  
42 frogs in late-seral forests compared to 40- to 60-year-old second growth stands. Findings  
43 of recent studies have suggested that increased sediment input may be the most important

---

## Chapter 3



1 factor behind tailed frog population declines following logging (Dupuis and Steventon  
2 1999; Ashton 2002). Dupuis and Steventon (1999) also found that buffered creeks had, on  
3 average, higher densities of tailed frogs than logged creeks. Several studies have also  
4 suggested that riparian buffers may be able to protect the streamside microhabitat variables  
5 required by tailed frogs, even if the surrounding habitat is not maintained as old growth  
6 (Bull and Carter 1996; Corn and Bury 1989). In somewhat of a contrast to the above  
7 information, tailed frog tadpoles and adults were found in abundance in several high-  
8 gradient streams surrounded by young (less than 20 years old) riparian and upland stands  
9 southeast of Mount St. Helens in the Gifford-Pinchot National Forest (Personal  
10 Communication, Sally Butts, USFWS, November 17, 2004).

11 In a study of 40 perennial non-fish-bearing streams in southwestern Washington, Wilkins  
12 and Peterson (2000) found tailed frogs only in streams with basaltic (i.e., bedrock)  
13 lithology. Similarly, Jackson et al. (2003) found tailed frogs only at steep basalt sites, and  
14 concluded that local geologic and topographic conditions play a large role in determining  
15 the presence and abundance of this species. In both studies, all surveyed streams occurred  
16 in second-growth forest stands. In contrast, Adams and Bury (2002) studied streams  
17 within unmanaged forests in Olympic National Park, and found that all stream amphibians  
18 were common in waters with unconsolidated surface geology. However, the range of  
19 competence in the unconsolidated surface geology class that Adams and Bury (2002)  
20 compared was broad, making it impossible to exclude the possibility that geology is not the  
21 pivotal control on tailed frog distribution. Welsh and Lind (2002) also found tailed frogs  
22 to be common in streams with unconsolidated geologies in the Klamath-Siskiyou Region,  
23 noting instead that stream temperature and forest age were the strongest predictors of tailed  
24 frog presence and abundance. Collectively, these findings support Dupuis and Steventon's  
25 (1999) report that the competency of the parent geology had substantial effects on tailed  
26 frogs. Timber harvest may exacerbate these effects, but the possibility that geology is the  
27 dominant control has not yet been excluded.

28



## **Chapter 3**

---

1 This page is intentionally left blank.

2



1    **3.10    BIRDS, MAMMALS, OTHER WILDLIFE AND THEIR HABITATS**

2    **3.10.1   Introduction**

3    This section presents a general description of birds, mammals, and other wildlife species in  
4    Washington, including rare, threatened, and endangered species that would be most  
5    affected by the alternatives described in Chapter 2 (Alternatives). Discussions focus on  
6    species with a strong association with riparian habitats, because those species have the  
7    greatest potential to be affected by the alternatives. A brief overview of forested habitat  
8    types available in upland areas is also provided. Changes in timber harvest intensity under  
9    the alternatives may lead to differences in the availability and distribution of forest  
10   structure classes overall. Effects on threatened and endangered species are also addressed.  
11   Table 3-24 lists the species occurring on covered lands that have some special status within  
12   the State. This list is not intended to be a complete list of all species native to Washington;  
13   instead, it is a list of sensitive species or species that have regulatory status under State or  
14   Federal statutes, and that face the potential for significant impacts under the proposed  
15   alternatives.

16   Many Washington amphibian species use streams, ponds, and temporary waters for  
17   mating, egg deposition, and larval development (Nussbaum et al. 1983). Because of their  
18   limited range, limited mobility, and sensitivity to water temperature and quality,  
19   amphibians are particularly sensitive to alterations of riparian and aquatic habitat  
20   (Nussbaum et al. 1983). Several of the amphibian species with special status in  
21   Washington, such as the Oregon spotted frog, have limited distributions and thus may be  
22   more at risk from disturbance than other species (Knutson and Naef 1997).

23   Several groups of birds are closely associated with riparian areas. These include many  
24   neotropical migrants, cavity-nesting birds (i.e., woodpeckers and waterfowl), waterfowl,  
25   and raptors (mainly the bald eagle and osprey). The complexity of riparian vegetation  
26   (subsection 3.6.2, Riparian Vegetation) provides breeding, foraging, and cover habitat for  
27   many of these species (Knutson and Naef 1997).

28   A wide variety of mammals are closely associated with riparian areas. At least five  
29   endemic small mammals are considered obligate inhabitants of streamside areas: water  
30   shrew, marsh shrew, muskrat, beaver, and water vole (O'Connell et al. 1993). The habitat  
31   characteristics of riparian areas, including presence of water, abundance of food, moist  
32   microclimate, and edge habitat support the life requisites of these species and a wide  
33   variety of other mammal species, including river otter, mink, raccoon, black bear, fisher  
34   marten, mule deer, and elk (Knutson and Naef 1997). Timber harvest has the potential to  
35   reduce (and in some cases increase) the populations of these species by affecting cover,  
36   decreasing or increasing the prey base or food sources, and affecting breeding areas.

37




# Chapter 3

1 **Table 3-24. Washington Special Status and High Profile Species on Covered**  
 2 **Lands.**

Common Name	Scientific Name	Status <sup>1/</sup>	Distribution <sup>2/</sup>	Use of Riparian Areas <sup>3/</sup>
<b>Amphibians</b>				
Coastal tailed frog	<i>Ascaphus truei</i>		1-5, 7-9	Stream/Creek – b, f
Western toad	<i>Bufo boreas</i>	SC	1-12	Lake/Pond – b, f
Dunn’s salamander	<i>Plethodon dunnii</i>	SC	2-3	Stream/Creek - b, f
Van Dyke’s salamander	<i>Plethodon vandykei</i>	SC, FSC	1-5	Stream/Creek - b, f
Northern red-legged frog	<i>Rana aurora</i>	FSC	1-7, 9	Lake/Pond/Slough - b, f
Cascades frog	<i>Rana cascadae</i>	FSC	1, 3-5, 7-9	Lake/Pond/Stream - b, f
Northern leopard frog	<i>Rana pipiens</i>	SE	8-12	Lake/Pond – b, f
Oregon spotted frog	<i>Rana pretiosa</i>	SE, FC	4, 7, 9	Lake/Pond – b, f
Columbia spotted frog	<i>Rana luteiventris</i>	SC, FSC/FC	7-12	Lake/Pond – b, f
Cascade torrent salamander	<i>Rhyacotriton cascadae</i>	SC	3-4	Stream/Creek - b, f
Columbia torrent salamander	<i>Rhyacotriton kezeri</i>	SC, FSC	2-3	Stream/Creek - b, f
Olympic torrent salamander	<i>Rhyacotriton olympicus</i>	FSC	1-2, 5	Stream/Creek – b, f
<b>Reptiles</b>				
Western pond turtle	<i>Clemmys marmorata</i>	SE, FSC	3-5, 9	Lake/Slough/Stream - f
Sharptail snake	<i>Contia tenuis</i>	SC	4, 8-9	Wetlands - b, f
<b>Birds</b>				
Wood duck	<i>Aix sponsa</i>	P	1-12	River/Stream – b, f
Great blue heron	<i>Ardea herodias</i>	P	1-12	Stream/Wetlands – b, f
Marbled murrelet	<i>Brachyramphus marmoratus marmoratus</i>	ST, FT	1-5, 7	River/Stream – b
Pileated woodpecker	<i>Dryocopus pileatus</i>	SC	1-12	River/Stream – b, f
Northern spotted owl	<i>Strix occidentalis caurina</i>	SE, FT	1-5, 7-9	River/Stream – b, f
Willow flycatcher	<i>Empidonax traillii</i>	FSC	1-12	N/A in WA
Common loon	<i>Gavia immer</i>	SS	1, 4, 7-8, 10	Lake - b, f
Sandhill crane	<i>Grus canadensis</i>	SE	9	Wetlands - b, f
Bald eagle	<i>Haliaeetus leucocephalus</i>	ST, FT	1-11	River/Lake – f
Harlequin duck	<i>Histrionicus histrionicus</i>	P, FSC	1, 3-5, 7-10	River/Stream - b, f
<b>Butterflies</b>				
Oregon silverspot butterfly	<i>Speyeria zerene hippolyta</i>	SE, FT	2	N/A
Mardon skipper	<i>Polites mardon</i>	SE, FC	2, 4, 9	N/A
<b>Mammals</b>				
Beaver	<i>Castor canadensis</i>	HP	1-5, 7-12	Stream/Creek – b, f
River Otter	<i>Lutra canadensis</i>	HP	1-12	River/Stream – b, f
Mink	<i>Mustela vison</i>	P	1-12	River/Stream – b, f

3



1 **Table 3-24.** Washington Special Status and High Profile Species on Covered  
2 Lands (continued).

Common Name	Scientific Name	Status <sup>1/</sup>	Distribution <sup>2/</sup>	Use of Riparian Areas <sup>3/</sup>
<b>Mammals (continued)</b>				
Muskrat	<i>Ondatra zibethicus</i>	HP	1-12	Stream/Wetlands – b, f
Western gray squirrel	<i>Sciurus griseus</i>	ST,FSC	2, 6, 7, 8, 11	River/Stream/Creek – b, f
Mazama pocket gopher	<i>Thomomys mazama</i>	SC, FC	2, 4, 5	N/A
Canada lynx	<i>Lynx canadensis</i>	ST, FT	7, 8, 10	N/A
Gray wolf	<i>Canis lupus</i>	SE, FT	Unknown	N/A
Columbian white-tailed deer	<i>Odocoileus virginianus leucurus</i>	SE, FE	3	Stream/Slough – b, f
Woodland caribou	<i>Rangifer tarandus caribou</i>	SE, FE	10	N/A
Grizzly bear	<i>Ursus arctos horribilis</i>	SE, FT	7, 8, 10	River/Stream/Wetlands – f

<sup>1/</sup> SE = State Endangered; ST = State Threatened; SS = State Sensitive; SC = State Candidate; FE = Federal Endangered; FT = Federal Threatened; FC = Federal Candidate; FSC = Federal Species of Concern; P = Covered Species with WDFW, but not listed; HP = high profile/high public interest.

<sup>2/</sup> Numbers indicate EIS-designated Regions: 1 = Olympic Coast; 2 = Southwest; 3 = Lower Columbia; 4 = Sound Puget Sound; 5 = West Puget Sound; 6 = Islands; 7 = North Puget Sound; 8 = Upper Columbia – Downstream of Grand Coulee; 9 = Middle Columbia; 10 = Upper Columbia – Upstream of Grand Coulee; 11 = Columbia Basin; and 12 = Snake River.

<sup>3/</sup> Indicates type of riparian area used, and type of use (b = breeding; f = foraging), based on Brown 1985; Riparian habitat not commonly used = N/A.

Sources: Brown 1985; WDFW Web site (<http://wdfw.wa.gov/wlm/diversity/soc/soc.htm>).

### 3 3.10.2 Federally Threatened or Endangered Wildlife Species

#### 4 3.10.2.1 Marbled Murrelet

5 The marbled murrelet (*Brachyramphus marmoratus marmoratus*) was listed as threatened  
6 under the ESA in 1992. Although marbled murrelets feed primarily on fish and  
7 invertebrates in nearshore marine waters, they fly inland to nest on large limbs of mature  
8 conifers (USFWS 1997). The main cause of population decline and the primary threat to  
9 the bird in Washington was the loss and alteration of nesting habitat (older forests)  
10 primarily as a result of timber harvesting (USFWS 1997). Other threats are mortality  
11 associated with gill-net fishing, nest predation, oil spills, and stochastic disturbance events  
12 that result in a loss of nesting habitat (USFWS 1997). In Washington, the murrelet is  
13 found in all nearshore marine areas with the greatest concentrations in northern Puget  
14 Sound (Table 3-24). Nesting behavior has been documented beyond 50 miles inland,  
15 though most nesting habitat occurs within 50 miles of shore throughout the breeding range  
16 (USFWS 1997). The majority of marbled murrelet sightings have been within 39 miles of  
17 the coast in the north Cascades. The most recent estimates of marbled murrelet numbers in  
18 Washington are 10,600 birds (6,300 to 15,700 birds – 95 percent confidence interval) in  
19 2001 and 12,300 birds (6,800 to 17,600 birds – 95 percent confidence interval) in 2002  
20 (Huff 2002).



## Chapter 3

---

### 1 3.10.2.2 Northern Spotted Owl

2 The northern spotted owl (*Strix occidentalis caurina*) was listed as threatened under the  
3 ESA in 1990. Its range in Washington encompasses the coastal mountains (including the  
4 Olympics and Willapa Hills) and Cascade Range (both western and eastern Washington).  
5 Preferred habitat includes structurally complex mature and old-growth coniferous forests  
6 with moderate to high canopy closure, a multi-layered, multi-species canopy with large  
7 overstory trees, a high incidence of snags or large trees with deformities, large  
8 accumulations of fallen trees and other debris, and a well developed shrub layer (Thomas  
9 et al. 1990). The owl's favored prey is the northern flying squirrel (*Glaucomys sabrinus*),  
10 although the birds also feed on a variety of other small mammals.

11 Declines in the number of spotted owls have been documented on long-term study areas in  
12 Washington State (Forsman et al. 1999). Preliminary results from the last northern spotted  
13 owl demography workshop held in January 2004 (Anthony et al. 2004) concluded that  
14 northern spotted owl populations on many of the study areas have declined more rapidly  
15 than was observed in previous reports (Franklin et al. 1999). The demographic analysis  
16 suggests that the range-wide northern spotted owl population declined at about 4.1 percent  
17 per year during 1990 to 2003. The average estimated rate of decline for eight monitoring  
18 areas under the effectiveness monitoring plan was 2.5 percent per year. The largest  
19 declines occurred in Washington State (7.5 percent per year). Oregon declined by 2.8  
20 percent per year, and California declined by 2.2 percent per year. According to the 2004  
21 report, the populations in the Cle Elum, Wenatchee, and Mount Rainier study areas  
22 declined substantially over the last decade. The population sizes were approximately 40 to  
23 60 percent of initial populations in the Cle Elum and Wenatchee study areas. The Olympic  
24 Peninsula population in 2002 was approximately 70 to 80 percent of initial populations.  
25 "Initial populations" here refers to the populations at the time the demography studies  
26 began, which for these areas is late 1980s and early 1990s.

27 As noted above, spotted owl populations have been declining in the past decade. The  
28 major reasons for the decline are thought to include continuing loss of habitat, climatic  
29 conditions, and the barred owl (*Strix varia*) invasion (Forsman et al. 2002; Forsman et al.  
30 2003). Habitat losses can be exacerbated by catastrophic events such as fire, volcanic  
31 eruptions, and wind storms. Recent studies have documented a rapid increase in the  
32 number of barred owl detections within known spotted owl territories throughout the  
33 Pacific Northwest (Gremel 2001 and 2003; Forsman et al. 2003; Kelly et al. 2003; Lint et  
34 al. 2003; and Pearson and Livezey 2003). The major concern from this increase in barred  
35 owl detections is that barred owls may outcompete spotted owls for limited resources.  
36 According to Pearson and Livezey (2003), spotted owl site occupancy appeared to be more  
37 affected by the presence of barred owls than by land management allocations, but suspect  
38 that human-caused loss of old-growth forest might reduce the ability of spotted owls to  
39 compete successfully with barred owls.

40 The USFWS designated spotted owl critical habitat solely on Federal lands (United States  
41 Department of the Interior 1992b) and anticipated that the major burden of conservation  
42 and recovery of northern spotted owl populations would be carried by these lands. The





1 final draft Recovery Plan for the Northern spotted owl (United States Department of the  
2 Interior 1992a) recommended establishment of conservation areas on Federal lands as the  
3 primary means for achieving recovery of the spotted owl. Hanson et al. (1993) identified  
4 important non-Federal landscapes for “essential owl habitats” on non-Federal lands, and  
5 provided recommendations for site and landscape-specific plans.

6 In 1996, the Washington Forest Practices Board conducted an evaluation of alternatives for  
7 providing additional protection for the spotted owl. The Forest Practices Board’s intent  
8 was to “define a level of conservation contribution from non-Federal lands that is essential  
9 to complement the Federal recovery and conservation strategy; identify those landscapes  
10 that are essential to complement the Federal conservation and recovery strategy; maximize  
11 the use of local planning to promote flexibility, and minimize conflicts and economic  
12 impacts” (Washington DNR 1996).

13 Final rules were adopted in July 1996 partly based on the USFWS proposed ESA Section  
14 4(d) rule for spotted owls on nonfederal lands and partly based on the Hanson et al. (1993)  
15 report on essential owl habitat on non-Federal lands. The rules designated 10 spotted owl  
16 special emphasis areas (SOSEAS) to provide for demographic support, dispersal support or  
17 a combination of both. Timber harvest, road construction, and aerial application of  
18 pesticides on suitable owl habitat inside owl circles within the SOSEAS would be Class  
19 IV-Special forest practices (with the exception of the Entiat SOSEA), and must comply  
20 with SEPA. Within the Entiat SOSEA, these SEPA requirements only apply to  
21 demographic support areas. The Forest Practices Board’s goals in adopting SOSEAs was  
22 to “maintain owls where they can make a contribution to the species, not maintain all  
23 individual owls where they currently exist; and to allow strategic allocation of habitat to  
24 those owls that have the potential to contribute to the viability of the species” (Washington  
25 DNR 1996). The Forest Practices Board recently requested that the WDFW conduct a  
26 review of the wildlife rules and report back to the Forest Practices Board by November  
27 2004. Consistent with WAC 222-16-080, part of this review will be to “determine whether  
28 circumstances exist that substantially interfere with meeting the goals of the SOSEAs.”

### 29 **3.10.2.3 Bald Eagle**

30 The bald eagle (*Haliaeetus leucocephalus*) was listed as threatened under the ESA in 1967.  
31 In 1998, there were 664 occupied nest sites in the State with some indications that the  
32 population has reached its carrying capacity in parts of western Washington (Washington  
33 Department of Fish and Wildlife 2001b). Winter populations are higher (3,500 to 4,000  
34 birds) due to an influx of migrants from Alaska and the Canadian provinces.

35 Breeding bald eagles need large trees near open water that are not subject to intense human  
36 activity. In Washington, nearly all bald eagle nests (99 percent) are within 1 mile of a lake,  
37 river, or marine shoreline (Washington Department of Fish and Wildlife 2001b). The  
38 distance to open water varies somewhat with shore type. Nests tend to be closer to marine  
39 shores and rivers than to lake shores (mean 457 feet [marine] versus 633 feet [river] versus  
40 997 feet [lakes]) (Washington Department of Fish and Wildlife 2001b). Assuming the  
41 presence of an adequate food supply, the single most critical habitat factor associated with  
42 eagle nest locations and success is the presence of large super-dominant trees (Watson and



## Chapter 3

---

1 Pierce 1998). Although current Washington Forest Practices Rules require large trees  
2 within riparian areas to be retained, the majority of forested riparian areas in western  
3 Washington are in an early seral stage with only 2 percent estimated in late seral stage (See  
4 subsection 3.7, Riparian and Wetland Processes). Bald eagles are not old-growth  
5 obligates, but use large trees capable of supporting their massive nests.

6 Past impacts to bald eagle populations include poaching, timber harvesting, reduced  
7 salmon runs, and the use of the pesticide DDT. The greatest current threat to eagle  
8 populations in Washington is the loss of suitable nesting habitat. Eagles prefer to nest in  
9 large trees along shorelines, but will utilize smaller second-growth trees (Washington  
10 Department of Fish and Wildlife 2001b). Conservation of bald eagle nesting habitat is  
11 difficult because 80 percent of the land within one-half mile of shorelines is privately  
12 owned and contains desirable view property subject to development. The State bald eagle  
13 protection rule (WAC 232-12-292) requires a management plan for development, forest  
14 practices, or other potentially disturbing activities on State and private lands near eagle  
15 nests and roosts (Washington Department of Fish and Wildlife 2001b).

16 The State's bald eagle protection rules of 1986 (WAC 232-12-292) established a legal  
17 requirement for private, State, and municipal landowners to reach agreement with WDFW  
18 on measures to protect breeding and roosting habitat. These rules are the most important  
19 mechanism for the protection of habitat on private and State lands in Washington. The  
20 retention of future nest and perch trees, in addition to currently used trees, has probably  
21 been an important contribution to eagle habitat from the regulation. The amount of  
22 privately owned, but undeveloped lands near shore (much already subdivided), indicate  
23 that the need for planning will continue to be important for some time to provide habitat  
24 for eagles.

### 25 **3.10.2.4 Oregon Silverspot Butterfly**

26 The Oregon silverspot butterfly (*Speyeria zerene hippolyta*) was listed as threatened under  
27 the ESA in 1980. In Washington, the species is found on the Long Beach peninsula in the  
28 Southwest Region (Table 3-24). The butterfly occupies early successional, coastally-  
29 influenced grassland habitats that contain the caterpillar host plant early blue violet (*Viola*  
30 *adunca*), adult nectar sources, and adult courtship areas (USFWS 2001). Soil and climatic  
31 conditions, salt-spray or mist, and natural disturbances such as fire historically contributed  
32 to maintaining low, open grasslands within the species' range by suppressing encroaching  
33 trees and shrubs. Invasion by exotic species, natural succession, fire suppression, and land  
34 development have resulted in habitat loss and modification and represent the primary  
35 threats to the species (USFWS 2001).

### 36 **3.10.2.5 Canada Lynx**

37 The Canada lynx (*Lynx canadensis*) was listed as threatened under the ESA in 2000. There  
38 are currently thought to be fewer than 100 individuals in the State (Washington  
39 Department of Fish and Wildlife 2001a). In Washington, lynx are primarily associated  
40 with subalpine and boreal forest types in the north-central and northeastern parts of the  
41 State. These areas lie within the North Puget and Upper Columbia Regions (Figure 3-1).



1 The WDFW originally identified six Lynx Management Zones that represented the  
2 distribution of primary lynx habitat in Washington (Brittell et al. 1989 in Washington  
3 Department of Fish and Wildlife 2001a). Boundaries were initially drawn based on the  
4 4,000-foot (1,220 meter) elevation contour and were refined based on the knowledge of  
5 biologists involved in lynx surveys. Boundaries were modified in 2000 by the Colville,  
6 Idaho Panhandle, and Wenatchee National Forests using a more refined definition of  
7 habitat that included elevations down to 3,500 feet in northeastern Washington, and took  
8 into account local detections of lynx, and deleted some areas of permanent non-lynx  
9 habitats (dry pine, openings) along the Lynx Management Zone periphery. These  
10 management zones do not encompass all areas potentially used by lynx, but habitat  
11 management within these zones is expected to hold the greatest promise for supporting  
12 lynx populations.

13 In contrast to the habitat of many rare species, potential lynx habitat has not been  
14 developed or converted to agriculture, but most is still forested and the potential to manage  
15 it for lynx still exists. Federal land jurisdictions include about 92 percent of the habitat in  
16 the six Lynx Management Zones, with the vast majority (approximately 88 percent)  
17 administered by the U.S. Forest Service (Washington Department of Fish and Wildlife  
18 2001a). The National Park Service (North Cascades National Park) administers about 3.6  
19 percent of primary lynx habitat across Washington. Reserve land type designations (e.g.,  
20 wilderness areas, refuges) protect almost 40 percent of the lynx habitat in Washington  
21 (Washington Department of Fish and Wildlife 2001a).

22 Lynx are largely dependent on a single prey species, the snowshoe hare (*Lepus*  
23 *americanus*), but also eat small mammals, birds, and carrion. The primary factors affecting  
24 populations in Washington include forest management, fire and fire suppression, insect  
25 epidemics, and management of lynx harvest and habitats in southern British Columbia  
26 (Washington Department of Fish and Wildlife 2001a).

27 The Washington Forest Practices Board, which has regulatory authority over timber  
28 harvest on State and private lands, designates critical wildlife habitat for State-listed  
29 species that may be affected by forest practices (WAC 222-16-010). Forest practices  
30 regulations, however, allow landowners to prepare special wildlife management plans in  
31 lieu of being subject to a critical habitat rule (WAC 222-16-080(2)). The three major non-  
32 Federal landowners in the Washington lynx range are the Washington DNR, Boise  
33 Cascade Corporation, and Plum Creek Timber Company; each of these has developed lynx  
34 plans, which were subsequently approved by the WDFW in 1996. Based on the approval  
35 of the above plans, the Forest Practices Board determined that no critical habitat rule would  
36 be needed for lynx since all of the significant State and private land in lynx range were  
37 covered by the three landowner plans.

### 38 **3.10.2.6 Gray Wolf**

39 The gray wolf (*Canis lupus*) is currently listed as threatened under the ESA; they were  
40 originally listed as endangered in 1967. Although there have been occasional reports of  
41 individual wolves in Washington, no documented breeding pairs or packs currently are  
42 known to occur in the State (Washington Department of Fish and Wildlife 1999b) (Table



## Chapter 3

---

1 3.24). However, wolves may appear in Washington within the next few years as they  
2 disperse from sites where they have recently been reintroduced in central Idaho, Wyoming,  
3 and Montana. The USFWS has proposed to de-list the Western Distinct Population  
4 Segment (DPS) of the gray wolf once all three states have approved wolf management  
5 plans. Wolves are highly adaptable and can survive in a variety of habitats, although they  
6 prefer relatively flat, open areas such as river valleys and basins (Washington Department  
7 of Fish and Wildlife 1999b). Primary prey species for wolves include elk, deer, and  
8 moose.

9 Rendezvous sites are usually near water and are characterized by systems of trails, beds,  
10 and play areas; they often border meadows (Washington Department of Fish and Wildlife  
11 2003). The current Washington Forest Practices Rules provide timing restrictions on  
12 harvest, road construction, and site preparations within at least 0.25 mile of a documented  
13 den site throughout the year and up to 1 mile from an active denning area (WAC 222-16-  
14 080).

### 15 **3.10.2.7 Columbian White-tailed Deer**

16 The Columbian white-tailed deer (*Odocoileus virginianus leucurus*) exists in two distinct  
17 population segments. The first is found along the lower Columbia River in southwest  
18 Washington and northwest Oregon; the second is found along the Umpqua River in  
19 Douglas County, Oregon. Both populations were listed as endangered under the Federal  
20 ESA in 1967. In 2003, the USFWS de-listed the Umpqua/Douglas County population,  
21 which is now estimated at several thousand animals.

22 In Washington State, the Columbian white-tailed deer is only known to occur in  
23 Wahkiakum County on islands in and along the banks of the Columbia River. The islands  
24 are located within the Columbia River and are in private ownership or are managed by the  
25 USFWS, Washington DNR, and WDFW. These lands lie within the Lower Columbia  
26 Region (Figure 3-1).

27 The Lower Columbia distinct population segment numbers about 700 animals and is  
28 distributed throughout the Julia Butler Hansen Columbian White-tailed Deer National  
29 Wildlife Refuge; on Puget, Crims, Lord, and Fisher Islands; and on the Oregon mainland.  
30 The islands are located within the Columbia River and are in private ownership or are  
31 managed by the USFWS, Washington DNR, and WDFW. These lands lie within the  
32 Lower Columbia Region (Table 3-24).

33 Columbian white-tailed deer prefer wet prairie and lightly wooded bottomlands or  
34 tidelands along streams and rivers. Woodlands are particularly attractive when  
35 interspersed with grasslands and pastures (NatureServe 2003). Major tree species along  
36 the Columbia River include Sitka spruce, dogwood, black cottonwood, red alder, and  
37 willow. The primary factors affecting the Lower Columbia population are land  
38 conversion, timber harvesting, vehicular traffic, poaching, and flooding (NatureServe  
39 2003).



### 1 **3.10.2.8 Woodland Caribou**

2 The woodland caribou (*Rangifer tarandus caribou*) was listed as endangered under the  
3 ESA in 1983. Currently, the population includes approximately 50 animals in the Selkirk  
4 Mountains of northeastern Washington, northern Idaho, and southeastern British  
5 Columbia, occurring as two herds (USFWS 1994). This area lies within the Upper  
6 Columbia-Upstream from Grand Coulee Dam Region (Table 3.24). The animals are  
7 generally found above 4,000 feet in elevation in Engelmann spruce/subalpine fir and  
8 western redcedar/western hemlock forest types (USFWS 1994). Woodland caribou feed  
9 almost exclusively on tree-borne lichens. The population is threatened by habitat  
10 fragmentation and loss and excessive mortality (USFWS 1994).

11 The recovery area for caribou in the Selkirk Mountains is comprised of approximately  
12 2,200 square miles in northern Idaho, northeastern Washington, and southern British  
13 Columbia. About 47 percent of the area lies in British Columbia, and 53 percent lies in the  
14 United States. The United States portion includes the Salmon-Priest Wilderness and other  
15 portions of the Colville and Idaho Panhandle National Forests, Idaho Department of Lands  
16 holdings, and scattered private parcels (USFWS 1994).

17 The current Washington Forest Practices Rules provide timing restrictions on harvesting,  
18 road construction, aerial application of pesticides, and site preparations within at least 0.25  
19 mile of a documented breeding area (WAC 222-16-080).

### 20 **3.10.2.9 Grizzly Bear**

21 The grizzly bear (*Ursus arctos horribilis*) was listed as threatened under the ESA in 1975.  
22 Grizzly bears are rare in Washington, but there is a small population in the Selkirk  
23 Mountains (Upper Columbia Upstream of Grand Coulee Dam Region) of northeast  
24 Washington. Grizzly bears have also been documented in the Okanogan Highlands and in  
25 the North Cascades (North Puget and Upper Columbia Downstream of Grand Coulee Dam  
26 Regions). Contiguous, relatively undisturbed mountainous habitat with a high level of  
27 topographic and vegetative diversity is characteristic of most areas where the species exists  
28 (USFWS 1993). Direct and indirect human-caused mortality and habitat loss have caused  
29 the decline in bear numbers (USFWS 1993).

30 Portions of two grizzly bear recovery zones exist in Washington State: the Selkirk and  
31 Cabinet Yaak Grizzly Bear Recovery Zone and the North Cascades Grizzly Bear Recovery  
32 Zone. The North Cascades recovery area covers almost 10,000 square miles (one of the  
33 largest in the United States). More than 40 percent of the recovery area is designated  
34 wilderness, 90 percent is Federal or State owned, and 68 percent has no motorized access.

35 The current Washington Forest Practices Rules provide timing restrictions for harvesting,  
36 road construction, aerial application of pesticides, or site preparation within 1 mile of a  
37 known active den site, documented by the WDFW, between the dates of October 1 and  
38 May 30, or within 0.25 mile at other times of the year (WAC 222-16-080(1)(c)).



## Chapter 3

---

### 3.10.3 Importance of Riparian Habitats to Wildlife

Riparian areas are among the most important wildlife habitats in Washington. A variety of physical and biotic features unique to the terrestrial-aquatic interface contribute to this high degree of biological diversity. Out of the 593 wildlife species that occur in Oregon and Washington, 319 (53 percent) use riparian zones (Johnson and O'Neil 2001). O'Connell et al. (1993) and Oakley et al. (1985) provide extensive reviews of the literature on wildlife use of riparian areas. This section highlights several attributes of riparian habitat that are of particular importance to riparian-dependent species. These include complex vegetation structure, snags and downed woody debris, edge effect, and connectivity.

#### 3.10.3.1 Complex Vegetation Structure

Riparian zones are noted for their structural complexity. They often are characterized by a variety of vegetation layers, including herbaceous, shrub, sapling, tree, and overstory layers (Oakley et al. 1985). This floristic diversity is encouraged by the frequent disturbance in most riparian areas, particularly along larger streams, due to flood events, mass wasting events, fire, and windthrow (Wissmar et al. 1994; Agee 1994). A high degree of vegetative structures in a riparian zone can provide abundant sites for breeding, roosting, foraging, and hiding for numerous species. In particular, riparian vegetation structure has been correlated to an abundance of breeding songbirds (Sanders and Edge 1998; Knopf 1985; Martin 1988; Hagar 1999). Doyle (1990) and McComb et al. (1993) reported that structural diversity of riparian vegetation was important to small mammals. However, narrow riparian zones along small streams often do not provide structural diversity enhancement beyond that provided by adjacent upland areas.

#### 3.10.3.2 Snags and Downed Woody Debris

Snags and downed woody debris serve very important biological functions for a wide variety of species. Many birds and small mammals use cavities in snags for nesting and resting. Brown (1985) estimates that over 100 species of wildlife in Oregon and Washington use snags, with approximately 53 of them being cavity-dependent. These species include woodpeckers, cavity-nesting ducks, owls, bats, and most mustelids. Marten and fisher use cavities in live and dead trees as nest sites (Ruggiero et al. 1994). Snags and downed woody debris provide other important habitat functions, including foraging, roosting, and perching habitat. Wildlife will use a wide variety of trees in different stages of decay, including trees with heart rot, hollow trees, completely dead snags, and downed logs of all decay classes (Bull et al. 1997). For instance, Bull et al. (1997) found that pileated woodpeckers in the Blue Mountains of Oregon selectively roosted in live and dead grand firs that were extensively decayed by Indian paint fungus. In the same region, downed logs provide important habitat for forest-dwelling ants, which are a primary prey of pileated woodpeckers (Torgersen and Bull 1995). Similarly, density of cavity-nesting birds in other regions has been positively correlated with the density of large snags (Raphael 1980; Madsen 1985). Marten use large downed logs for predator avoidance, thermal protection, and natal dens (Buskirk and Ruggiero 1994).

Timber harvesting has been shown to reduce the density of snags in the landscape and this has been correlated with reduced abundance of cavity-nesting species (Dickson et al. 1983;



1 Brown 1985; O’Connell et al. 1993). Retention of riparian buffer zones has the potential to  
2 maintain greater densities of snags and downed logs in the landscape. Environmental  
3 conditions in riparian and wetland areas can contribute to the production of snags and  
4 downed logs. Undercut slopes, soil saturation, ponding, high water, and other types of soil  
5 disturbance that are common in riparian areas can all contribute to the weakening of trees  
6 and subsequent production of snags or deformities. Furthermore, riparian buffer zones that  
7 border clearcuts are very vulnerable to windthrow. One study of 40 buffers on small  
8 streams in northwest Washington found that an average of 33 percent of all trees in the  
9 buffers were affected by windthrow (Grizzel and Wolff 1998). This windthrow increased  
10 the large instream woody debris counts in this study by 52 percent compared to counts at  
11 the time of harvest (1 to 3 years earlier). This study concluded that windthrow may be the  
12 most important mechanism for LWD recruitment to stream channels. However, these  
13 authors caution that much of this LWD is suspended over narrow, confined channels and  
14 does not contribute to sediment retention (Grizzel and Wolff 1998). Partially submerged  
15 snags in wetlands, particularly beaver ponds, are important habitat for species such as  
16 cavity-nesting ducks, tree swallows, woodpeckers, and osprey (Knutson and Naef 1997).

17 Windthrow is not the only mechanism that can reduce the amount of snags in a riparian  
18 zone. Some snags in a given riparian zone will have to be removed prior to and during  
19 adjacent timber harvest activities to meet State safety regulations. According to chapter  
20 296-54 of the WAC, any tree that presents a hazard to workers because of some observable  
21 natural or manmade defect is labeled a “danger tree” and must be removed.

### 22 **3.10.3.3 Edge Effect**

23 The edge effect is a term used to describe the potentially positive and negative effects  
24 associated with the change between two different habitat types. These effects can include  
25 increased exposure to predation, increased prey availability, increased vegetative structural  
26 complexity, and increased exposure to light and heat. It is generally used in reference to  
27 the ecotone between recently harvested areas and older forests, but it can also be applied to  
28 the ecotone between riparian areas and upland habitats. Riparian areas, due to their usually  
29 long and sinuous shape, are dominated by edge habitat. Edge habitat is characterized by  
30 the presence of species representative of both the riparian zone and the adjacent upland  
31 habitat. The diverse vegetation and complex structure that characterizes the edge of  
32 riparian zones makes this area attractive and beneficial to many species, particularly  
33 generalist species (Knutson and Naef 1997; Wilcove et al. 1986). These species benefit  
34 from the myriad of different nesting and perching substrates as well as multiple vegetation  
35 layers (e.g., grass, herb, shrub, tree) and usually more abundant food sources such as  
36 berries or insects (Knutson and Naef 1997). Species richness is thus often greatest in edge  
37 habitat (Fraver 1994).

38 On the other hand, some studies have demonstrated the negative effects of edge habitat on  
39 species that are adapted to the conditions of forest away from the edge (i.e., interior  
40 habitat). Increased edge habitat can increase exposure to predators such as crows and  
41 ravens, brown-headed cowbirds, and raccoons. A literature review by Paton (1994)  
42 suggested that predation and parasitism rates on forest interior-nesting birds are often



## Chapter 3

---

1 substantially greater within 164 feet (50 meters) of an edge than for nests found further  
2 from an edge. Nelson and Hamer (1995) found that successful marbled murrelet nests  
3 were located substantially farther from edges (greater than 180 feet [56 meters]) than  
4 unsuccessful nests. The effects of predation have been shown to extend up to 2,000 feet  
5 (620 meters) into a stand (Wilcove et al. 1986).

### 6 **3.10.3.4 Connectivity**

7 Riparian areas can provide important habitat linkages in the landscape. Many different  
8 species have been documented using riparian areas for travel and dispersal (Lovejoy et al.  
9 1986; Brown et al. 1985; Gibbs 1998; Harris 1984). Although very few species are limited  
10 to riparian corridors for movement, many mobile species such as marten, fisher, cougar,  
11 deer, and birds will utilize riparian corridors. Beier (1993) documented cougars in the  
12 Santa Ana Mountains of southern California using relatively narrow riparian corridors for  
13 movement. Machtans et al. (1996) found that forest birds would utilize habitat corridors  
14 more often than clearcuts. The potential value of riparian corridors increases in a  
15 fragmented landscape as they become the only safe way for some species to cross  
16 unsuitable habitat, which is the case for the cougars in the Beier (1993) study.

### 17 **3.10.4 Wildlife in Upland Forested Habitats**

#### 18 **3.10.4.1 Early Seral Forests**

19 Early seral forests represent the initial phases of forest development following a major  
20 disturbance such as a fire or regeneration harvest. They correspond to the grass/forb and  
21 shrub/sapling forest structure classes. Young forest stands with an open canopy and  
22 plentiful shrub cover support a diverse assemblage of small bird species—bird species  
23 diversity and overall abundance reach their highest levels in stands in the ecosystem  
24 initiation stage (Carey et al. 1996). Many of these species are habitat generalists, and most  
25 are very common. Such stands also provide abundant forage for wide-ranging species such  
26 as deer and elk. Other species closely associated with forests in the ecosystem initiation  
27 stage include the dark-eyed junco, white crowned sparrow, and Townsend's vole (Johnson  
28 and O'Neil 2001). Structural legacies (e.g., large snags and down logs) retained from the  
29 previous stand can increase biological diversity by providing habitat for small mammals,  
30 cavity-nesting birds, and terrestrial amphibians (Carey et al. 1996). In managed  
31 landscapes, retention of such legacies combined with a management program designed to  
32 promote biological diversity may speed the development of more complex forest  
33 ecosystems (Carey and Curtis 1996; Carey et al. 1996; Carey 1998).

#### 34 **3.10.4.2 Mid-Seral Forests**

35 Forests of the mid-seral stages generally have a single, dense canopy layer dominated by  
36 trees between 10 and 19 inches in dbh (Johnson and O'Neil 2001). Small snags and  
37 downed logs are often present as smaller trees die because they are suppressed by other  
38 trees competing for available resources called competitive exclusion. Large decaying logs  
39 and stumps may be present as remnants of previous disturbances, such as windstorms or  
40 harvests. Forest structure classes that make up this habitat type include the closed-canopy  
41 shrub/sapling class, all pole-sized classes, and all large-tree classes described by Johnson





1 and O’Neil (2001), except for multistoried, large-tree stands with less than 70 percent  
2 canopy cover.

3 In younger competitive exclusion stands, the high density and uniform size of relatively  
4 short trees allows only small amounts of sunlight to reach the forest floor, creating sparse  
5 understory conditions and low levels of biological diversity. Canopy gaps—either as a  
6 result of thinning or natural mortality—allow understory plants to become established.  
7 The result is a gradual increase in biological diversity. The competitive exclusion stages  
8 have the lowest biodiversity and the least favorable conditions for wildlife when compared  
9 to all the forest stages described by Carey et al. (1996). No wildlife species in western  
10 Washington are found exclusively in competitive-exclusion forests (Carey and Curtis  
11 1996).


### 12 **3.10.4.3 Late Seral Forests**

13 Late seral forests typically feature multiple canopy layers, with the top layer dominated by  
14 trees 20 to 30 inches in dbh (Johnson and O’Neil 2001). However, tree sizes associated  
15 with these forests are quite variable. Snags and down logs play a vital role in providing  
16 structural and biological diversity.

17 Biological diversity in this forest habitat type is promoted by structural complexity along  
18 both the vertical axis (i.e., trees of different heights, as well as shrubs and herbaceous  
19 plants) and the horizontal axis (e.g., gaps in the forest canopy) (Carey et al. 1996; Franklin  
20 et al. 2002). A diversity of plant species and growth forms in structurally complex forest  
21 provides niches for a wide variety of wildlife species. For example, structurally complex  
22 forests have an understory of small trees, shrubs, ferns, and herbs, providing foraging  
23 opportunities for herbivores and breeding habitat for ground-nesting birds (Carey et al.  
24 1996). Large snags and down logs in the more fully developed stages of this class may  
25 provide suitable habitat conditions for a variety of important species, including nest sites  
26 for spotted owls, roost sites for bats, and den sites for Pacific fishers. Very large trees may  
27 also provide nest sites for other wildlife species, including bald eagles and marbled  
28 murrelets.

29

30



## **Chapter 3**

---

1 This page is intentionally left blank.

2



1    **3.11    RECREATION**

2    **3.11.1    Introduction**

3    Public lands account for approximately 40 percent (17.2 million acres) of the land base in  
4    Washington State, of which, approximately 74 percent is federally managed, 22 percent is  
5    managed by the State, and local governments manage the remaining 4 percent (Interagency  
6    Committee for Outdoor Recreation [IAC] 2001) (Table 3-1). Typically laws, policies, or  
7    regulations determine the way these lands are to be managed. Few public lands in  
8    Washington are managed for recreational uses alone; most are managed for multiple uses.  
9    However, for purposes of classification, the IAC (2001) breaks these uses into the  
10   following four general categories (Table 3-25):

- 11   •   Outdoor Recreation, Habitat, or Environmental Protection—Lands with designations  
12       such as camping, picnicking, wildlife areas, wilderness areas, municipal watershed  
13       areas, or natural areas preserves.
- 14   •   Resource Production or Extraction—Lands primarily used for production or extraction  
15       of natural resources such as agricultural, timber, mineral, or fish and wildlife  
16       commodities.
- 17   •   Transportation or Utilities Infrastructure—Lands that support general services to the  
18       public such as roads, utility corridors, power plants, dams, landfills, and sewage  
19       treatment plants.
- 20   •   Other Government Services or Facilities—Lands primarily used to support government  
21       functions, services, or facilities not included in any of the other categories listed above.  
22       These include government offices, community centers, colleges and universities,  
23       military facilities, and cemeteries.

24   Approximately 50 percent of the reported recreation in Washington State is local, meaning  
25   that it occurs close to cities or towns, and includes local parks, playgrounds, and bike or  
26   jogging trails, and other developed facilities, which form about 3.6 percent (660,000 acres)  
27   of the public lands in the State (IAC 2002). The remaining 50 percent of the recreational  
28   use in the State is split evenly between State and Federal lands (IAC 2002). State lands  
29   designated principally as recreational lands represent about 7 percent (648,580 acres) of the  
30   total recreational lands in the State (Table 3-25). These lands are typically located farther  
31   away from urban centers and are used primarily for camping, hiking, hunting, and fishing.  
32   However, most State lands are available to recreational users even though their use  
33   designation may fall within a different category. Federal lands include approximately 91  
34   percent (9,143,462 acres) of all lands designated as recreation lands although, like State  
35   owned lands, they support only about 25 percent of the recreational use in the State (IAC  
36   2002). Federal lands are typically located in remote areas away from urban centers, and  
37   much of their lands tend to be in higher elevations (above 3,000 feet), making it less  
38   accessible to most recreational users (IAC 2002).

39



# Chapter 3

1 **Table 3-25.** Acreage<sup>1/</sup> of Public Lands by Categories Defined by the Interagency  
 2 Committee on Outdoor Recreation.

Landowner	Outdoor Recreation, Habitat, Environmental Protection (Acres)	Resource Production and Extraction (Acres)	Transportation and Utilities Infrastructure (Acres)	Other Government Services and Facilities (Acres)	Total (Acres)	Percent of Total Public Land Base (%)
Federal	9,143,462	2,435,550	656,165	640,358	12,875,535	74
State	648,498	2,836,694	168,876	34,806	3,688,874	22
Local	237,038	65,903	424,580	67,259	794,780	4
<b>Total Public</b>	10,028,998	5,338,147	1,249,621	742,423	17,359,189	100
<b>Total Public (%)</b>	57.8	30.8	7.2	4.3	100	--

<sup>1/</sup> Includes roadway right-of-way easement acres.

Source: Interagency Committee for Outdoor Recreation (IAC) 2001.

3 The IAC (2002) describes lands above 3,000 feet as typically unsuitable for many  
 4 recreational activities due to rough terrain, extreme weather conditions, and seasonal  
 5 limitations. Land above 3,000 feet makes up 28 percent of the State, and much of this land  
 6 (approximately 77 percent) is publicly owned. Lands between sea level and 3,000 feet  
 7 comprise 72 percent of the State, of which nearly 70 percent is privately owned (IAC  
 8 2002).

### 9 3.11.2 Estimates of Future Recreational Needs

10 Future participation in recreational activities is a function of population demographics  
 11 along with land use decisions that are currently being made or are likely to be made in the  
 12 future. In the case of population demographics, an example might be that activities  
 13 requiring some degree of physical stamina such as hiking, mountain biking, and mountain  
 14 climbing are more likely to be attractive to active members of society. In contrast,  
 15 activities such as walking or picnicking generally require less stamina and therefore may  
 16 attract less active people (IAC 2002). As the population ages, demand for less physically  
 17 challenging activities may increase creating demand for local parks or green spaces closer  
 18 to urban centers. Population growth of about 20 percent over the last decade has resulted  
 19 in increased numbers of people engaged in recreation, even though the percent of the  
 20 population actively participating in outdoor recreation declined over this period. More  
 21 than half of the State's population currently participates in some form of outdoor recreation  
 22 (IAC 2002).

23 Outdoor recreation activities that occur on forested lands include walking/hiking,  
 24 horseback riding, off-road vehicle use, picnicking, camping, hunting, fishing, and other  
 25 activities. The IAC assessment (2002) found that 53 percent of the State's population  
 26 participated in the walking/hiking recreation category, with 20 percent picnicking, 13  
 27 percent camping, 13 percent fishing, 9 percent using off-road vehicles, and 6 percent  
 28 hunting/shooting. The IAC assessment was a statewide outdoor recreation survey  
 29 conducted in 1999 and 2000.



1 Participation in all of these activities, with the exception of fishing and hunting/shooting, is  
2 projected to increase by 2023. Increases by 2013 are expected to range from 5 to 10  
3 percent for camping to 20 percent for picnicking. The numbers of people fishing and  
4 hunting/shooting are projected to decrease by 5 percent and 15 percent, respectively, over  
5 the same period (IAC 2003).

6 Land use decisions, including the conversion of forestland into uses incompatible with  
7 timber production also affect recreational trends. Subsection 3.2 (Land Ownership and  
8 Land Use) discusses the rate of land conversion over the past century. Additionally, many  
9 cities in the State have designated urban growth areas where it is expected that future land  
10 uses will be oriented toward urban development rather than being managed or preserved as  
11 forestland. As urban development and forestland conversion continues, there may be a  
12 shift in the availability and types of recreational activities.

### 13 **3.11.3 Private Lands and Recreation**


14 Currently in Washington State, private landowners play a key role in recreation. Smaller  
15 landowners sometimes provide formal facilities, such as campgrounds and golf courses  
16 that are available to the public on a fee basis, while some larger and industrial timberland  
17 owners operate on a “good neighbor” policy and allow the public access to their lands for  
18 general recreational purposes (IAC 2002).

19 In an IAC assessment performed in Washington State in 2002, the recreating public  
20 demonstrated concerns regarding increasing needs for better maintenance of public  
21 facilities, while local managers of public lands statewide expressed concerns about  
22 decreasing funds available for public facilities. As resources for operation and  
23 maintenance of recreation facilities on public lands decline, the public may eventually be  
24 forced to rely more heavily on undeveloped private lands for recreational needs.

### 25 **3.11.4 Forest Road-related Recreation**

26 Existing forest road systems generally receive heavy recreation-related use, providing the  
27 public with access to specific recreation areas, such as trailheads, campgrounds, and picnic  
28 areas. In addition, a large portion of recreational users of managed forestlands use the road  
29 systems as the primary focus of their recreational activity; driving the road systems and  
30 occasionally dispersing across the landscape to hunt, bird watch, gather mushrooms or  
31 berries, or engage in some other non-facility oriented activity. A recent survey, for  
32 example, estimated that approximately 50 percent of back road and “off of road” fuel use  
33 in the State of Washington was associated with hunting, driving, sightseeing, camping, and  
34 fishing, with the other 50 percent used for off-road motorized activities (off-road vehicles  
35 and snowmobiling) and access for non-motorized activities (hiking, mountain biking,  
36 cross-county skiing, and equestrian) (Hebert Research, Inc. 2003).

37



## **Chapter 3**

---

1 This page is intentionally left blank.

2



1    **3.12    VISUAL RESOURCES**

2    **3.12.1    Introduction**

3    High quality scenery, especially scenery with natural appearing landscapes, is generally  
4    regarded as an important resource that enhances peoples’ quality of life, influences the  
5    quality of their recreational experiences, and, in some cases, effects the adjacent property  
6    values. Forestlands are considered to be an important visual or scenic resource by many  
7    Washington State residents and visitors. Approximately 53 percent of the lands in  
8    Washington State are forested (23 million out of 43 million acres) (Table 3-2).

9    Table 3-26 presents the percentage of each analysis region that is forested and the  
10   percentage of the forestlands that are subject to Washington Forest Practices Rules. The  
11   visual resource provided by a forest depends on observer characteristics, as well as the  
12   physical properties of the forest landscape. Important observer characteristics include: the  
13   amount of concern individual observers have for visual quality, the number of observers,  
14   the frequency and duration of their viewing, the context of viewing (e.g., hiking, driving),  
15   and attitudes towards visible forest management activities (Bergen et al. 1995; Sheppard  
16   1989; Magill 1992). Physical properties of the forest landscape that generally influence the  
17   quality of the visual resource provided by forests include: type and density of vegetation,  
18   topography, slope and aspect, presence of water, number and type of viewpoints, distance,  
19   and weather (Bergen et al. 1995; Shepard 1989).

20   Primary areas where forest-related visual concerns exist include major highway corridors,  
21   cities and towns, adjacent housing developments, and trails and other recreation areas.  
22   Forested landscapes in these areas are often highly visible to the public and can be  
23   managed to reduce the visual impact of harvest and road-building activities.

24   The public’s concern with forestland-related visual resources and quality has increased and  
25   continues to increase for a number of reasons. Washington State’s population continues to  
26   grow, increasing by approximately 21 percent, or 1 million people, between 1990 and 2000  
27   (U.S. Census Bureau 2000). Scenic touring and viewing has continued to increase since it  
28   first gained popularity following World War II when more families began to purchase cars  
29   and use them for pleasure trips (Dakin 2003; Wilson 1991). Demands for outdoor  
30   recreation in Washington State have increased, as shown by visits to State parks, which  
31   increased from 12.9 million visits in 1965 to 42.3 million visits in 1997 (Washington DNR  
32   1998h). Outdoor recreation takes place on public and private forestland where allowed.  
33   Sightseeing was identified as the fifth most popular recreation activity by Washington  
34   residents out of 15 recreation categories. Fifty percent of those individuals, who listed  
35   sightseeing as one of their recreation activities, choose to sightsee in scenic areas, often  
36   forestlands, while the other 50 percent sightsee in other areas such as historical areas  
37   (IAC 2002).

38



## Chapter 3

1 **Table 3-26.** Percent Forestland and Percent of Forestland that is Subject to  
 2 Washington Forest Practices Rules by Analysis Region.

<b>Region</b>	<b>Total Acres</b>	<b>Percent of Region that is Forestland (%)</b>	<b>Percent of Forestland Subject to Washington Forest Practices Rules (%)</b>
North Puget Sound	4,397,962	78	45
South Puget Sound	2,178,228	70	73
West Puget Sound	1,735,288	88	57
Islands	246,822	73	92
Olympic Coast	1,756,868	95	45
Southwest	2,313,254	89	93
Lower Columbia	3,072,412	85	62
Middle Columbia	6,497,115	41	35
Upper Columbia – Downstream of GC Dam	6,400,577	43	17
Upper Columbia – Upstream of GC Dam	5,747,801	71	44
Snake River	4,575,868	8	35
Columbia Basin	4,350,198	0	99
<b>Total State</b>	<b>43,272,394</b>	<b>53</b>	<b>50</b>

Source: USGS Land Use/Land Cover Layer, 2004, and WRIA GIS Layer, 2004.

3 Forestlands in Washington State span the vegetation zones from near sea level to the  
 4 subalpine and include a wide range of landscape types and scenic resources, including wet  
 5 coastal forests, high elevation forests, and dry eastern forests. Lowland forests in western  
 6 Washington differ substantially from those in eastern Washington. Western Washington is  
 7 one of the most densely forested regions in the United States. Conifer trees (the most  
 8 common tree type) in this area grow in dense stands, can live long lives, and reach large  
 9 sizes. Douglas-fir is the most common species in this area and can live to 500 years or  
 10 more and reach heights of over 200 feet at maturity. Most of western Washington's forests  
 11 have been harvested at least once since European settlers immigrated to Washington. As a  
 12 result, most of the forestland in western Washington is currently comprised of second or  
 13 third growth even-aged stands that are denser and smaller in size than old growth.

14 Conifer forests in eastern Washington start on the eastern slope of the Cascade Range and  
 15 extend around the northern edge of Washington to the northern Rocky Mountains.  
 16 Ponderosa pine or western yellow pine is the typical species found in lowland eastern  
 17 Washington conifer forests. Ponderosa pine forests often look quite different from western  
 18 Washington conifer forests, because the stands are typically more open and can seem  
 19 almost park-like due to the historical fire regime that maintains them. However, with the  
 20 past century of fire suppression in the State, many of the open, park-like stands that were  
 21 once common in eastern Washington are now more densely stocked.

22 Land management can affect visual resources and different owners manage lands  
 23 differently. Ownership of forestland in Washington State can be divided into six  
 24 categories: Federal and State protected lands where timber management is generally not





1 allowed (e.g., wildernesses, parks, and refuges), Federal lands managed under the  
2 Northwest Forest Plan, State lands managed for timber under the Washington Forest  
3 Practices Rules, tribal forestlands, county and city forestlands, and private forestlands  
4 (Table 3-1). Private forestlands may be divided into industrial forestlands where intensive  
5 forest practices take place and other private forestlands where forest practices are generally  
6 periodic.

7 Federal- and State-protected lands are not managed for timber production and are,  
8 therefore, not likely to be affected by Washington Forest Practices Rules. Federal lands  
9 managed under the Northwest Forest Plan are managed under an extensive set of scenery  
10 management rules (USDA Forest Service and USDI Bureau of Land Management 1994).  
11 State lands managed for timber are also managed under policies that require consideration  
12 of potential visual impacts, but they permit more extensive visual alteration than the  
13 Federal Northwest Forest Plan (USDA Forest Service and USDI Bureau of Land  
14 Management 1994). Private, tribal, and other lands can also be managed for timber, often  
15 with fewer restrictions on harvest and road building for visual reasons. When these  
16 forestlands are converted to other land uses, the visual resource can be substantially  
17 impacted (conversion is discussed in subsection 3.2, Land Ownership and Use).

### 18 **3.12.2 Visual Resources and the Current Washington Forest Practices** 19 **Rules**

20 When forests are managed for timber in Washington State, landowners are subject to  
21 Washington Forest Practices Rules, which include prescriptions that, although not their  
22 primary intent, provide some protection for visual resources during timber harvest.  
23 Clearcut harvest areas are prohibited from being larger than 240 acres. Clearcuts must  
24 retain at least five uncut trees per acre in western Washington and four uncut trees per acre  
25 in eastern Washington for wildlife. Uncut trees are also sometimes required to be left on  
26 unstable slopes (WAC 222-10-030 (4)), and landowners are required to leave riparian  
27 buffers adjacent to most streams.

28 Riparian buffers requirements, which have existed for nearly two decades, have increased  
29 since the current Washington Forest Practices Rules were implemented in April 2000.  
30 Under these rules the required widths for buffers adjacent to streams are wider and apply to  
31 more streams than was previously required (WAC 222-30). The Washington Forest  
32 Practices Rules contribute to the protection of visual resources on State and private  
33 forestlands. In addition to these rules, many private forest landowners voluntarily leave  
34 additional buffers specifically for visual resource protection. Many Industrial forest  
35 landowners use *Guidelines For The Design Of Harvest Practices In Visually Sensitive*  
36 *Areas* (Bradley 1996) to assist them in planning harvests when they are dealing with  
37 visually sensitive landscapes.

38 Practices employed by industrial forest landowners in visually sensitive landscapes may  
39 include leaving uncut buffers along major highways, leaving buffers along topographic  
40 features to reduce the visual impact of the timber harvest, using selective harvest methods  
41 instead of clearcut harvesting, harvesting timber in a way that simulates natural patterns,  
42 providing roadside signs that explain the forest practices that have taken place, minimizing



## **Chapter 3**

---

- 1 forest openings next to highways, protecting special vistas with no-harvest areas, and using
- 2 intensive management to help the harvest area regenerate quickly (Merrill and Ring 1999;
- 3 Plum Creek 1999; Champion Pacific Timberlands 2000).
- 4



1     **3.13     ARCHEOLOGICAL, HISTORICAL, CULTURAL AND INDIAN**  
2             **TRUST RESOURCES**

3     **3.13.1   Introduction**

4     Cultural resources are districts, sites, buildings, structures, and objects that contain  
5     evidence of past human activities or that play an active part in the traditional cultures of the  
6     disparate ethnic groups that comprise Washington’s populace. Cultural resources have  
7     been recognized by legislative bodies at the Federal and State levels as being important for  
8     the education and inspiration of future generations of Americans, whatever their  
9     backgrounds. Four categories of cultural resources are recognized and discussed here and  
10    in the environmental effects section, as defined below.

11    **Archaeological Resources**—the physical residues of past human activity. Archaeological  
12    resources provide evidence of the cultural continuum of people occurring across time and  
13    space throughout the diverse landscapes of Washington. Examples of archaeological  
14    resources include shell middens, scatters of stone chips (lithic scatters), rock art, talus slope  
15    gravesites, and culturally modified trees.

16    **Historic Sites**—locations where Native or non-Native events and activities have taken  
17    place since contact with Euro-Americans. Historic sites often, but not always, have written  
18    records that document the events and activities that occurred at a particular location.  
19    Examples of historic sites include homesteads, forts, lumber mills, cabins, mine shafts, and  
20    old logging or mining equipment.

21    **Traditional Places**—landscapes, resource gathering areas, sacred sites, and legendary  
22    areas that are identified (often with traditional names) by Indian Tribes in the State of  
23    Washington as being important for the maintenance and perpetuation of their traditional  
24    values and practices. These landscapes, places, and objects provide subsistence and  
25    spiritual relationships, as well as stability and meaning to community ceremonies, customs,  
26    and beliefs. Examples of traditional places include sacred ceremonial sites, groves used for  
27    gathering edible/medicinal plants, and sources of materials used for traditional tools and  
28    arts.

29    **Traditional Materials**—the resources used by Native peoples to sustain their culture.  
30    Traditional materials come from the broad variety of plants, animals, and minerals that are  
31    indigenous to this region’s native landscapes. Traditional and current cultural values for  
32    plants include their use as medicines, foods, tools, textiles, building materials, carvings,  
33    and sacred objects. Examples of traditional materials (such as some of the plants utilized  
34    by Tribes) include bear grass, tule, and cedar and birch trees.

35    **3.13.2   Archaeological Overview**

36    Lands lying west and east of the Cascade Range have been ecologically distinct for most of  
37    the postglacial period as well as being geographically separated by the mountain range.  
38    Consequently, cultural developments and archaeological site distributions are also distinct.



## Chapter 3

---

### 1 3.13.2.1 Western Washington

2 Despite nearly a century of scientific research in the region, the archaeology of western  
3 Washington is not well understood. Most of what is known comes from low elevation  
4 lands that are now largely under agricultural and residential development. The foothills  
5 and lower mountain settings, where most of the region's productive forests can be found,  
6 have received relatively little attention from archaeologists. What is known about the  
7 prehistoric archaeology of the region is, consequently, biased toward the lowlands,  
8 particularly coastlines. Much of the property managed by the State and the majority of  
9 privately owned forestlands has not been intensively surveyed for archaeological resources;  
10 the same is true for nearby lands of the National Forests. Most archaeological sites in these  
11 forests have been found along streams or on high ridges, but this may be due in part to a  
12 tendency for land managers to survey what they consider high probability areas more  
13 intensively than lower probability slopes. What is currently known about the archaeology  
14 of western Washington is summarized below.

15 The first human occupation of Washington State may date back about 14,000 years<sup>1</sup> to the  
16 Manis Mastodon site at Sequim. Artifacts of the 13,000- to 13,500-year-old Clovis culture,  
17 which is thought to have been focused on big game hunting, have been found on the  
18 ground surface in such places as Thurston County and Whidbey Island, but no campsite of  
19 this culture has yet been found in Washington.

20 The post-Clovis prehistory of western Washington has been divided into three periods,  
21 designated as Early, Middle, and Late. The Early Period, which lasted from approximately  
22 12,000 to 7,000 years ago, includes the Proto-Western and Old Cordilleran Traditions  
23 (Matson and Copeland 1995). Sites left by these traditions typically occur on high marine  
24 and river terraces, sometimes at a significant distance from modern water courses, and  
25 consist of concentrations of cobble cores, flakes, large, ovate knives, and broad-stemmed  
26 and leaf-shaped projectile points (Wessen 1990). In some areas, sites of this period have  
27 also been documented along mountain streams in open sites, rockshelters, and caves  
28 (Wessen and Stilson 1986; Lewarch and Benson 1989). Because of an apparent inland  
29 focus, the people of this era are thought to have been more oriented to land animal hunting  
30 and less to marine and fish resources.

31 The Middle Period, lasting from 7,000 to 3,500 years ago sees a continuation of the Old  
32 Cordilleran Tradition until around 4,500 years ago, but few sites can be attributed to this  
33 time interval (Morgan 1999). The focus of subsistence activity after 4,500 years ago seems  
34 to have changed from terrestrial to marine resources and most sites appear along the coasts  
35 or major river systems. There is little evidence of activity in the higher mountains. Tools  
36 are more complex, including tools and ornaments of bone and antler along with chipped  
37 stone (e.g., Larson and Lewarch 1995).

38 The concentration on marine resources intensified during the Late Period (3,500 to 150  
39 years ago), and the number and diversity of sites increased markedly. People maintained

---

<sup>1</sup> Dates given here are in calendric years, based on approximate calibration of radiocarbon ages.



1 permanent villages on the coast and along the lower reaches of inland rivers, which they  
2 used as home bases and storage warehouses. Huge shell middens built up at some villages  
3 and at the best clam beaches. Cemeteries and petroglyph sites are often associated with  
4 village and midden sites and fishing camps and occur occasionally in higher montane  
5 settings. Blazed cedars, stripped of bark for basketry or with planks removed from their  
6 living trunks, can still be found throughout the lowlands. Small open camps, left by  
7 hunters, fishers, berry pickers (Mack and McClure 2002), and traders have been found  
8 from the lowlands well into the subalpine zone of the mountains, but usually remain close  
9 to larger, permanent sources of water. They typically are concentrated along trade routes.

10 Archaeological sites in western Washington consist of the remains of villages and their  
11 associated cemeteries and petroglyph sites, saltwater shell middens, temporary camps  
12 consisting of stone flaking debris (lithic scatters), berry processing camps, rockshelters and  
13 caves, and stone quarries. With the exception of quarrying areas, berry processing  
14 localities and some caves, rockshelters, and petroglyphs, most archaeological sites in  
15 western Washington occur close to streams and shorelines (Nelson 1990).

16 A type of archaeological site that has recently been recognized is the culturally-modified  
17 tree. These are typically living cedar trees that have had bark stripped from one or more  
18 sides of the tree for use in making baskets or clothing. Culturally-modified trees are  
19 expected in stands of old-growth cedar that predate intensive Euro-American settlement  
20 (Larson 1998; Gunther 1973).

### 21 **3.13.2.2 Eastern Washington**

22 Lands east of the Cascade Range tend to be drier and more open, except at higher  
23 elevations, and the archaeology of the region is proportionately better known than the  
24 western region. The prehistory of this region begins much the same as that of western  
25 Washington, with scattered manifestations of the Clovis Culture. Post-Clovis archaeology  
26 is again divided into Early, Middle and Late, but the beginning and ending points of these  
27 periods differ from those of western Washington (Ames et al. 1998; Chatters and Pokotylo  
28 1998).

29 The Early Period, from 11,000 to 8,000 years ago, is characterized by large, stemmed  
30 projectile points, ovate knives, barbed harpoons, grooved net weights, and a stone tool  
31 technology reminiscent of the upper Paleolithic technologies of Eurasia. Few sites of this  
32 time period have been found, most of them consisting of small campsites along rivers and  
33 streams of the non-forested lowlands. A few sites of this period have been reported from  
34 the Blue Mountains, but none are yet known from the Cascades.

35 During the Middle Period, 8,000 to 5,000 years ago, the region was occupied by a variant  
36 of the Old Cordilleran Culture, known locally as Cascade. Cascade sites are ubiquitous  
37 along the Snake and Columbia Rivers, and their larger tributaries and occasionally occur as  
38 open camps and in rockshelters in all mountain ranges. There is some indication that  
39 quarries in the eastern Cascade Range were in use at this time. Artifacts of this period  
40 resemble comparable assemblages from western Washington, consisting of cobble tools,



## Chapter 3

---

1 cobble flakes, leaf-shaped projectile points and knives, simple scraping tools, occasional  
2 microblades, and little else (Ames et al. 1998).

3 During the Late Period, which is comprised of a sequence of distinct cultural adaptations,  
4 people had begun to settle in pithouse villages along major streams. They followed a  
5 seasonal subsistence routine that included early springtime camps away from the villages  
6 but still in riverine settings, root gathering camps near springs in foothills and at mountain  
7 meadows, berry picking camps in the subalpine forests, and hunting camps throughout the  
8 landscape in caves, near springs, and along water courses. Special purpose sites are  
9 common from this Period and include cemeteries near villages, isolated graves and rock art  
10 panels in many settings, quarrying areas, spirit quest cairns and game drives on ridgetops,  
11 hunting blinds and storage pits in talus slopes, storage caves, and culturally modified trees.  
12 In eastern Washington forests, culturally modified trees may include cedars stripped of  
13 bark for making baskets or pines stripped of edible cambium (Hunn et al. 1998).

14 There is an even more marked tendency in this region for sites to concentrate near springs  
15 and streams. Cemeteries, villages, and most camps are rarely far from water, although  
16 special purpose sites may occur at considerable distance (Galm et al. 1981).


### 17 **3.13.3 Cultural and Trust Resources of Native American Tribes**

18 Historic native cultures of the region can generally be seen as a continuation of the  
19 lifeways indicated by Late Period archaeological sites. The region's peoples belonged to  
20 five linguistic groups: Wakashan, Salishan, Chimakuan, Chinookan, and Sahaptian.  
21 Wakashan, Chinookan, Chimakuan, and most Salishan peoples were marine oriented,  
22 occupying villages on the major rivers or saltwater shorelines of western Washington and  
23 focusing on shellfish, and salmon and/or saltwater fish for their subsistence. These peoples  
24 abandoned their villages in summer, moving among fishing sites, and hunting, root  
25 gathering, and berrying camps in mountains and prairies (Haeberlin and Gunther 1930;  
26 Silverstein 1990). The Salishan Snoqualmie and the Sahaptian-speaking Klikitat differed  
27 from other western Washington Tribes, spending most of their time in foothill and  
28 mountain settings, where they emphasized hunting, berrying, and root gathering, and  
29 served as intermediaries in the transmontane trade (Suttles and Lane 1990; Walker 1998).

30 All eastern Washington Tribes belong either to the Salishan or Sahaptian language  
31 families. Members of these groups focused on salmon fishing, root and berry gathering,  
32 and big game hunting for their subsistence in varying proportion, depending on the  
33 resources of their respective territories. In a continuation with the Late Period of  
34 prehistory, they occupied villages of pithouses or mat lodges in winter along major stream  
35 courses and moved among a series of smaller camps or multi-band conclaves as the year  
36 progressed. They were linked to the peoples of the western flank of the Cascades and  
37 Lower Columbia River through trade and intermarriage (Miller 1998; Schuster 1998).

38 Today, 29 federally recognized Tribes reside in Washington (Table 3-27) and over a dozen  
39 Tribes that once lived in what is now Washington State, reside in adjacent states and  
40 Canada. The Tribes that entered into treaties with the United States in 1855 (i.e., all the

# Chapter 3



1 **Table 3-27.** Federally-Recognized Tribes of Washington and Adjacent States  
 2 with Cultural Interests in Washington Forests.

Region	Tribes
Eastern Washington	Coeur D’Alene <sup>1</sup> , Colville, Nez Perce <sup>1/</sup> , Spokane, Pend d’Oreille, Umatilla <sup>1</sup> , Yakama
Western Washington	Shoalwater Bay Chinook, Chehalis, Cowlitz, Hoh, Lummi, Makah, Muckleshoot, Nisqually, Nooksack, Puyallup, Quileute, Quinault, Sauk-Suiattle, Upper Skagit, Lower Elwha, Jamestown, Port Gamble, S’Klallum, Skokomish, Snoqualmie, Squaxin Island, Stillaguamish, Suquamish, Swinomish, Tulalip

<sup>1/</sup> With reservations outside Washington.

3 Tribes in Table 3-27 except the Colville, Chehalis, Cowlitz, Shoalwater Chinook, Spokane,  
 4 and Coeur D’Alene) all retained the right to certain resources on ceded territories. All  
 5 treaties reserve the right to fish in usual and accustomed grounds and places in common  
 6 with other citizens, and to hunt and gather roots and berries on open and unclaimed land.

7 The right to fish carries with it a trust responsibility on the part of the Federal government  
 8 to help ensure the continued productivity of the fishery.

9 Forests provide raw materials for Washington’s Tribes (e.g., bark and grasses for baskets  
 10 and mats, wood for carvings, and medicinal plants) as well as subsistence resources  
 11 (Gunther 1973). While each tribal group is different in the plants and animals that it  
 12 considers culturally important, and while many of the medicinal and ceremonial species are  
 13 kept secret for religious or other cultural reasons, some plants and animals that have  
 14 importance across many groups are listed in Table 3-28. Salmon, deer, elk, cedar,  
 15 beargrass, and huckleberries are the most prominent among these resources.

16 Of the culturally important resources listed in Table 3-28, most occur or are most abundant  
 17 in the water (fishes, shellfish, some mammals), streamside wetlands (tule, cattail),  
 18 meadows (camas, nettle, many berries and roots), or early seral stages of forests (most  
 19 berries, beargrass, nettle, bracken, alder, big game). Some species, however, occur or are  
 20 most useful when found in old-growth stands (cedar, spruce, tree lichen) (Turner 1978;  
 21 Gunther 1973).

22 Mature forests also provide solitude that is necessary for individuals’ quests for personal  
 23 spirit guidance, and for the observance of spirit dances among some Salishan groups  
 24 (Hajda 1990; Kew 1990; Suttles and Lane 1990). The quest for spiritual guidance begins  
 25 before puberty and continues throughout a person’s life (Haeberlin and Gunther 1930;  
 26 Walker 1998). Forestlands also contain many traditional places, which can take many  
 27 forms, from rock outcroppings or caves to cedar groves or camas meadows. Some Tribes  
 28 have worked with land managers to identify sacred places and stands of culturally  
 29 important plants (e.g., Blukis-Onat and Hollenbeck 1981), while others prefer to keep such  
 30 locations secret.

31

32



# Chapter 3

1 **Table 3-28.** Culturally Important Plant and Animal Resources for Native  
 2 American Tribes in Washington.

Resources	Western Washington <sup>1/</sup>	Eastern Washington <sup>2/</sup>
Fishes <sup>3/</sup>	Chinook, coho, pink, chum, and sockeye salmon, steelhead, cutthroat trout, dolly varden trout, smelt	Chinook, coho, and sockeye salmon, steelhead, bull trout, cutthroat trout, lamprey, suckers, mountain whitefish
Shellfish <sup>3/</sup>		Freshwater mussels
Mammals <sup>4/</sup>	Blacktail deer, elk, bear, otter, beaver	Deer, elk, blackbear, otter
Berries, Roots, and Other Foods	Huckleberries, blueberries, trailing blackberry, black cap, salmonberry, thimbleberry, salal berry, serviceberry, soapberry; wild carrot, wapato, camas, bracken root, sand verbena, thistle root, surf grass, buttercup, clover roots; cow parsnip	Blue huckleberry, serviceberry, chokecherry, currant, blackberry; camas, wild carrot, chocolate lily; tree lichen; pine cambium; wild celery
Raw Materials	Yew, alder, and cedar wood, spruce root, cedar, hemlock, and willow bark, beargrass, nettle, cattail, and tule	Cedar bark, wood and boughs, beargrass, tule

<sup>1/</sup> Renker and Gunther 1990; Gunther 1945; Powell 1990; Suttles and Lane 1990; Silverstein 1990.

<sup>2/</sup> Schuster 1998; Ross 1998; Kennedy and Bouchard 1998; Stern 1998; Walker 1998.

<sup>3/</sup> Freshwater or anadromous, only.

<sup>4/</sup> Terrestrial only.

### 3 3.13.4 Overview of Regional History

4 Washington’s coastline was first charted and described by English and American Explorers  
 5 in the last decades of the eighteenth century. Fur traders, primarily associated with Forts  
 6 Vancouver, Nisqually, Walla Walla, Okanagan, and Spokane, traveled into the interior in  
 7 the first half of the nineteenth century. Except for the increasing presence of beads, metal,  
 8 and other trade goods among the local Indian Tribes, they left few traces outside their fort  
 9 compounds. By the 1830s, the Hudson’s Bay Company had expanded into agricultural  
 10 production, maintaining large farms in the lowlands around Forts Vancouver and Nisqually  
 11 and in the lower Cowlitz River Valley. Missionaries soon followed the fur traders, setting  
 12 up missions in the Yakima and Walla Walla Valleys during the 1840s. Settlers, some  
 13 drawn by the promise of farmland, but most coming to exploit the region’s timber and  
 14 mineral wealth began flowing into the Puget Lowlands and Portland Basin by the late  
 15 1840s and into eastern Washington after the 1855 signing of treaties with the region’s  
 16 Indian Tribes. The farmers needed water and, like their Native predecessors, settled first in  
 17 valley bottoms and along small stream courses, particularly in the drier eastern part of the  
 18 state. Loggers and fishermen built along the coasts and exploited lowland timber from the  
 19 onset of settlement. By the latter part of the nineteenth century, Euro-Americans also  
 20 began to exploit the timber and mineral resources of the non-arable mountains in western





1 Washington, and began to mine, log, and graze stock in the open forests of eastern  
2 Washington (Avery 1965). Railroads and haul roads were built into and across the  
3 Cascades for hauling timber, coal, stock, and produce to markets and to connect  
4 Washington with the rest of the United States.

5 Past farming and stock raising activities are marked by buildings, water control structures,  
6 fences, and stock camps. Mining has left its traces throughout the uplands of Washington.  
7 Coal mining is marked by large, open pit mines and haul roads, whereas traces of past gold  
8 mining occur as placer tailings, mining prospects, mine shafts, and miners' camps. Sites  
9 associated with logging include skid roads, railroad grades and tracks, trestles, construction  
10 and logging camps, stumps cut with springboard notches, and a variety of equipment.

### 11 **3.13.5 Washington State Protective Structure**

12 Washington State makes use of many venues for protection and preservation of  
13 archaeological, historical, and cultural resources, including the Office of Archaeology and  
14 Historic Preservation, protective statutes in the RCW, and protective regulations in the  
15 WAC.

16 The Office of Archaeology and Historic Preservation provides expertise, service, and  
17 training for the protection and preservation of Washington's historic places. The office  
18 manages programs to aid historic property owners, community leaders, and preservation  
19 advocates. A key role of the Office of Archaeology and Historic Preservation is the  
20 identification, evaluation, and protection of significant properties worthy of preservation  
21 (Washington State Community, Trade, and Economic Development 1996). The office is a  
22 center for data collection and administers the National Register of Historic Places and the  
23 Washington Heritage Register. The Office of Archaeology and Historic Preservation is a  
24 part of the Washington State Department of Community, Trade, and Economic  
25 Development. The Director of this department, or their designee, is a member of the  
26 Forest Practices Board, the promulgating agency for Washington Forest Practices Rules.

27 Various state agencies interface with archaeological, historical, and cultural resources. The  
28 Washington Department of Transportation has a Cultural Resources Program that keeps the  
29 public, including Washington Tribes, informed about Department of Transportation  
30 projects and actively seeks the public's input on concerns about project impacts on  
31 archaeological, historical, or cultural resources (Personal Communication, Sandie Turner,  
32 Transportation Planning Specialist, Department of Transportation, August 19, 2004).  
33 Washington State Parks and Recreation protects many archaeological, historical, and  
34 cultural resources on their properties. Washington DNR interfaces with cultural resources  
35 in both the regulatory arena and the land management arena.

36 The Washington DNR Forest Practices Regulatory Program regulates forest practices in  
37 the state including forest practices affecting archaeological, historical, and cultural  
38 resources on both private and State land (WAC 222). In addition to the forest practices  
39 regulations, the Cultural Resource Protection and Management Plan, written and agreed to  
40 by TFW participants on July 3, 2003, provides a process to enhance protection of cultural



## Chapter 3

---

1 and archeological sites on managed forestlands (Timber, Fish, and Wildlife Cultural  
2 Resources Committee 2003).

3 Washington DNR State Lands program manages 2.1 million acres of forestland in the State  
4 where archaeological, historical, and cultural resources may be found. Washington DNR  
5 has a tribal Relations Manager who is a liaison between Washington DNR State Lands and  
6 Washington Tribes and is working on strengthening and focusing Washington DNR State  
7 Lands' Cultural Resources program (Personal Communication, Joseph Davis, Washington  
8 DNR, August 18, 2004). The Commissioner of Public Lands, the head of Washington  
9 DNR, recently reaffirmed Washington DNR's commitment of collaborative tribal  
10 relationships in a Commissioner's Order, stating that the Department will "actively work  
11 with Tribes to encourage understanding and the cooperative pursuit of common objectives"  
12 (Department of Natural Resources, Commissioner's Order, April, 26, 2004a.)

13 The Department of Transportation, Washington State Parks and Recreation, and the Office  
14 of Archaeology and Historic Preservation offer a weeklong semi-annual training course on  
15 cultural resources. The course provides information for recognition and protection of  
16 resources as well as information on State agency legal obligations centered on  
17 archaeological, historical, and cultural resources. The course includes site visits to help  
18 students learn how to recognize an archaeological, historical, or cultural resource.



1    **3.14    SOCIAL AND ECONOMIC ENVIRONMENT**

2    **3.14.1    Introduction**

3    This section presents a general overview of the social and economic conditions of  
4    Washington State, and provides a baseline that the potential effects of the alternatives may  
5    be measured against. The discussion is organized into three sections that address  
6    population, employment and the economy, and environmental justice.

7    **3.14.2    Population**

8    The total population of Washington State was approximately 5.9 million in 2000, an  
9    increase of 21 percent, or approximately 1 million people, since 1990. All Washington  
10    counties experienced population growth during the 1990s, with increases ranging from  
11    1 percent in Columbia County to 45 percent in Clark County. Net in-migration accounted  
12    for 63 percent of statewide population growth over this period and contributed to  
13    population increases in all Washington counties, with the exception of Whitman County,  
14    where net out-migration accounted for approximately -1 percent of population change.  
15    The majority of Washington counties also experienced natural increase (more births than  
16    deaths) over this period. There were, however, some exceptions, including Columbia,  
17    Garfield, Pacific, Wahkiakum, Clallam, Jefferson, and Lincoln Counties, where there were  
18    more deaths than births (Washington Office of Financial Management 2004).

19    The statewide population density in 2000 was 88.6 persons per square mile compared to a  
20    national average of 79.6 percent. Population density varies considerably by county,  
21    ranging from 4.7 persons per square mile in Columbia County in southeast Washington to  
22    817 persons per square mile in King County. Population densities were, not surprisingly,  
23    highest in those counties that include major urban areas: the Puget Sound area (King,  
24    Kitsap, Pierce, Snohomish, and Thurston Counties), Vancouver (Clark County), and  
25    Spokane (Spokane County). Population densities in counties elsewhere in the State were  
26    less than 100 persons per square mile and many counties, particularly those located in the  
27    northeast part of the State, had population densities below 20 persons per square mile  
28    (Washington Office of Financial Management 2004).

29    Population projections developed by the Washington Office of Financial Management in  
30    2002 anticipate continued growth in the State, with the total State population projected to  
31    reach 7.5 million by 2020. Population is projected to increase in all counties over this  
32    period, with increases ranging from 2 percent in Columbia County to 50 percent in  
33    Thurston County (Washington Office of Financial Management 2002a).

34    **3.14.2.1    Race and Ethnicity**

35    Approximately 85 percent of Washington's population identified as white in the 2000  
36    Census, compared to 75 percent nationwide. Compared to the national average, Black or  
37    African American persons and persons of Hispanic or Latino origin were relatively under-  
38    represented, comprising smaller shares of the State population than the national average.  
39    Asian and American Indian persons were, by the same measure, relatively over



## Chapter 3

---

1 represented, comprising slightly larger shares of the State population than the national  
2 average (U.S. Census Bureau 2000).

3 The percentage of the population identifying as white varied by county, ranging from  
4 approximately 62 percent in Franklin County to approximately 97 percent in Garfield  
5 County. Persons of Hispanic or Latino origin comprised relatively large shares of the  
6 population in Adams (40 percent), Franklin (39 percent), Yakima (28 percent), and Grant  
7 (23 percent) Counties. American Indians made up relatively large shares of the population  
8 in Ferry (18 percent), Okanogan (12 percent), and Yakima (4.5 percent) Counties  
9 (Washington Office of Financial Management 2002b).

### 10 **3.14.3 Employment and the Economy**

11 This section provides a general overview of employment and the economy in Washington  
12 State. The section discusses two summary measures developed by Federal and State  
13 agencies—socioeconomic resiliency and the distressed areas index. It also addresses  
14 overall employment trends and employment in the lumber and wood products, commercial  
15 fishing, and recreation and tourism industries.

#### 16 **3.14.3.1 Socioeconomic Resiliency**

17 Socioeconomic resiliency refers to the ability of an area's population and economy (e.g.,  
18 community, county, or region) to adapt to economic changes or shocks. Resiliency is  
19 generally related to diversity, with areas or socioeconomic systems with higher diversity  
20 less affected by a change in the system, such as a change in timber harvest or grazing  
21 opportunities. A high degree of resiliency implies that an area or socioeconomic system  
22 adapts quickly to economic fluctuations or changes, such as specific firms or business  
23 sectors experiencing downturns, with unemployment rates rising only briefly until  
24 displaced individuals find other employment. Areas or socioeconomic systems with low  
25 resiliency may, on the other hand, experience long-term negative impacts, with  
26 unemployment or out-migration rates remaining high for several years.

27 A recent study prepared for Washington DNR assessed the socioeconomic resiliency of  
28 Washington counties employing the methodology used to develop measures of  
29 socioeconomic diversity for the Interior Columbia Basin Ecosystem Management Project  
30 (ICBEMP) (Daniels 2003; Horne and Haynes 1999). The objective of the Washington  
31 DNR study was to identify those counties where changes in timber harvest on State lands  
32 may have disproportionate negative effects on the well being of county residents. Part of  
33 this analysis involved identifying counties of concern, which were defined as those  
34 counties with low socioeconomic resiliency and high forest dependency. A socioeconomic  
35 resiliency index was developed based on a composite of three related factors: lifestyle  
36 diversity, economic diversity, and population density. Forest dependence was identified  
37 based on the percent of total county area classified as forestland. Ferry, Pend Oreille,  
38 Pacific, Skamania, Stevens, and Wahkiakum Counties all had low socioeconomic  
39 resiliency and high forest dependency and were identified as counties of concern. Adams,  
40 Okanogan, Klickitat, Columbia, Garfield, and Lincoln Counties also had low  
41 socioeconomic resilience, but had low or medium forest dependence ratings (Daniels  
42 2003).



1 **3.14.3.2 Distressed Areas Index**

2 The Washington Employment Security Department identifies distressed counties using 3-  
 3 year average unemployment rates. A county is considered distressed if the average  
 4 unemployment rate is 120 percent or greater than the average statewide unemployment rate  
 5 (Washington State Employment Security Department 2003). Nineteen counties were  
 6 identified as distressed areas in 2003 (Table 3-29).

7 **Table 3-29. State of Washington Distressed Counties, 2003.**

State/County	3-Year Average Unemployment Rate <sup>1/</sup>	County	3-Year Average Unemployment Rate <sup>1/</sup>
Washington State	6.3	Grays Harbor County	9.9
Adams County	10.3	Klickitat County	13.4
Chelan County	8.8	Lewis County	9.1
Clallam County	7.6	Mason County	7.7
Columbia County	11.1	Okanogan County	10.8
Cowlitz County	9.4	Pacific County	8.6
Douglas County	8	Pend Oreille County	9.5
Ferry County	13.2	Skamania County	10.5
Franklin County	9.1	Stevens County	10.1
Grant County	9.9	Yakima County	10.7

<sup>1/</sup> The 3-year averages are for January 2000 through December 2002.

Source: Washington State Employment Security Department 2003.

8 **3.14.3.3 Employment**

9 Employment is summarized by sector for Washington State in Table 3-30 for 1990 and  
 10 2000. Overall Washington State employment increased by 24 percent or 688,915 jobs  
 11 between 1990 and 2000, with the largest absolute gains occurring in the services (+325,865  
 12 jobs), retail trade (+125,545 jobs), and State and local government (+85,630 jobs) sectors.  
 13 Sectors with relatively large increases included services and also the agricultural services,  
 14 forestry, fishing, and other sectors. The agricultural services and forestry subsectors, also  
 15 shown in Table 3-30, experienced relatively large employment increases, while the fishing  
 16 subsector experienced large relative and absolute job loss over this period (U.S. Bureau of  
 17 Economic Analysis 2004a).

18 The largest absolute job loss occurred in the manufacturing sector (-17,577 jobs), with job  
 19 loss in the lumber and wood products accounting for approximately 47 percent of this  
 20 decline. Job loss also occurred in the Federal government sector, with a net loss of 4,610  
 21 jobs in the civilian sector, and a further net loss of 8,896 jobs in the military (Table 3-30)  
 22 (U.S. Bureau of Economic Analysis 2004a).

23 The following sections discuss lumber and wood products, commercial fishing, and  
 24 recreation and tourism.

25



## Chapter 3

1 **Table 3-30.** Washington Employment by Sector, 1990 and 2000.

	1990		2000		1990-2000	
	Jobs	Percent of Total (%)	Jobs	Percent of Total (%)	Absolute Change	Percent Change (%)
TOTAL EMPLOYMENT <sup>1/</sup>	2,864,111	100	3,553,026	100	688,915	24
<b>By Type</b>						
Wage and Salary Employment	2,371,705	83	2,940,967	83	569,262	24
Proprietors Employment	492,406	17	612,059	17	119,653	24
<b>By Industry</b>						
Farm Employment	81,399	3	80,004	2	-1,395	-2
Nonfarm Employment	2,782,712	97	3,473,022	98	690,310	25
Agricultural Services, Forestry, Fishing, and Other	49,242	2	63,384	2	14,142	29
Agricultural Services <sup>2/</sup>	32,777	1	48,273	1	15,496	47
Forestry <sup>2/</sup>	4,075	0	5,693	0	1,618	40
Fishing <sup>2/</sup>	12,390	0	9,418	0	-2,972	-24
Mining	5,637	0	5,411	0	-226	-4
Manufacturing	388,748	14	371,171	10	-17,577	-5
Lumber and Wood Products <sup>2/</sup>	45,664	2	37,460	1	-8,204	-18
Transportation and Public Utilities	128,055	4	168,164	5	40,109	31
Wholesale Trade	141,561	5	168,279	5	26,718	19
Retail Trade	468,693	16	594,238	17	125,545	27
Finance, Insurance, and Real Estate	218,487	8	267,150	8	48,663	22
Services <sup>3/</sup>	747,216	26	1,073,081	30	325,865	44
Government and Government Enterprises	475,689	17	547,813	15	72,124	15
Federal, Civilian	73,745	3	69,135	2	-4,610	-6
Military	81,702	3	72,806	2	-8,896	-11
State and Local	320,242	11	405,872	11	85,630	27

<sup>1/</sup> Total employment includes self-employed individuals. Employment data are by place of work, not place of residence and, therefore, include people who work in the area but do not live there. Employment is measured as the average annual number of jobs, both full- and part-time, with each job that a person holds counted at full weight.

<sup>2/</sup> Agricultural Services, Forestry, and Fishing are subcategories of the Agricultural Services, Forestry, Fishing and Other Standard Industrial Classification (SIC) division. Lumber and Wood Products is a subcategory of the Manufacturing SIC division.

<sup>3/</sup> Examples include hotels/motels, business services, automotive repair and other services, amusement and recreation services, health services, legal services, educational services, social services, engineering and management services, and accounting services.

Source: U.S. Bureau of Economic Analysis 2004a.

### 2 **Lumber and Wood Products**

3 Statewide, a total of 37,460 people were directly employed in the lumber and wood  
 4 products industry in 2000, approximately 8,200 or 18 percent fewer than were employed in  
 5 this sector in 1990 (Table 3-30). Timber harvest results in direct employment in the  
 6 logging and lumber and wood products sectors. It also generates indirect and induced



1 employment. Indirect employment includes jobs associated with industries that supply  
2 inputs to the harvesting and processing sector. Induced employment includes jobs  
3 associated with spending in the economy from the salaries created by direct and indirect  
4 effects.

5 Direct employment in the lumber and wood products sector accounted for about 1 percent  
6 of total employment statewide, ranging from no covered employment in a number of  
7 counties (e.g., Adams, Garfield, and Grant Counties) to 10 or more percent of total covered  
8 employment in Mason (10 percent), Stevens (12 percent), and Wahkiakum (25 percent)  
9 Counties.<sup>1</sup> Viewed in absolute terms, the counties with the largest number of covered jobs  
10 in this sector in 2001 were King (5,500 jobs), Pierce (3,500 jobs), and Snohomish (3,000  
11 jobs), which together accounted for approximately 41 percent of covered employment in  
12 the lumber and wood products sector. Lewis and Grays Harbor Counties followed with  
13 approximately 2,100 lumber and wood products jobs each (Washington State Employment  
14 Security Department 2004). Covered employment and wage data indicate that the average  
15 wage in the lumber and wood products sector in Washington State was approximately  
16 \$39,700 in 2001 (Washington State Employment Security Department 2004).

17 Washington State was the second largest softwood lumber producer in the United States in  
18 2001. Lumber production fluctuated during the 1990s, ranging from 3,820 million board  
19 feet in 1991 to 4,384 million board feet in 2000. A total of 4,257 million board feet were  
20 produced in 2001 (Warren 2003).

21 Washington timber harvest levels have shown an overall pattern of decline since 1990  
22 (Figure 3-10). Harvest levels fluctuated during the 1990s, but ranged from a peak of 5,850  
23 million board feet in 1990 to a low of 3,582 million board feet in 2002. Harvest declined  
24 for all ownerships over this period, with the exception of tribal harvests, which increased  
25 from 182 million board feet in 1990 to 319 million board feet in 2002. Harvest on lands  
26 owned by large private (non-industrial) landowners was higher in 2002 than during most of  
27 the preceding decade (Figure 3-10). The largest relative decline in harvest occurred on  
28 Federal lands with harvest levels decreasing by 90 percent, from 833 million board feet in  
29 1990 to just 85 million board feet in 2002. Statewide, private lands accounted for the  
30 majority of harvest in 2002, with forest industry lands providing 36 percent of total harvest  
31 and large private and small private lands accounting for 27 percent and 11 percent,  
32 respectively (Washington DNR 2004b).

33 Counties in western Washington accounted for approximately 75 percent of total harvest in  
34 2002 and the majority of harvest on forest industry, private large, private small, and State  
35 lands (Table 3-31). This pattern was, however, reversed for tribal and Federal lands with

36

---


<sup>1</sup> Covered employment includes workers covered by State unemployment insurance laws and Federal workers covered by the Unemployment Compensation for Federal Employees Program. Self-employed persons are not included in covered employment data. This differs from the data summarized in Table 3-29, which include both covered employment and self-employed persons. Covered employment data tend to underestimate total employment in the logging, commercial fishing, and recreation and tourism sectors because people who work in these industries are often self-employed.



## Chapter 3

1

2 **Figure 3-10.** Washington Timber Harvest by Owner, 1990 to 2002.



3 Notes:

4 MMBF = million board feet

5 Private Large—Non-industrial companies and individuals not operating wood-using plants but with statewide holdings totaling more than 1,000 acres.

6 Private Small—Non-industrial companies and individuals not operating wood-using plants and with statewide holdings totaling less than 1,000 acres.

7 Data for the Other Non-Federal category are not shown in this figure. Statewide harvest from Other Non-Federal lands accounted for 26 million board feet in 2001.

8 Source: Washington DNR Washington Timber Harvest, annual reports 1990 to 2002.

12

13 **Table 3-31.** Timber Harvest by Ownership and Location, 2001.

	Tribal	Forest Industry	Private Large	Private Small	State	Federal	Other Non-Federal	Total
<b>Harvest (MMBF)</b>								
Eastside	292	163	172	113	59	66	14	878
Westside	27	1,142	795	297	398	19	26	2,704
<b>Total</b>	<b>319</b>	<b>1,304</b>	<b>967</b>	<b>410</b>	<b>457</b>	<b>85</b>	<b>40</b>	<b>3,582</b>
<b>Percent of Total by Ownership</b>								
Eastside	92	12	18	28	13	78	34	25
Westside	8	88	82	72	87	22	66	75
<b>Total</b>	<b>100</b>	<b>100</b>	<b>100</b>	<b>100</b>	<b>100</b>	<b>100</b>	<b>100</b>	<b>100</b>

MMBF = million board feet

Private Large—Non-industrial companies and individuals not operating wood-using plants but with statewide holdings totaling more than 1,000 acres.

Private Small—Non-industrial companies and individuals not operating wood-using plants with statewide holdings totaling less than 1,000 acres.

Source: Washington DNR 2004b.





1 counties in eastern Washington accounting for the majority of timber harvest on these land  
2 ownerships (Washington DNR 2004b).

### 3 **Commercial Fishing**

4 The commercial fishing industry accounted for 9,418 jobs in Washington State in 2000,  
5 less than 1 percent (0.3 percent) of total employment (Table 3-30). Employment in this  
6 sector was 24 percent lower in 2000 than it was in 1990 (Table 3-30). Covered  
7 employment in the fishing, hunting, and trapping sector accounted for less than 1 percent  
8 of total employment in all counties in 2001, with the exception of Pacific County where it  
9 comprised 5.2 percent of all covered employment. Viewed in absolute terms, the counties  
10 with the largest number of covered jobs in this sector in 2001 were King (1,122 jobs) and  
11 Pacific (313 jobs) (Washington State Employment Security Department 2004). Covered  
12 employment and wage data indicate that the average wage in the fishing, hunting, and  
13 trapping sector in Washington State was \$55,250 in 2001 (Washington State Employment  
14 Security Department 2004). Data compiled by the Pacific Fishery Management Council  
15 suggest that the estimated economic value of commercial fishing to Washington state was  
16 \$289.2 million in 2001 (Washington Department of Fish and Wildlife 2001c).

17 As the above employment data suggest, ports on the Puget Sound and the Pacific Ocean  
18 handle almost all commercial landings in Washington, with less than 1 percent of total  
19 catch by value coming from freshwater harvest. Salmon account for about one-third of the  
20 catch by value, followed by oysters, crab, shrimp, and other shellfish. Other fish caught  
21 include halibut, flounder, tuna, cod, rockfish, pollock, and sablefish (Washington DNR  
22 2004b).

23 Sport and tribal fishing also generate employment and income in Washington. Sport  
24 fishing is addressed in the following Recreation and Tourism section. Tribal issues,  
25 including ceremonial and subsistence harvest, are addressed in subsection 3.13, Cultural  
26 and Indian Trust Resources.

### 27 **Recreation and Tourism**

28 Recreation and tourism is not classified or measured as a standard industry category and,  
29 therefore, employment and income data are not specifically collected for this sector.  
30 Components of recreation and tourism activities are instead captured in other industrial  
31 sectors, primarily the retail sales and services sectors. Estimates of travel impacts  
32 developed for the Washington State Office of Trade and Economic Development indicated  
33 that travel-related expenditures supported approximately 160,720 jobs in Washington State  
34 in 2000, representing approximately 4.5 percent of total State employment. Travel-related  
35 employment ranged from below 3.5 percent of total employment in Clark, Thurston,  
36 Wahkiakum, and Yakima Counties to slightly more than 20 percent in Pacific (20.1  
37 percent) and Skamania (21.6 percent) Counties (Dean Runyon Associates 2002).

38 Employment in the recreation and tourism sector generally tends to be seasonal and  
39 relatively low paid, with a high proportion of the labor force self-employed. The study  
40 prepared for Washington State indicated that the average annual salary for this sector in  
41 2001 was \$20,604 compared to the State average salary of \$37,849 for all sectors (Dean



## Chapter 3

1 Runyon Associates 2002; U.S. Bureau of Economic Analysis 2004b). Average salaries in  
 2 2001 ranged from \$22,252 in Pacific County to \$47,336 in King County. The majority of  
 3 counties (31 out of 38) had average salaries of \$30,000 or below in 2001 (U.S. Bureau of  
 4 Economic Analysis 2004b).

5 The 2001 National Survey of Fishing, Hunting, and Wildlife-Associated Recreation  
 6 estimated that a total of 938,000 residents and nonresidents 16 years or older fished in  
 7 Washington in 2001, spending approximately \$854 million on fishing-related expenses  
 8 (Table 3-32) (U.S. Fish and Wildlife and Census Bureau 2003). The survey identified  
 9 approximately 659,000 freshwater anglers, with 211,000 and 156,000 anglers indicating  
 10 that they fished for salmon and steelhead, respectively. The survey also identified 386,000  
 11 saltwater anglers, with 250,000 anglers indicating that they fished for salmon in 2001 (U.S.  
 12 Fish and Wildlife and Census Bureau 2003). These categories are not mutually exclusive.  
 13 Some anglers fish in both fresh and salt water, and the majority fish for more than one  
 14 species at any one time. These numbers do, however, provide a good indication of the  
 15 importance of recreational fishing in Washington State, as well as the importance of  
 16 salmon and steelhead fishing to this activity.

17 This survey also found that a total of 227,000 residents and nonresidents 16 years or older  
 18 hunted in Washington in 2001, spending approximately \$350 million on hunting-related  
 19 expenditures (Table 3-31). In addition, 2.5 million residents and nonresidents 16 years or  
 20 older fed, observed, or photographed wildlife in Washington, spending approximately \$980  
 21 million on wildlife-watching activities (U.S. Fish and Wildlife and Census Bureau 2003).

22 **Table 3-32.** Fishing, Hunting, and Wildlife-Associated Recreation Expenditures in  
 23 Washington in 2001.

Activity	Number of Participants	Estimated Expenditures (\$ thousand)
Recreational Fishing	947,000	853,761
Hunting	269,000	349,771
Wildlife Viewing	2,496,000	979,730

Source: U.S. Fish and Wildlife and Census Bureau 2003.

### 24 3.14.3.4 Income and Poverty

25 Per capita income, which is calculated by dividing total personal income by total  
 26 population, was \$31,230 in Washington State in 2000, ranging from \$16,600 or 53 percent  
 27 of the State average in Ferry County to \$45,500 (146 percent of the State average) in King  
 28 County. San Juan County was the only other county with per capita income above the  
 29 State average (\$33,800 or 115 percent) (Washington State Employment Security  
 30 Department 2004).

31 Total personal income includes earnings (wage and salary disbursements, other labor  
 32 income, and proprietors' income); dividends, interest, and rent; and transfer payments  
 33 received by residents. Earnings accounted for approximately 68 percent of total personal  
 34 income in Washington State in 2001; dividends, interest, and rent comprised approximately  
 35 19 percent, and transfer payments accounted for approximately 13 percent. Earnings,  
 36 dividends, interest, and rent, and transfer payments also accounted for the same relative



1 shares of total personal income in Washington State in 1991 (U.S. Bureau of Economic  
2 Analysis 2004c). The share of total personal income comprised of earnings varied by  
3 county, ranging from 46 percent in Clallam County in 2000 to 74 percent in Snohomish  
4 and King Counties.

5 In 1999, 10.6 percent of the population in Washington was below the poverty rate,  
6 compared to 10.9 percent in 1989. The percent of the population below the poverty rate by  
7 county ranged from 6.9 percent in Snohomish County to more than 20 percent in  
8 Okanogan (21.5 percent) and Whitman (24.2 percent) Counties (USDA Economic  
9 Research Service 2004).

#### 10 **3.14.4 Environmental Justice**

11 Executive Order 12898, Federal Actions to Address Environmental Justice in Minority  
12 Populations and Low-Income Populations, requires each Federal agency to make the  
13 achievement of environmental justice part of its mission by identifying and addressing  
14 disproportionately high and adverse human health or environmental effects of its programs,  
15 policies, and activities on minority and low income populations. The Order further  
16 stipulates that the agencies conduct their programs and activities in a manner that does not  
17 have the effect of excluding persons from participation in, denying persons the benefits of,  
18 or subjecting persons to discrimination because of their race, color, or national origin.

19 The EPA working with the Enforcement Subcommittee of the National Environmental  
20 Justice Advisory Council has developed technical guidance for conducting environmental  
21 justice assessments that are referenced in the following discussion, as appropriate. Much  
22 of this guidance is concerned with identifying low income and minority populations based  
23 on the location of the proposed action. Suggested measures include identifying affected  
24 areas as low income if more than 20 percent of the affected area is below the poverty level  
25 or identifying areas as minority areas if minority populations represent more than 15.72  
26 percent of the total population. These types of measure are useful for identifying potential  
27 environmental justice concerns associated with proposed actions that occur in specific  
28 locations, such as the siting of a hazardous waste site or an electric transmission line. They  
29 have very limited or no applicability to analyses such as this that are concerned with  
30 programmatic actions that establish direction for broad land areas rather than scheduling  
31 activities on specific parcels of land. Race, ethnicity, income, and poverty are, however,  
32 discussed for the State and by county in subsections 3.14.2.1 (Race and Ethnicity) and  
33 3.14.3.4 (Income and Poverty).

34 More relevant to this analysis are measures that assess target populations in terms of those  
35 groups that would likely be disproportionately affected by the proposed action. In this  
36 case, these groups include loggers, mill workers, and others involved in timber harvest, and  
37 groups that would be affected by potential changes in salmonid populations, primarily  
38 people involved in commercial and recreational fishing.

39 Using this approach the minority populations most likely to be affected by the proposed  
40 action are Native American Tribes. The general relationship between the potentially  
41 affected Tribes, those that presently reside in Washington, as well as those that once lived



## **Chapter 3**

---

1 in what is now Washington State and now reside in adjacent states and Canada, is  
2 discussed in subsection 3.13 (Cultural and Indian Trust Resources). The federally  
3 recognized Tribes of Washington and adjacent states with cultural interests in Washington  
4 forests are identified in Table 3-27.

5 Plant and animal resources that are culturally important for Native American Tribes in  
6 Washington are summarized in Table 3-28. These include salmon, which are important  
7 traditional resources to all Washington Tribes, and bull trout, which are particularly  
8 important in watersheds outside the salmon's historic range. In addition to being important  
9 traditional tribal resources, salmon and bull trout are also important for subsistence and  
10 commercial fishing. Forests are also important resources for Washington's Tribes  
11 providing raw materials, such as bark and grasses for baskets and mats, wood for carvings,  
12 and medicinal plants, as well as subsistence resources.