

DEPARTMENT OF NATURAL RESOURCES

SOUTH PUGET SOUND REGION

950 Farman Avenue North Enumclaw, WA 98022

360-825-1631SOUTHPUGET.REGION@DNR.WA.GOV
WWW.DNR.WA.GOV

October 31, 2024

Notice of Final Determination Evergreen Gold Timber Sale #30-103585 SEPA File No. 24-101602

The Department of Natural Resources issued a [x] Determination of Non-significance (DNS), [] Mitigated Determination of Non-significance (MDNS), [] Modified DNS/MDNS on October 16, 2024 for this proposal under the State Environmental Policy Act (SEPA) and WAC 197-11-340(2).
This threshold determination is hereby:
[] Retained.
[x] Modified. Modifications to this threshold determination include the following:
It was discovered the SEPA checklist for Evergreen Gold did not fully address Proposed landscaping, use of native plants, or other measures to preserve or enhance vegetation on the site. The answer to B.4.d. should include the following: Trees over 60 inches diameter at breast height (DBH), even if not marked for leave, will be retained through contract requirements.
Following harvest, the variable retention harvest units will be replanted with native conifer species that will be supplemented by natural regeneration which is expected to occur as a result of the conservation areas in and around the harvest units. Species, stocking type, and density for plantings are prescribed to be suitable for the unique site conditions. After planting, the need for noxious weed treatments will be assessed and occur if necessary.
[] Withdrawn. This threshold determination has been withdrawn due to the following:
[] Delayed. A final threshold determination has been delayed due to the following:

Summary of Comments and Responses (if applicable):

Comment: Received October 17, 2024 from an Olympia resident concerning harvesting impacts on climate change and recreation, as well as negative public use impacts.

Response: DNR provided a response to the commenter on October 31, 2024 regarding DNR management.

Comment: Received October 23, 2024 from the advocacy organization Legacy Forest Defense Coalition concerning structurally complex stands and DNR's 10 to 15 percent older forest targets.



DEPARTMENT OF NATURAL RESOURCES

SOUTH PUGET SOUND REGION

950 Farman Avenue North Enumclaw, WA 98022

360-825-1631SOUTHPUGET.REGION@DNR.WA.GOV
WWW.DNR.WA.GOV

Response: DNR provided a response to the commenter on October 31, 2024 regarding the Department's older forest management. The 'Landscape Assessment to Identify and Manage Structurally Complex Stands to Meet Older-Forest Targets in Western Washington', May 2024 (Revised September 2024), referenced in Class Dismissed SEPA checklist A.8. was also provided to this commenter and attached for reference.

Responsible Official: Don Melton

Position/title: South Puget Sound Region Manager Phone: (360) 825-1631

Address: 950 Farman Avenue North, Enumclaw, WA 98022

Date: 10/31/2024 Signature: Don Welton A

There is no DNR administrative SEPA appeal.

Landscape Assessment to Identify and Manage Structurally Complex Stands to Meet Older-Forest Targets in Western Washington

Washington State Department of Natural Resources | May 7, 2024

Revised September 3, 2024¹

Summary

The Washington Department of Natural Resources (DNR) actively manages suitable forest stands to achieve, over time, older-forest structures across 10 to 15 percent of each 1997 *State Trust Lands Habitat Conservation Plan* (HCP) planning unit in western Washington. These "older-forest targets" are described in DNR's *Policy for Sustainable Forests* (PSF).

This document (2024 Landscape Assessment) provides updated projections on the amount of older-forest stands, as defined by the PSF and discussed in this document, that will develop in western Washington by 2100 as a result of implementing landscape-level management strategies. It describes how DNR has implemented these landscape-level management strategies at the HCP planning unit scale, and how DNR has identified and prioritized suitable stands to be managed to meet older-forest targets.

This document is an update to a landscape assessment titled "Identifying Stands to Meet Older-Forest Targets in Western Washington" dated May 11, 2021 (2021 Landscape Assessment). This 2024 Landscape Assessment:

- Utilizes the same methodologies and data sources as the 2021 Landscape Assessment, which are discussed in Part 2 of this document.
- Refines and updates conservation areas from the 2021 Landscape Assessment, including the
 addition of newly designated areas, to better identify areas deferred from stand replacement
 harvest. Conservation areas include areas being conserved under the HCP; Natural Areas such as
 Natural Resource Conservation Areas and Natural Area Preserves; areas deferred by the PSF; and
 areas conserved under the Forest Practices Act.
- Updates the results tables from the 2021 Landscape Assessment that detail DNR's projected timeline for reaching the PSF's older-forest targets to reflect the refined and updated conservation areas.

As with the 2021 Assessment, DNR identified and designated the forest stands that will contribute to older-forest targets within its GIS spatial database. The identified stands are also illustrated on the maps in Appendix 2. The identified stands comprise some existing, suitable structurally complex stands, and

_

¹ Refer to Appendix 4 for data revised on September 3, 2024.

additional stands suitable to be managed for older-forest targets over time. The identified stands are all located within conservation areas.

Part 1: Overview of DNR's Management Framework

Development of older-forest structures is an expected outcome of DNR's management under the HCP and a policy objective of the PSF. DNR also analyzed stand structure as part of its consideration of alternatives for the 2004 Sustainable Harvest Level calculation. For more information on older-forest structure, refer to Part 2, Identifying Older-Forest Stands.

Habitat Conservation Plan

The HCP mentions stand structure in the Multispecies Conservation Strategy for Unlisted Species. Specifically, the HCP discusses the likely outcomes of DNR's management under the HCP:

DNR has modeled the age-class distribution that will likely result from expected management under the HCP and existing policies. Results from this modeling have been used to develop a table (refer to HCP Table IV.14) of ranges of expected percentages of each of several forest habitat/structural types, using age-class as a surrogate, that would likely exist 100 years following implementation of such management (HCP pg. IV.179).

The HCP uses age classes as a surrogate for stand structure. Specifically, it utilizes an age of 150 years in the five west-side planning units, and 200 years in the Olympic Experimental State Forest (OESF) as representative of a stand in the fully functional stand stage. In the HCP's analysis, the modeling projected that forest stands in the fully functional stand stage would occur on approximately 10 to 15 percent of the five west-side planning units and the OESF after 100 years of HCP implementation (in other words, by 2097).

Sustainable Harvest Calculation

The 2004 Final Environmental Impact Statement on Alternatives for Sustainable Forest Management of State Trust Lands in Western Washington and for Determining the Sustainable Harvest Level (2004 Sustainable Harvest FEIS) includes an analysis of forest stands over time using eight stand development stages. (Refer to Appendix 1 of this document for forest inventory queries used to identify stand development stages in the 2004 Sustainable Harvest FEIS.) The Board of Natural Resources adopted its preferred alternative, which includes a proposed policy change, in Resolution 1110:

The proposed Preferred Alternative would target 10 to 15 percent of each westside HCP Planning Unit as older forests based on structural characteristics. The desired structural characteristics are represented by stand development classes understory development through old natural forests (as described in the Sustainable Forest Management DEIS)." (Resolution 1110, pg. 3 of Document 2). The 2004 Sustainable Harvest FEIS defined structurally complex stands as those stands in the botanically diverse, niche diversification, and fully functional stages of stand development (2004 Sustainable Harvest FEIS, pg. 4-22).

The FEIS further indicates that the preferred alternative includes a proposed policy to "Manage 10-15% of each Planning Unit in Mature Forest Component" (2004 Sustainable Harvest FEIS, pg. 4-23 Table 4.2-12). The proposed policy was analyzed within the 2004 Sustainable Harvest FEIS, and then later adopted with the then-anticipated PSF (which was an update to the 1992 *Forest Resources Plan*):

The policies amended through the Final Environmental Impact Statement on Alternatives for Sustainable Forest Management of State Trust Lands in Western Washington (2004) have already been analyzed and adopted by the Board of Natural Resources and will be included in the PSF" (Final Environmental Impact Statement on the Policy for Sustainable Forests [PSF FEIS] pg. 1-3).

Policy for Sustainable Forests

The Board of Natural Resources adopted the PSF in 2006. The PSF contains a suite of landscape-level management strategies, including the General Silvicultural Strategy, which adopted the proposed policy analyzed in the 2004 Sustainable Harvest FEIS (PSF pg. 46). The PSF states that "DNR intends to actively manage suitable structurally complex forests to achieve older-forest structures across 10 to 15 percent of each western Washington HCP planning unit in 70 to 100 years." Meeting older-forest targets within 70 to 100 years equates to target dates of 2076 to 2106. The PSF emphasized that DNR anticipates reaching the older-forest targets over time. The PSF, relying on the analysis in the 2004 Sustainable Harvest FEIS (pg. 3-44), states, "Older-forest structures that contribute to this target are represented by stands in the fully functional or niche diversification stage of stand development." It also states that, "Through landscape assessments, the department will identify suitable structurally complex forest stands to be managed to help meet older-forest targets" (PSF pg. 47).

DNR will identify stands to be managed to achieve older-forest structures in conservation areas. As noted earlier, conservation areas include Natural Areas; areas conserved under the HCP, areas deferred from harvest by the PSF, and areas conserved under the Forest Practices Act.

Part 2. Identifying Older-Forest Stands

Definitions and Query Criteria

In the 1990s and 2000s, several research teams developed and published definitions of stand development stages for Pacific Northwest forests (for example, Carey and Curtis 1996, Oliver and Larson 1996, and Franklin and others 2002, refer to references at the end of this document). These definitions are also used in the description of the policy in the PSF FEIS.

Older-forest stands are defined as those in the "niche diversification" or "fully functional" stand-development stages (PSF FEIS pg. 3-177). However, the published literature did not provide guidance about how to identify these stands based on forest inventory data.

To address this issue, DNR developed stand-level variables and associated threshold values that can be applied to DNR's inventory data to identify forests in these two stand development stages. The variables were based on the type of inventory data that DNR collects. This work was done as part of the 2004

Sustainable Harvest FEIS and was based on scientific literature. Using these variables and threshold values, DNR was able to identify stands in these stand development stages across the landscape. Table B.2.2-2 of the 2004 Sustainable Harvest FEIS provides the stand-development stages and representative stand-level variables and associated threshold values used in the project and is also included in Appendix 1 of this document.

DNR conducted this 2024 Landscape Assessment to monitor forest conditions existing on the landscape. For this work, DNR used the definitions of stand development stages adopted in the PSF and queried them using the identified, representative stand-level variables and associated threshold values as shown in Appendix 1.

DNR did not use descriptions or methods of identifying these forests that were produced after the 2004 Sustainable Harvest FEIS. For example, *Identifying Mature and Old Forests in Western Washington* (Van Pelt 2007), and environmental impacts statements for the *South Puget Forest Land Plan* (DNR 2010), *OESF Forest Land Plan* (DNR 2016), and the 2019 sustainable harvest level (DNR 2019) used different methods to classify stand development and older-forest stands for the purposes of those analyses. Van Pelt (2007) also provides a method to identify older-forest stands and trees in the field. The descriptions and analyses in these documents complement but do not redefine the direction set in the PSF or any of the strategies or policies contained in the PSF.

Identification of Suitable Stands

The HCP projected that the 10 to 15 percent older-forest structure target will likely be achieved over time through implementation of its conservation strategies (HCP IV.179). The HCP's projection is based on the premise that fully functional structure is likely to develop in riparian zones, wildlife habitat, and other areas conserved by the HCP. The PSF directs DNR to manage suitable stands to meet older-forest targets, as defined in the PSF (PSF pg. 46). The PSF notes that "the size of the stand, its proximity to old growth or other structurally complex forest stands, or the scarcity of old growth or other structurally complex forest stands are all factors in determining whether a stand is suitable for contributing to older-forest targets" (PSF pg. 46).

DNR manages approximately 1.6 million acres of forested land in western Washington under the HCP. Approximately 681,000 acres (43 percent) of that land is permanently deferred from harvest by the HCP or PSF or has been incorporated into a Natural Area. These conserved areas provide good connectivity with other conserved and federal land, may be centered around old growth, provide optimal wildlife habitat, and contain the majority of older-forest structure on DNR-managed lands. Based on these factors, DNR has determined that conserved areas are the most suitable places to manage for older-forest targets.

While some of the forest stands DNR has identified to be managed for older-forest structure are not currently structurally complex, DNR anticipates that all identified stands will reach the niche diversification or fully functional stand development stage between 2076 and 2106. Stands within conserved areas that are capable of achieving older-forest structure within the anticipated time frame were prioritized over existing structurally complex stands in general ecological management areas (areas that are generally available for harvest) because their heightened connectivity to existing older forest make them more suitable to be managed to meet the older-forest targets.

Analysis

Following is an updated analysis that was completed by applying the stand-development stage stand-level variables and associated threshold values from the 2004 sustainable harvest FEIS to DNR's inventory data from 2021. The methods in this 2024 Landscape Assessment are the same as those used in the 2021 Landscape Assessment, with modifications to the conservation areas.

Methods

In both the 2021 and 2024 landscape assessments, DNR developed queries based on the stand-level variables and associated threshold values defined in the 2004 Sustainable Harvest FEIS. These queries were applied to DNR's forest inventory data and allowed DNR to identify stands (including older-forest stands) across the landscape.

The stand-level variables and associated threshold values in 2004 Sustainable Harvest FEIS were designed for use with DNR's plot-based inventory. In 2017, DNR changed the inventory system from a fully plot-based system to a system called RS-FRIS that pairs plots with remote sensing data. Due to this change in methods, the data are not structured in the same manner. As a result, DNR modified the data queries to be compatible with RS-FRIS Version 3.

Conserved Areas

For the 2021 Landscape Assessment, DNR identified conserved areas using the large data overlay. Conserved areas included Natural Areas; areas conserved under the HCP, including long-term forest cover designated under the marbled murrelet long-term conservation strategy, riparian areas, areas conserved under the multispecies conservation strategy, potentially unstable slopes, and spotted owl nest patches; areas deferred from harvest by the PSF, including old-growth forest; and areas conserved under the Forest Practices Act.

In preparing this 2024 Landscape Assessment, DNR included recently designated Natural Areas and newly identified old-growth forest in the delineated conserved areas. Further, DNR added 2,000 acres of forests designated to be set aside to the conserved areas under the fiscal year 2023-25 Capital Budget (Chapter 474, Laws of 2023, Section 3130). Lastly, DNR removed areas that are not permanently deferred, such as suitable northern spotted owl habitat that will be available for harvest once habitat thresholds are met, from the delineated conserved areas.

This 2024 Landscape Assessment confirms that DNR is on track to achieve the PSF's older-forest targets through implementation of the HCP and other conservation strategies within 100 years, by prioritizing and designating suitable structurally complex stands within conservation areas. Accordingly, DNR has not included areas outside of conservation areas to be managed to reach the PSF's older-forest targets.

Key Metrics

Max RD Age

A key starting point for identifying older-forest stands is "max RD age." This is the age at which unthinned stands reach their maximum relative density (RD)². DNR used yield curves³ developed for the 2019 sustainable harvest calculation to determine max RD age. These yield curves are produced by the Forest Vegetation Simulator (Dixon 2002) and adjusted to fit growing conditions on DNR-managed lands in Western Washington, based on a comparison with inventory plot data. DNR has yield curves for three cover types (Douglas-fir, red alder, and western hemlock) and four site index classes⁴.

In this analysis, DNR used the max RD age for each of these cover types and site index classes. Since stands in Western Washington tend to stay at or near max RD age for many decades, DNR defined max RD age as the point at which the RD yield curve levels off. The 2004 Sustainable Harvest FEIS defined the niche diversification stage, in part, as stands at least 80 years older than the max RD age, and the fully functional stage as stands at least 160 years older than the max RD age.

Table 1. Age of max RD for the three forest cover types and four site index classes on DNR managed land (refer to DNR 2019 Appendix F for more details).

Cover type	Site index class	Max RD age
Douglas fir	1	43
Douglas fir	2	43
Douglas fir	3	44
Douglas fir	4	47
Red alder	1	44
Red alder	2	43
Red alder	3	43
Red alder	4	44
Western hemlock	1	46
Western hemlock	2	44
Western hemlock	3	48
Western hemlock	4	50

Canopy Layers

Both the niche diversification and fully functional stand development stages can be defined in part by the presence of more than one canopy layer. Figure 1 shows the average number and variability of canopy


² Relative density (RD) is a quantification of the current density of a forest stand in comparison to a maximum level.

³ A yield curve projects the volume growth of a forest over time.

⁴Site index describes the potential for trees to grow at a particular location.


layers in stands of different ages. By about age 80, stands average over 2 canopy layers, with 90 percent confidence intervals above 2 layers.

Figure 1. Canopy layers by age from DNR inventory plots. Each box represents the average of 30 plots. Lines show 90 percent confidence intervals.



Legacy Structures

The niche diversification and fully functional stand development stages include biological legacies in the form of snags and woody debris. DNR's data show that stands roughly 90 years old and older average more than 3 snags per acre that are over 20 inches in diameter, with 90 percent confidence intervals extending only slightly below 3 snags per acre (Figure 2). Likewise, stands over 115 years old average more than 2,400 cubic feet of dead and down woody debris per acre (Figure 3).







Figure 3. Cubic feet of dead and down wood material (DDWM) by age from DNR inventory plots. Each box represents the average of 30 plots. Lines show 90 percent confidence intervals.



Plot Data

Figures 2 and 3 show that aggregated RS-FRIS inventory plots in stands averaging over 115 years old have the characteristics of the niche diversification stand development stage. However, at the stand level, not all characteristics are present in all stands.

DNR used data from its older, field-plot-based inventory system, FRIS, to further analyze stand-level conditions. As stated previously, the HCP uses age classes as a surrogate for stand structure. Specifically, the HCP utilizes an age of 150 years in the five west-side HCP planning units, and 200 years in the OESF, to represent a fully functional stand development stage. DNR calculated the area of each HCP planning unit that contained stands at least five acres in size that were over 150 years old (200 years old in the OESF). Sixty-eight percent of those stands with at least 5 FRIS plots that were identified as older-forest based on age alone contained the legacy components of the older-forest stand definition. To reflect this information, DNR reduced the raw projections displayed in Table 3 by 32 percent, which is termed a plot discount. These discounts are shown in Table 2.

This current analysis made no adjustment for the time since the collection of FRIS plot data, some of which is over 20 years old. Therefore, this result is likely an underestimate of older-forest conditions at the stand level.

Disturbance

For this analysis, DNR assumed that all stands, including older-forest stands, will be subject to stand-replacing disturbances over time. As a result, not all areas projected to develop older-forest structure based on growth alone will actually achieve that structure within 100 years. In 2016, Raphael and others completed a study looking at stands that provide higher-quality marbled murrelet habitat, which is generally similar to older-forest conditions. They reported a decadal disturbance rate of 0.43 percent. DNR then assumed that disturbance rates in marbled murrelet habitat would increase by 20 percent per decade, resulting in doubling of the decadal disturbance rate over 5 decades. The marbled murrelet long-term conservation strategy disturbance rate assumption was applied across this analysis. DNR refers to this rate as a disturbance factor.

Active Management

This analysis does not project the acceleration of stand development stages due to forest management. Published literature, such as Carey and Curtis (1996), indicate that active management can advance stands through stand development stages. The PSF also envisioned active management contributing to the development of older-forest structures. DNR's management in certain conservation areas is designed to accelerate older-forest structure development. For example, the goal of the Riparian Forest Restoration Strategy (RFRS) is to hasten the development of older-forest structures in riparian areas: "The main objective of silvicultural activities will be to put the Riparian Management Zone on an accelerated trajectory toward the riparian desired future conditions" (DNR 2006c, pg. 20). Based on modeling for the 2019 Sustainable Harvest FEIS, DNR anticipates over 4,000 acres of thinning will occur in riparian areas per the RFRS between 2019 and 2067, the end of the initial HCP agreement period. Thinned stands are


likely to develop older-forest structures faster than if left untreated. As this acceleration is not accounted for in this analysis, the query may underestimate the actual area achieving older-forest structures.

Identification of Stands

DNR maintains age data in both polygon and raster format. For this 2024 Landscape Assessment, DNR used the raster format to avoid misidentifying a stand's development stage based on polygons that do not precisely match stand boundaries. For this analysis, DNR used the same raster aggregation method that it uses to delineate northern spotted owl habitat.

DNR used the combined origin year raster in RS-FRIS as the source for age data. This data source shows age for all DNR-managed lands in western Washington in a 0.1-acre grid. In the delineation process, DNR calculated the median age of each cell by utilizing the age of the cell and the adjacent eight cells, and then assigned the median age to that cell. The resulting cells over a specified age are then converted to polygons, with adjacent cells dissolved into a single, larger polygon. Further processing is then done to fill gaps 132 feet across (two chains or double the width of an RS-FRIS cell), and to remove patch portions that are narrower than 132 feet. DNR then removed any stands under 5 acres identified in this process. As a result, only older-forest stands larger than 5 acres are counted towards the estimated area of older-forest stands in a given decade. DNR repeated this process for 10 decades, projecting the age of each cell forward to future years. This process creates smoothed, contiguous stands from the pixelated original data that DNR used, to estimate the area of older-forest stands in conservation areas into the future (Figure 4).

Figure 4. An example of spatial arrangement of input data for and results of the stand identification process converting RS-FRIS grid data into stand polygons.



Part 3: Results

Older-Forest Stands

The following tables contain the results of DNR's 2024 Landscape Assessment, utilizing the methodologies and queries described in Part 2 with updated conservation areas. As illustrated in Table 2, through implementation of the HCP and other policies and laws, over 10 percent of the OESF HCP Planning Unit currently contains older-forest stands. The North Puget HCP Planning Unit is on track to achieve 10 percent older-forest stands by 2070; South Puget, Columbia, and Straits are on track to achieve 10 percent older-forest stands by 2090; and South Coast is on track to achieve 10 percent older-forest stands by 2100. (Table 2). **DNR will meet the PSF's older-forest targets before 2106.**

Table 2⁵: Percent area western Washington HCP planning units with older-forest stands in conservation areas by decade through 2120. Adjusted query output with plot discounts and disturbance factor.

HCP Planning		Year												
Unit	2021	2030	2040	2050	2060	2070	2080	2090	2100	2110	2120			
Columbia	1.1%	1.2%	1.4%	1.8%	2.6%	4.3%	6.8%	10.1%	14.0%	17.3%	18.9%			
North Puget	3.2%	3.9%	4.9%	6.2%	7.9%	10.2%	13.2%	16.7%	20.6%	23.9%	25.0%			
OESF	10.2%	10.7%	11.0%	11.7%	12.6%	13.9%	16.0%	20.1%	25.0%	28.4%	29.6%			
South Coast	0.2%	0.3%	0.6%	1.2%	2.2%	3.6%	6.0%	8.8%	12.3%	16.0%	18.7%			
South Puget	1.7%	2.1%	2.7%	3.6%	4.6%	6.1%	8.4%	11.3%	14.4%	17.2%	18.7%			
Straits	1.8%	2.5%	3.2%	4.3%	5.6%	7.4%	9.9%	12.6%	15.0%	17.9%	19.3%			

Table 2 represents the most conservative time estimate of DNR meeting older-forest targets. The plot discount removed query results that were determined to be "false positives;" however, it is likely that the query also produced "false negatives." In other words, the query likely eliminated stands that did not meet certain criteria from the inventory data, but do, in fact, exhibit older-forest structure. There is no way at this time to determine the rate of false negatives and so DNR did not account for them. Table 2 is the most conservative estimate of projected older-forest stands across the westside HCP Planning Units and the assessments show that DNR will meet older-forest targets in 70 to 100 years. Table 3 illustrates a less-conservative time estimate by eliminating the plot discount and disturbance factor. Refer to Appendix 3 for maps of those stands that are referenced in Table 3, year 2100.

_

⁵ Refer to Appendix 4 for data revised on September 3, 2024.

Table 3⁶. Percent area Western Washington HCP planning units with older-forest stands in conservation areas by decade through 2120. Values over 10% in bold. Raw query output without plot discount or disturbance factor.

HCP Planning	Year												
Unit	2021	2030	2040	2050	2060	2070	2080	2090	2100	2110	2120		
Columbia	1.5%	1.7%	2.0%	2.6%	3.9%	6.5%	10.5%	15.8%	22.1%	27.8%	31.1%		
North Puget	4.7%	5.8%	7.2%	9.2%	11.9%	15.6%	20.3%	26.1%	32.5%	38.5%	41.0%		
OESF	14.9%	15.7%	16.3%	17.5%	18.9%	21.2%	24.5%	31.3%	39.6%	45.6%	48.6%		
South Coast	0.3%	0.5%	1.0%	1.8%	3.2%	5.5%	9.2%	13.8%	19.5%	25.8%	30.7%		
South Puget	2.5%	3.2%	4.0%	5.3%	6.9%	9.2%	12.9%	17.7%	22.8%	27.6%	30.7%		
Straits	2.7%	3.7%	4.8%	6.3%	8.4%	11.2%	15.2%	19.6%	23.8%	28.8%	31.7%		

⁶ Refer to Appendix 4 for data revised on September 3, 2024.

References

Carey, A.B. and R.O. Curtis. 1996. Conservation of biodiversity: a useful paradigm for forest ecosystem management. Wildlife Society Bulletin 1996, 24(2): 610-620.

Dixon, G.E. (Comp.), 2002. Essential FVS: A User's Guide to the Forest Vegetation Simulator. Internal Report. U.S. Department of Agriculture, Forest Service, Forest Management Service Center, Fort Collins, CO, 189 pp.

DNR: refer to Washington Department of Natural Resources.

Franklin, J. F., T. A. Spies, R. Van Pelt, A. B. Carey, D. A. Thornburgh, D. R. Berg, D. B. Lindenmayer, M. E. Harmon, W. S. Keeton, D. C. Shaw, K. Bible, and J. Chen. 2002. Disturbances and Structural Development of Natural Forest Ecosystems with Silvicultural Implications, Using Douglas-fir Forests as an Example. Forest Ecology and Management 155:399–423. Oliver, C. D. and B. C. Larson. 1996. Forest Stand Dynamics, update edition. John Wiley & Sons, New York, New York. 520 p.

Raphael, M.G., G.A. Falxa, D. Lynch, S.K. Nelson, S.F. Pearson, A.J. Shirk, and R.D. Young. 2016. Status and trend of nesting habitat for the marbled murrelet under the Northwest Forest Plan. Chapter 2, in Falxa, G.A. and M.G. Raphael (tech. eds.), 2016: Northwest Forest Plan—the first 20 years (19942013): Status and trend of Marbled Murrelet populations and nesting habitat. Gen.Tech. Rep. PNW-GTR933. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 132 p

Van Pelt, R. 2007. Identifying Mature and Old Forests in Western Washington. Washington State Department of Natural Resources, Olympia, WA. 104 p.

Washington Department of Natural Resources. 1997. Habitat Conservation Plan. Washington State Department of Natural Resources, Olympia, Washington.

Washington Department of Natural Resources. 2004. Final Environmental Impact Statement on Alternatives for Sustainable Forest Management of State Lands in Western Washington and for Determining the Sustainable Harvest Level. Washington Department of Natural Resources, Olympia, Washington.

Washington Department of Natural Resources. 2005. Draft Environmental Impact Statement on the Policy for Sustainable Forests. Washington State Department of Natural Resources, Olympia, Washington.

Washington Department of Natural Resources. 2006a. Policy for Sustainable Forests. Washington State Department of Natural Resources, Olympia, Washington.

Washington Department of Natural Resources. 2006b. Final Environmental Impact Statement on the Policy for Sustainable Forests. Washington Department of Natural Resources, Olympia, Washington.

Washington Department of Natural Resources. 2006c. Implementation Procedures for the Habitat Conservation Plan Riparian Forest Restoration Strategy. Olympia, Washington.

Washington Department of Natural Resources. 2007a. Addendum to a Final Environmental Impact Statement (SEPA File #02-022201). Washington Department of Natural Resources, Olympia, Washington.

Washington Department of Natural Resources. 2008. Final Environmental Impact Statement on the Policy for Sustainable Forests. Washington Department of Natural Resources, Olympia, Washington.

Washington Department of Natural Resources. 2010. South Puget HCP Planning Unit Forest Land Plan Final Environmental Impact Statement. Washington Department of Natural Resources, Olympia, Washington.

Washington Department of Natural Resources. 2013. Olympic Experimental State Forest HCP Planning Unit Forest Land Plan Revised Draft Environmental Impact Statement. Washington Department of Natural Resources, Olympia, Washington.

Washington Department of Natural Resources. 2016. Olympic Experimental State Forest HCP Planning Unit Forest Land Plan Final Environmental Impact Statement. Washington Department of Natural Resources, Olympia, Washington.


Washington Department of Natural Resources. 2019. Final Environmental Impact Statement on Alternatives for the Establishment of a Sustainable Harvest Level for Forested State Trust Lands in Western Washington. Washington Department of Natural Resources, Olympia, Washington.


Appendix 1: Stand Development Stage Definitions in DNR 2004


Stag	ges				Stand	-level Variable and As	ssociated	Thresholo	d Valu	e		
							Mai	nagement	Activ	ity		
Summarized	Detailed		QMD	Canopy Layer	RD	Stand Age	BioThin Age	Years Since BioThin	Thin Age	Years Since Thin	Snag Ratio1	CWD
Ecosystem Initiation	Ecosystem Initiation		<2									
	Sapling Exclusion		>=2									
	Pole		>5									
	Exclusion	or							>0	>=0		
Competitive	Large Tree		>11									
Exclusion	Exclusion	or	>11						>0	>=0		
			>=2	>1								
	Understory	or	>=2		>=MaxRD							
	Development		>=2			>MaxRD Age	ļ					
		or	>=2				>0	>=0				
			>=2	>1								
		or	>=2	>1		>=MaxRD Age+60	-	_				
		or	>=2	>1			>0	>=0				
		or	>=2	>1	>=MaxRD							
	Botanically	or	>=2		>=MaxRD	>=MaxRD Age+60						
	Diverse	or	>=2		>=MaxRD	. MDD A00	>0 >0	>=0 >=0				
	-	or	>=2			>=MaxRD Age+60		_				
	-	or	>=2	. 4		. Marrido Anna	>0	>5				
	-	or	>=2 >=2	>1		>MaxRD Age >=MaxRD Age+60	-					
	-	or	>=2			>MaxRD Age+00	>0	>5				
		OI	>=2	>1		>=MaxRD Age+80	-0	-3			>0.07	>2400
	-	or	>=2	>1		>=MaxRD Age+80 >=MaxRD Age+80	>0	>0			>0.07	>2400
	-	or	>=2	>1		>=IVIAXRD Age+ou	>0	>5				_
	-	or	>=2	>1	>=MaxRD	>=MaxRD Age+80	>0	>5			>0.07	>2400
	-	or	>=2		>=MaxRD	>=MaxRD Age+80	>0	>0			20.07	~2400
	Niche	or	>=2		>=MaxRD	Z-INIAXIND Age+00	>0	>5				
Structually	Diveris-	or	>=2		>-IVIANED	>=MaxRD Age+80	-0	-3			>0.07	>2400
Complex	ification	or	>=2			>=MaxRD Age+80	>0	>0			20.01	22400
		or	>=2			>MaxRD Age	>0	>5				
		or	>=2			>=MaxRD Age+80	>0	>=0			>0.07	>2400
	l	or	>=2			>=MaxRD Age+80	>0	>0				
	l	or	>=2				>0	>5			>0.07	>2400
			>=2	>1		>=MaxRD Age+160					>0.07	>2400
	l	or	>=2	>1		>=MaxRD Age+160	>0	>0				
	l	or	>=2	>1			>0	>40		<u> </u>		
	l	or	>=2		>=MaxRD	>=MaxRD Age+160					>0.07	>2400
		or	>=2		>=MaxRD	>=MaxRD Age+160	>0	>0				
	Fully	or	>=2		>=MaxRD		>0	>40				
	Functional	or	>=2			>=MaxRD Age+160					>0.07	>2400
	[or	>=2			>=MaxRD Age+160	>0	>0				
	ļ	or	>=2			>MaxRD Age	>0	>40				
	ļ	or	>=2			>=MaxRD Age+160	>0	>=0			>0.07	>2400
		or	>=2			>=MaxRD Age+160	>0	>0				
		or	>=2				>0	>40			>0.07	>2400


Appendix 2: Maps of Stands to Attain Older-forest Conditions in 2021


Older-forest stands and conservation areas (conservation areas identified on the maps as westside forest cover) represent stands that have been set aside to meet older-forest targets.


















Appendix 3: Maps of Projected Older-Forest Stands in 2100


Older-forest stands and conservation areas (conservation areas identified on the maps as westside forest cover) represent stands that have been set aside to meet older-forest targets.














Appendix 4: September 3, 2024 Revisions to May 7, 2024 Landscape Assessment

Misidentification of 3,475 acres in Columbia Planning Unit

DNR has corrected an error with the data query used in the 2024 Landscape Assessment to delineate conservation areas that are permanently deferred from variable retention harvest (VRH, a type of stand-replacement harvest) and to quantify older forests that will contribute to older forest stand structure targets over time. The error was isolated to portions of the Columbia HCP Planning Unit.

The data query to locate permanently VRH-deferred conservation areas within each HCP planning unit included several types of conservation areas, such as areas permanently conserved for marbled murrelets. Murrelet conservation areas often overlap with northern spotted owl (NSO) conservation areas.

While murrelet conservation areas are permanently deferred from VRH, NSO habitat is not deferred unless it overlaps a murrelet conservation area or other permanently deferred area. NSO habitat that does not overlap these areas may be available for harvest in planning units that have exceeded threshold habitat amounts set by the Northern Spotted Owl Conservation Strategy in the HCP. As noted in the main body of this document, DNR removed such NSO habitat from the query.

The query correctly identified murrelet conservation areas in all HCP planning units except for part of the Columbia HCP Planning Unit. In this unit, the query erroneously identified high-quality NSO habitat as permanently conserved murrelet habitat, even though a large portion of the Columbia Unit is outside of the range of the marbled murrelet and therefore does not include marbled murrelet conservation areas. Having been identified in this query, these acres were included in the May 7 calculation of permanently VRH-deferred conserved acreage that will contribute to older forest stand structure targets over time.

As a result of this error, approximately 3,475 acres within the Columbia Unit were miscategorized as permanently VRH-deferred conservation areas. Therefore, projections of older forest stand structure over time were inaccurate in that planning unit. Now that this error has been corrected, the Columbia Unit is projected to meet the 10 percent older-forest target by the year 2100, rather than by year 2090 as previously published.

Processing Revision

The spatial processing used in the analysis was streamlined, thereby reducing accumulated rounding errors in the geometric calculations. The rounding errors previously yielded inconsistent total acreages across different years of the analysis. DNR also included any boundary adjustments over the last six months as old-growth forest has been field verified and acres have been added to Natural Area Preserves and Natural Resource Conservation Areas. Running the query again resulted in minor (.1 to .2 percent) changes to HCP planning units at different decades. None of these results changed the timing estimate of when the HCP planning units (other than Columbia) are expected to meet the 10 percent older-forest target.

Data Table Revisions

Table 2 Revisions

Following is a comparison of the percent area of western Washington HCP planning units with older-forest stands in conservation areas by decade through 2120, as estimated in May and September. DNR adjusted the query output with plot discounts and a disturbance factor. Cells in the September 3 table with changes from the May 7 table are highlighted.

May 7, 2024 version of Table 2

HCP Planning		Year													
Unit	2021	2030	2040	2050	2060	2070	2080	2090	2100	2110	2120				
Columbia	1.1%	1.2%	1.4%	1.8%	2.6%	4.3%	6.8%	10.1%	14.0%	17.3%	18.9%				
North Puget	3.2%	3.9%	4.9%	6.2%	7.9%	10.2%	13.2%	16.7%	20.6%	23.9%	25.0%				
OESF	10.2%	10.7%	11.0%	11.7%	12.6%	13.9%	16.0%	20.1%	25.0%	28.4%	29.6%				
South Coast	0.2%	0.3%	0.6%	1.2%	2.2%	3.6%	6.0%	8.8%	12.3%	16.0%	18.7%				
South Puget	1.7%	2.1%	2.7%	3.6%	4.6%	6.1%	8.4%	11.3%	14.4%	17.2%	18.7%				
Straits	1.8%	2.5%	3.2%	4.3%	5.6%	7.4%	9.9%	12.6%	15.0%	17.9%	19.3%				

September 3, 2024 version of Table 2

HCP Planning	Year												
Unit	2021	2030	2040	2050	2060	2070	2080	2090	2100	2110	2120		
Columbia	<mark>1.0%</mark>	1.2%	1.4%	<mark>1.7%</mark>	<mark>2.4%</mark>	<mark>3.9%</mark>	<mark>6.2%</mark>	<mark>9.4%</mark>	<mark>13.3%</mark>	<mark>16.5%</mark>	<mark>18.2%</mark>		
North Puget	3.2%	3.9%	4.9%	6.2%	7.9%	10.2%	13.2%	16.7%	<mark>20.5%</mark>	23.9%	25.0%		
OESF	10.2%	10.7%	11.0%	11.7%	12.6%	13.9%	<mark>15.9%</mark>	20.0%	<mark>24.9%</mark>	28.3%	<mark>29.5%</mark>		
South Coast	0.2%	0.3%	0.6%	1.2%	<mark>2.1%</mark>	3.6%	<mark>5.9%</mark>	8.8%	<mark>12.2%</mark>	<mark>15.9%</mark>	<mark>18.6%</mark>		
South Puget	1.7%	<mark>2.2%</mark>	2.7%	3.6%	4.6%	6.1%	8.4%	11.3%	14.4%	<mark>17.1%</mark>	18.7%		
Straits	<mark>1.9%</mark>	<mark>2.6%</mark>	3.2%	4.3%	5.6%	7.4%	9.9%	12.6%	<mark>15.1%</mark>	<mark>18.0%</mark>	<mark>19.5%</mark>		

Table 3 Revisions

Following is a comparison of the percent area of Western Washington HCP planning units with older-forest stands in conservation areas by decade through 2120, as estimated in May and September. Values over 10 percent are shown in bold. This is raw query output without plot discount or a disturbance factor applied. Cells in the September 3 table with changes from the May 7 table are highlighted.

May 7, 2024 version of Table 3

HCP Planning	Year													
Unit	2021	2030	2040	2050	2060	2070	2080	2090	2100	2110	2120			
Columbia	1.5%	1.7%	2.0%	2.6%	3.9%	6.5%	10.5%	15.8%	22.1%	27.8%	31.1%			
North Puget	4.7%	5.8%	7.2%	9.2%	11.9%	15.6%	20.3%	26.1%	32.5%	38.5%	41.0%			
OESF	14.9%	15.7%	16.3%	17.5%	18.9%	21.2%	24.5%	31.3%	39.6%	45.6%	48.6%			
South Coast	0.3%	0.5%	1.0%	1.8%	3.2%	5.5%	9.2%	13.8%	19.5%	25.8%	30.7%			
South Puget	2.5%	3.2%	4.0%	5.3%	6.9%	9.2%	12.9%	17.7%	22.8%	27.6%	30.7%			
Straits	2.7%	3.7%	4.8%	6.3%	8.4%	11.2%	15.2%	19.6%	23.8%	28.8%	31.7%			

September 3, 2024 version of Table 3

HCP Planning		Year												
Unit	2021	2030	2040	2050	2060	2070	2080	2090	2100	2110	2120			
Columbia	1.5%	1.7%	2.0%	2.6%	<mark>3.7%</mark>	<mark>5.9%</mark>	<mark>9.5%</mark>	<mark>14.7%</mark>	<mark>21.0%</mark>	<mark>26.6%</mark>	<mark>29.9%</mark>			
North Puget	4.7%	5.8%	7.2%	9.2%	11.9%	<mark>15.5%</mark>	20.3%	<mark>26.0%</mark>	32.5%	<mark>38.4%</mark>	<mark>40.9%</mark>			
OESF	14.9%	15.7%	16.3%	17.5%	<mark>19.0%</mark>	21.2%	24.5%	<mark>31.2%</mark>	<mark>39.4%</mark>	<mark>45.5%</mark>	<mark>48.4%</mark>			
South Coast	0.3%	0.5%	1.0%	1.8%	3.2%	<mark>5.4%</mark>	9.1%	13.7%	<mark>19.4%</mark>	<mark>25.6%</mark>	<mark>30.5%</mark>			
South Puget	2.5%	3.2%	4.0%	5.3%	6.9%	9.2%	<mark>12.8%</mark>	<mark>17.6%</mark>	22.8%	27.6%	<mark>30.6%</mark>			
Straits	2.7%	<mark>3.8%</mark>	4.8%	<mark>6.4%</mark>	<mark>8.5%</mark>	11.2%	<mark>15.3%</mark>	<mark>19.7%</mark>	<mark>23.9%</mark>	<mark>29.0%</mark>	<mark>32.0%</mark>			

September 3, 2024 Data Tables

For convenience, both tables updated on September 3 are shown below. These tables are identical to the September 3 tables presented in the previous section.

Table 2: Percent area western Washington HCP planning units with older-forest stands in conservation areas by decade through 2120. Adjusted query output with plot discounts and disturbance factor.

HCP Planning		Year												
Unit	2021	2030	2040	2050	2060	2070	2080	2090	2100	2110	2120			
Columbia	1.0%	1.2%	1.4%	1.7%	2.4%	3.9%	6.2%	9.4%	13.3%	16.5%	18.2%			
North Puget	3.2%	3.9%	4.9%	6.2%	7.9%	10.2%	13.2%	16.7%	20.5%	23.9%	25.0%			
OESF	10.2%	10.7%	11.0%	11.7%	12.6%	13.9%	15.9%	20.0%	24.9%	28.3%	29.5%			
South Coast	0.2%	0.3%	0.6%	1.2%	2.1%	3.6%	5.9%	8.8%	12.2%	15.9%	18.6%			
South Puget	1.7%	2.2%	2.7%	3.6%	4.6%	6.1%	8.4%	11.3%	14.4%	17.1%	18.7%			
Straits	1.9%	2.6%	3.2%	4.3%	5.6%	7.4%	9.9%	12.6%	15.1%	18.0%	19.5%			

Table 3: Percent area Western Washington HCP planning units with older-forest stands in conservation areas by decade through 2120. Values over 10% in bold. Raw query output without plot discount or disturbance factor.

HCP Planning		Year												
Unit	2021	2030	2040	2050	2060	2070	2080	2090	2100	2110	2120			
Columbia	1.5%	1.7%	2.0%	2.6%	3.7%	5.9%	9.5%	14.7%	21.0%	26.6%	29.9%			
North Puget	4.7%	5.8%	7.2%	9.2%	11.9%	15.5%	20.3%	26.0%	32.5%	38.4%	40.9%			
OESF	14.9%	15.7%	16.3%	17.5%	19.0%	21.2%	24.5%	31.2%	39.4%	45.5%	48.4%			
South Coast	0.3%	0.5%	1.0%	1.8%	3.2%	5.4%	9.1%	13.7%	19.4%	25.6%	30.5%			
South Puget	2.5%	3.2%	4.0%	5.3%	6.9%	9.2%	12.8%	17.6%	22.8%	27.6%	30.6%			
Straits	2.7%	3.8%	4.8%	6.4%	8.5%	11.2%	15.3%	19.7%	23.9%	29.0%	32.0%			